

Homework 4

Lecturer: Ronitt Rubinfeld

Due Date: April 23, 2006

1. Given a boolean function $f(\cdot)$ on boolean inputs, a sequence $C = C_1, C_2, \dots$ of circuits is a *circuit family for $f(\cdot)$* if C_n has n inputs and computes $f(x_1, \dots, x_n)$ at its output for all n bit inputs (x_1, \dots, x_n) . The family C is said to be *polynomial-sized* if the size of C_n is bounded above by $p(n)$ for every n , where $p(\cdot)$ is a polynomial. A *randomized circuit family for $f(\cdot)$* is a circuit family for $f(\cdot)$ that, in addition to the n inputs x_1, \dots, x_n , takes m inputs r_1, \dots, r_m , each of which is equiprobably and independently 0 or 1. In addition, for every n , circuit C_n must satisfy

- (a) if $f(x_1, \dots, x_n) = 0$ then output 0 regardless of the values of the random inputs r_1, \dots, r_m .
- (b) if $f(x_1, \dots, x_n) = 1$ then output 1 with probability $\geq 1/2$.

Show: If a boolean function has a randomized polynomial sized circuit family, then it has a polynomial sized circuit family.

2. You are given a 2-SAT formula $\phi(x_1, \dots, x_n)$. Consider the following algorithm for finding a satisfying assignment:
- Start with an arbitrary assignment. If it's satisfying, output it and halt.
 - Do s times:
 - Pick an arbitrary unsatisfied clause
 - Pick one of the two literals in it uniformly at random
 - Complement the setting of the chosen literal
 - If the new assignment satisfies ϕ , output the assignment and halt.

Show that if you pick s to be $O(n^2)$, you will output a satisfying assignment with probability at least $3/4$.

3. Let $G(V, E)$ be a graph with n vertices such that for some constant $\alpha > 0$, and every set $S \subseteq V$ with $n/2$ vertices,

$$|\{w \in V \mid \exists v \in S, (v, w) \in E\}| \geq \frac{n}{2} + \alpha n.$$

For any positive integer k , let W_1, \dots, W_k be subsets of V of size at least $(1 - \alpha)n$ each. Show that there exists a path (v_1, \dots, v_k) in G such that for $1 \leq i \leq k$, $v_i \in W_i$.

4. A d -regular graph has (K, A) -vertex expansion if $\forall S \subset V, |S| \leq K, |\lambda(S)| \geq A|S|$ where $\lambda(S) = |\{u \mid \exists v \in S \text{ s.t. } (u, v) \in E\}|$. For a probability distribution π over $[n]$, the *collision probability* is

$$\|\pi\|^2 = \sum_x \pi_x^2$$

Show that the following is true:

- $\|\pi\|^2 \geq \frac{1}{|S(\pi)|}$ where $S(\pi) = \{x | \pi_x > 0\}$.
- $\|\pi\|^2 = \|\pi - u\|^2 + \frac{1}{n}$ where u is the uniform distribution.
- If there is a constant $\lambda < 1$ such that the transition matrix of G is such that $|\lambda_2| \leq \lambda$ then for any $\alpha < 1$, G has vertex expansion $(\alpha n, \frac{1}{(1-\alpha)\lambda^2 + \alpha})$.

Remark: We use the convention that $|\lambda_1| \geq |\lambda_2| \geq \dots \geq |\lambda_n|$, where $\lambda_1, \dots, \lambda_n$ are eigenvalues of the random walk matrix.

5. (optional) Let $G = (V, E)$ be a d -regular, α -expander, i.e., $\forall S \subset V, |S| \leq |V|/2, |\lambda(S) \setminus S| \geq \alpha|S|$ (λ is defined as above). Let $n = |V|$. Suppose a set of pairs $S = \{(a_1, b_1), \dots, (a_q, b_q)\}$ are such that a_i and b_i are chosen uniformly and independently from V for all i . Show that for all a, b , there is a path connecting a, b of length $O(\log n)$. Show that there exists a way of connecting each a_i, b_i pair via a path of length $O(\log n)$, such that no edge is used more than $O(\log n)$ times in total over all paths.

Hint: for each (a_i, b_i) pair, choose a random x_i . Show how to pick a “random” path from a_i to x_i and from b_i to x_i . Show that your method is such that no edge is used more than $O(\log n)$ times in total.