

Impossible Shaded Images

Berthold K. P. Horn
MIT AI Laboratory
545 Technology Square
Cambridge, MA 02139

Richard S. Szeliski
Digital Equipment Corporation
Cambridge Research Lab
One Kendall Square, Building 700
Cambridge, MA 02139

Alan L. Yuille
Division of Applied Sciences
Harvard University
Cambridge, MA 02138

Abstract: *In this paper, we show that there exist images that could not have arisen from shading on a smooth surface with uniform reflecting properties and lighting. Much work has been done on recovering surface shape from images and there has been some attention paid to the question of uniqueness. It has been shown, for example, that singular points and occluding boundaries often curtail ambiguity. But little has been said about the existence of solutions, perhaps because in practice the given image is assumed to have arisen from a uniform, smoothly curved surface, and so one knows that there must be at least one solution. What if, however, the reflecting properties of the surface vary from place to place? What if the actual surface does not reflect light the way one has assumed, or the light source is not where it was thought to be? Will the solution only be warped by these departures from the ideal model, or may there in fact be situations where there is no smooth surface that could have given rise to the given shading pattern? Can the fact that a shaded image of some surface with spatially varying surface reflectance is impossible in this sense be used to detect surface albedo variations?*

1 Introduction

The problem of *shape from shading* has a history almost as long as that of computer vision itself [13]. Aside from the development of algorithms for recovering shape from shaded images, some attention has been paid to the problem of uniqueness of the solution. It has been shown that *singular points* of brightness in the image (corresponding to isolated global extrema in the reflectance map) play an important role in limiting the number of possible solution surfaces [4] [5] [3] [2] [19].

So far little has been said, however, about the existence of solutions (but see [19]). Surfaces with continuously varying surface orientation give rise to shaded images. Are there brightness patterns that could not have arisen this way? Can such *impossible shaded images* be detected directly from their brightness patterns without explicitly solving the shape from shading equations? We show here that this is indeed the case.¹. In this paper we will assume that the distribution of light sources and the reflecting properties of the surface are known and that the reflecting properties of the surface are uniform. We also assume that the surfaces are *smooth*, by which we will mean that they have continuous first derivatives.

2 Reflectance Map and Image Irradiance Equation

The brightness at a point in the image, the *image irradiance* is proportional to the brightness of the corresponding point on an object, the *scene radiance*[11]. The latter depends on: (a) the reflecting properties of the surface material, (b) the distribution and intensity of the light sources, and (c) the surface orientation.

Surface orientation has two degrees of freedom and can be specified in several different ways. The slopes $p = (\partial z / \partial x)$ and $q = (\partial z / \partial y)$ in two orthogonal directions are convenient for this purpose, where $z(x, y)$ is the height of the surface above some reference plane perpendicular to the direction of projection of orthographic image formation². Surface orientation may also be specified by means of the unit normal, which can be obtained from the components of the gradient, p and q , as follows: $\hat{n} = ((-p, -q, 1)^T / \sqrt{1 + p^2 + q^2})$.

The *reflectance map* $R(p, q)$ gives scene radiance as a function of surface orientation

¹This work arose from a conjecture by one of us. Following initial confirmation of the conjecture by the co-authors and others (see acknowledgments), we received note of work by Oliensis [17] that demonstrates that small fluctuations in intensities could indeed make most surfaces “impossible.”

²The shape-from-shading problem can be formulated in the case where the imaging system performs a perspective projection [18], [7], and when the light sources are near the objects being viewed, but this makes the analysis harder, since scene radiance then depends on position as well as surface orientation.

and encodes information about both the surface reflecting properties and light source distribution [10]. It can be computed given the *bi-directional reflectance distribution function* (BRDF) [14] [11] or determined experimentally using the image of a calibration object.

The (normalized) *image irradiance equation* is $E(x, y) = R(p(x, y), q(x, y))$, where $E(x, y)$ is the image irradiance at the point (x, y) in the image, while $p(x, y)$ and $q(x, y)$ are the partial derivatives of $z(x, y)$ at the corresponding point on an object in the scene [10], [11].

The shape-from-shading problem is that of recovering the surface $z(x, y)$ given the image $E(x, y)$ and the reflectance map $R(p, q)$. The image irradiance equation can be viewed as a first-order non-linear partial differential equation, and so can be solved using the method of characteristic strips [7], [9], [11].

2.1 Phenomenological Models of Reflection

Some of the impossible images we will present depend on particular properties of a class of reflectance maps, such as rotational symmetry. At other times it is useful to have a very specific reflectance map in mind, such as that of a Lambertian surface under point source illumination. Let us consider a simple imaging situation where we are dealing with an idealized surface material that satisfies two conditions: (a) it appears equally bright from all viewing directions, and (b) it reflects all incident light. Such a surface is called an (ideal) Lambertian surface and it can be shown that when illuminated by a single light source it satisfies Lambert's cosine law [11]. In this case brightness depends on the cosine of the incident angle, the angle between the incident rays and the surface normal, and is independent of the direction towards the viewer. If there is a single light source in the direction given by the unit vector $\hat{\mathbf{s}} = ((-p_s, -q_s, 1)^T / \sqrt{1 + p_s^2 + q_s^2})$, then we use the fact that the cosine of the incident angle is equal to $(\hat{\mathbf{n}} \cdot \hat{\mathbf{s}})$ and so obtain the rotationally symmetric (normalized) reflectance map

$$R(p, q) = \max \left[0, \frac{1 + p_s p + q_s q}{\sqrt{1 + p^2 + q^2} \sqrt{1 + p_s^2 + q_s^2}} \right].$$

A particularly simple case arises when the light source lies in the same direction as the viewer, that is, when $(p_s, q_s) = (0, 0)$, for then $R(p, q) = (1 / \sqrt{1 + p^2 + q^2})$. This special case can be used when a concrete example of a rotationally symmetric reflectance map is needed with brightness decreasing monotonically with surface slope s , where $s = \sqrt{p^2 + q^2}$.

3 Impossible Shaded Images

3.1 A Circularly Symmetric Dark Blotch

Suppose we are given the image

$$E(x, y) = \begin{cases} 1/\sqrt{1 + \cos^2 r}, & \text{for } r < \pi/2; \\ 1, & \text{for } r \geq \pi/2; \end{cases}$$

where $r = \sqrt{x^2 + y^2}$ (shown in Figure 1), with reflectance map

$$R(p, q) = \frac{1}{\sqrt{1 + p^2 + q^2}}.$$

Then there is a “solution”

$$z(x, y) = \begin{cases} \sin r; & \text{for } r < \pi/2; \\ 1; & \text{for } r \geq \pi/2. \end{cases}$$

While this function is smooth almost everywhere it has a conical singularity at the origin. Does there exist a solution that has continuous first derivatives everywhere? The answer is “no,” as we show next (generalizing to an even wider class of impossible images).

3.2 Compact Dark Blotch on Unit Brightness Background

Suppose we are told that the reflectance map has a unique isolated global maximum of one at the origin. That is,

$$R(p, q) \begin{cases} = 1, & \text{for } (p, q) = (0, 0); \\ < 1, & \text{otherwise.} \end{cases}$$

In this case a surface facing the viewer directly has brightness one, and surface patches oriented differently are always darker (as would be the case if the surface was a Lambertian reflector with the light source in the direction towards the viewer). Suppose that image brightness is less than one in some simply connected compact region D , while brightness equals one outside this region, that is,

$$E(x, y) \begin{cases} < 1, & \text{for } (x, y) \in D; \\ = 1, & \text{otherwise.} \end{cases}$$

Note that outside the region D , the surface gradient (p, q) must be zero, since that is the only gradient for which $R(p, q) = 1$. If $p = 0$ and $q = 0$ in some connected region,

Figure 1. (a) Cross-section of rotationally symmetric shaded image, (circular) shown in (b), that could not have arisen from a (smooth) surface with continuous first derivatives, if we assume that the reflectance map is $R(p, q) = 1/\sqrt{1 + p^2 + q^2}$.

then surface height $z(x, y)$ must be constant in that region. Now, either the surface $z(x, y)$ is constant in the region D , or it has at least one extremum there. But the surface cannot be constant in D , because that would imply that the brightness there equaled one; so it must have an extremum. The first partial derivatives must vanish at that extremum since we are assuming that the surface $z(x, y)$ has continuous first derivatives. But this implies that the brightness at the extremum must be one, which contradicts the assumption that brightness is less than one everywhere in the region D . Thus there is no surface with continuous first derivatives that will give rise to the given image.

We note that a compact dark blotch on a unity brightness background needs to have at least one interior point where brightness equals one if it is to be the shaded image of some surface with continuous first derivatives.

3.3 Source not at Viewer

It is possible to extend the result of the previous section to reflectance maps that have their unique isolated global extremum somewhere other than the origin in gradient space. Suppose the extremum in the reflectance map occurs at $(p, q) = (p_s, q_s)$, as happens with a Lambertian surface when the light source is away from the viewer.

Then the surface orientation outside the region D is fixed and the surface there planar, with surface normal

$$\hat{\mathbf{n}} = \frac{(-p_s, -q_s, 1)^T}{\sqrt{1 + p_s^2 + q_s^2}}.$$

We can then use a similar argument to that in the previous section, by using a new coordinate system oriented with the z -axis parallel to $\hat{\mathbf{n}}$. The only potential problem arises from the possibility that the point where the surface exhibits an extremum in the light-source coordinate system may be obscured by another portion of the surface when seen from the viewing direction. But this can only occur if there is a fold in the surface, as seen from the viewing direction, and the derivatives of surface height are discontinuous at the fold.

3.4 Singular Points are not Required

The above examples might appear to suggest that singular points are crucial to the construction of impossible shaded image examples, since the exterior of the region D consists entirely of singular points, where surface orientation can in fact be recovered locally. But this is not so. Consider, for example, a reflectance map that has brightness one for zero slope, with brightness falling off with slope in such a way that one can write

$$f(s) \leq R(p, q) \leq g(s) \quad \text{where} \quad s = \sqrt{p^2 + q^2}$$

for two monotonically decreasing function $f(s)$ and $g(s)$ (see Figure 2).

A special case of this is a rotationally symmetric reflectance map with brightness dropping monotonically with slope. A particular instance of this special case is a Lambertian surface with the source at the viewer, as mentioned above.

Now suppose that there is a compact, simply connected region D in which the brightness is low, nested inside another compact simply connected region F , outside of which brightness is high (see Figure 3).

In the part of F that is not in D , brightness makes a smooth transition. That is

$$\begin{cases} 0 < E(x, y) < E_i; & \text{for } (x, y) \in D, \\ E_i \leq E(x, y) \leq E_o; & \text{for } (x, y) \in F - D, \\ E_o < E(x, y) < 1; & \text{otherwise.} \end{cases}$$

There are no singular points, since $E(x, y) < 1$ everywhere, yet, as we show next, we can chose E_i and E_o such that there is no smooth surface giving rise to this shading.

We have for the slope of the surface inside the region D : $s \geq s_i = f^{-1}(E_i)$ while the slope outside F satisfies $s \leq s_o = g^{-1}(E_o)$. Now inside the region F , the slope is guaranteed to be non-zero, so the surface $z(x, y)$ has a unique non-zero gradient at

Figure 2. In this generalization, the reflectance map need not be radially symmetric, but must be bounded above and below by two radially symmetric, monotonically decreasing functions.

Figure 3. For the impossible image discussed here, brightness is low inside the region D , high outside the region F , with a smooth transition in the part of F that is not also in D . A curve of steepest descent can be constructed that passes through the center of the largest circle that can be inscribed in D . A contradiction is indicated if the change in height along this curve in the region D is greater than the change of height along the perimeter of region F .

every point. This gradient field can be integrated out to yield lines of steepest ascent on the surface. Such steepest ascent curves cannot cross or terminate in F , and so can be followed all the way from one point on the boundary of F to another. Such a steepest ascent curve passing through a point in D , will similarly cross the boundary of D in two places.

Suppose that w is the diameter of the largest inscribed circle of the region D . Then the steepest ascent curve passing through the center of this circle must have length at least w , and hence a change in z from one end to the other of at least $w s_i$. Now suppose that p is the perimeter of the region F . Then the shortest distance along the boundary of F between the two points where this steepest ascent curve touches the boundary is at most $p/2$, and so the change in z is at most $(p/2)s_o$. Given w and p , we can now choose E_i and E_o , and hence s_i and s_o , such that $w s_i > (p/2)s_o$, which leads to a contradiction.

(Note: the only reason that two nested regions are needed in this construction is to allow brightness to vary smoothly in the transition, as it must, since we have assumed that the first derivatives and hence brightness is continuous.)

3.5 Nested Iso-Brightness Contours

The above construction can be extended to nested iso-brightness contours of monotonically increasing brightness from the inside to the outside.

Suppose that we have a dark blotch in the image that increases monotonically in brightness from the inside outward, so that one can construct a set of nested iso-brightness contours for brightnesses $0 < E_0 < E_1 < E_2 \dots < E_n < 1$, where E_0 is the brightness of the darkest point in the image. Suppose that the minimum distance between the two iso-brightness contours for $E = E_i$ and $E = E_{i+1}$ is w_i . Note that the slope of the surface on points lying between these two iso-brightness contours is constrained by $s > s_i = f^{-1}(E_{i+1})$. Consider the curve of steepest ascent passing through the point where $E = E_0$. The change in height along this contour between the points where it crosses the iso-brightness contour $E = E_n$ is bounded below by $\delta z_n > 2 \sum_{i=0}^{n-1} s_i w_i$ while at the same time bounded above by $\delta z_n \leq (p_n/2)\bar{s}_n$, where p_n is the length of the iso-brightness contour $E = E_n$ and $\bar{s}_n = g^{-1}(E_n)$. We have an impossible image unless $2 \sum_{i=0}^{n-1} s_i w_i \leq (p_n/2)\bar{s}_n$ for all n . This provides a way of constructing a variety of impossible images. It also provides a limit on how dark a blotch on a bright background can be before it can no longer be interpreted as shading on an inclined portion of the surface. These topics are explored further in [15].

3.6 Folds in Riemann Sheets on Gaussian Sphere

Imagine that we have a rotationally symmetric reflectance map that drops to zero at infinity in gradient space, such as the Lambertian surface with the source at the viewer. For what we will do next, it is convenient to think of the reflectance map as a function of position on the Gaussian sphere, rather than as a function of the components of the gradient. The reflectance map plotted on the Gaussian sphere here has a peak of one at the “pole,” corresponding to the viewing direction, and drops off to zero at the “equator,” corresponding to points on the occluding boundary of the object being viewed³.

Now suppose that we are given an image that has non-zero brightness in the interior of some compact simply connected region D , with zero brightness on the boundary ∂D of this region. Then the boundary ∂D is a *silhouette*, that is, the projection of an occluding contour on the object being viewed. If we assume that surface orientation varies continuously, there is a mapping from the object’s surface to the surface of the Gaussian sphere that covers every point in one hemisphere (at least once). We can see this by noting that for any orientation in the hemisphere, there must be a point on the surface with that orientation since a plane with that orientation as its normal approaching from infinity will touch the surface somewhere [6] (If the object is convex, this *Gauss map* is invertible). Note that the occluding boundary maps onto the “equator” of the hemisphere.

It should, first of all, be clear that there must be at least one point in the image where the brightness equals the maximum brightness in the reflectance map, since the pole of the hemisphere must be covered. How many such extrema can there be in the image of a single object with continuous first derivatives? We can have more than one if the object is not convex, since the mapping from the surface onto the sphere then folds over on itself. But every time we fold it over in order to cover the pole more than once, we add *two* new places where the surface is oriented for maximal reflection of light. This suggests that there must be an odd number of bright spots in the image. There is one exception to this rule: If the fold on the Gaussian sphere happens to cross the pole it will yield only one maximum instead of two. Of course, in this special case, any slight change in the orientation of the object with respect to the viewer will change this. So this does not apply if we assume that the viewer is in “general position.”

The other possibility is that the part of the surface carrying one of the two points happens to be obscured by another part of the object. But in this case the surface $z(x, y)$ is not a continuous function of x and y within in the region D . Furthermore, the occluding boundary is not a simply closed curve, and parts of the occluding boundary

³The Gaussian sphere approach to the analysis of shading was introduced in [16].

lie within D . But this is impossible, since the brightness is non-zero inside D .

3.7 Multiple Viewpoints or Multiple Lighting Conditions

So far we have been trying to determine the “impossibility” of surfaces from a *single* image. This problem (naturally) becomes a lot easier if we have several images of the surface corresponding to different lighting conditions or different viewpoints. We can, for example, make use of certain photometric invariants [16] [20] that relate properties of the surface geometry to properties of the image brightness, assuming only a generic form for the reflectance map $R(p, q)$ and constant albedo. One specific result [20], extending a result of [16], is that the directions of the isophotes (the lines of constant image brightness) must always lie along the directions of principal curvature at parabolic lines (lines of zero Gaussian curvature), hence these isophote directions will be invariant as we alter the viewing conditions - this has been exploited by [2]. Moreover for generic surfaces these are the only lines with this property. Thus, given several images, we can use these results to determine the parabolic lines of the surface. For regular (that is, not “impossible”) surfaces the parabolic lines will either be closed contours or will terminate at the boundaries of the viewed object. If we find parabolic lines that terminate inside the object, or have other undesirable behavior, then we have an “impossible” surface.

3.8 Iterative Solution Applied to Impossible Image

It is of interest to see what the iterative algorithm [12] will do when presented with an impossible shaded image. It is shown, [15], that it finds the “solution”

$$z(x, y) = \begin{cases} \sin r; & \text{for } r < \pi/2; \\ 1; & \text{for } r \geq \pi/2, \end{cases}$$

This function is smooth everywhere except at the origin, where it has a conical singularity.

4 Detecting Spatial Variations of Albedo

If a surface has spatially varying reflecting properties, or if the illumination has spatial variations, then the normal shading rules for the image are altered. An extreme example of this is a photographic print, where all the brightness variations are due to spatially varying reflectance, and there is no shading resulting from spatial variations in

surface orientation. Given the limited information in a single image, it often is not possible to separate the contributions to the brightness pattern that come from spatially varying surface orientation and those that come from spatial variations in reflectance or illumination.

As we have demonstrated in this paper, however, it is sometimes possible to show that the given image could not have arisen from a uniformly illuminated smooth surface with uniform reflecting properties (that is, it is an impossible shaded image). The way this manifests itself when iterative algorithms are used for recovering the surface shape is that the functional cannot be reduced to zero, and that discontinuities and cone shaped singularities in surface orientation remain in the estimated solution.

When we look at images taken of the surface of rocky planets like Mars, we can get a clear impression of the shapes of surface features such as impact craters, yet also be aware of the fact that surface albedo varies from place to place. Our ability to separate shading and albedo variations suggest that there is some way of distinguishing the two. Quite often what distinguishes shading from surface markings is that the latter have sharp transitions between regions of relatively constant reflectance, while shading typically varies smoothly. What is needed then is a simultaneous solution of the shading and the lightness problems [8], [1]⁴. At least there now is a diagnostic test that tells us when the assumptions of uniform albedo and uniform illumination are being violated.

Similarly, if the assumed light source is in the wrong position, there will typically not be a solution to the shape from shading problem. This again manifests itself as a residual error in the iterative scheme. It has been found possible, for example, to refine an estimate of the light source position by searching for the position that minimizes the residual errors [12]

5 Conclusions

We have shown in this paper that there exist shaded images that cannot have originated from a uniformly illuminated, smooth continuous surface with uniform albedo. The typical condition where this occurs is when we have a dark area (corresponding to a region of high gradient) surrounded by a lighter region (with low gradient). For this to correspond to a real surface, we can establish that there must be a local extremum or area of lower gradient inside the dark region. This, in turn, will show up as either a light area in the image or an orientation discontinuity in the surface (thus violating either our intensity or smoothness constraints). We can also sometimes establish the impossibility of a shaded image by counting the number of extrema inside a region

⁴Simultaneous estimation of shape and albedo has been described in a recent paper [21].

corresponding to an isolated surface patch.

The theoretical arguments we have presented are in agreement with the effects observed with numerical shape from shading algorithms. When presented with an “impossible shaded image”, the algorithm will find a solution that is smooth almost everywhere but has isolated orientation singularities (“peaks” or “cusps”). We thus have two methods for detecting when the assumptions behind our shape from shading algorithm are being violated. First, we can examine the intensity image to check if any of the theoretical conditions for impossible shading exist. Second, we can monitor the output of our numerical shape from shading algorithm to see if isolated or connected singularities exist in the final solution. Detecting these violations will hopefully lead us to more robust and more general shape from shading algorithms, which can detect albedo variation and discontinuities in the reconstructed surface.

Acknowledgments

The idea for this paper grew out of a competition organized by one of the authors (BKPH), who had a conjecture that the image presented at the beginning of section 3 was indeed an impossible shaded image. Proofs of impossibility were received first from one of the co-authors (RSS), then the other (ALY), followed by Jim Smith, Richard Kozera and Mike Brooks, John Oliensis, and David Lee. We would like to thank the others, as well as Michael Gennert and Bror Saxberg, for their interest in this problem.

References

- [1] Blake, A. (1985) “Boundary Conditions for Lightness Computation in Mondrian World,” *Computer Vision, Graphics and Image Processing*, Vol. 32, No. 3, pp. 314-327, December.
- [2] Blake, A., A. Zisserman & G. Knowles (1985) “Surface Descriptions from Stereo and Shading,” *Image & Vision Computing*, Vol. 3, No. 4, pp. 183-191. Also in *Shape from Shading*, MIT Press, 1989.
- [3] Brooks, M.J. (1983) “Two Results Concerning Ambiguity in Shape from Shading,” *Proceedings of the National Conference on Artificial Intelligence*, Washington, D.C., August 22-26, pp. 36-39.
- [4] Bruss, A.R. (1982) “The Eikonal Equation: Some Results Applicable to Computer Vision,” *Journal of Mathematical Physics*, Vol. 23, No. 5, pp. 890-896, May. Also in *Shape from Shading*, MIT Press, 1989.

- [5] Deift, P. & Sylvester, J. (1981) "Some Remarks on the Shape-from-Shading Problem in Computer Vision," *Journal of Mathematical Analysis and Applications*, Vol. 84, No. 1, pp. 235-248, November.
- [6] Hilbert, D., & S. Cohn-Vossen, *Geometry and the Imagination*, Chelsea Publishing Co., New York, 1952.
- [7] Horn, B.K.P. (1970) "Shape from Shading: a Method for Obtaining the Shape of a Smooth Opaque Object from One View," Ph.D. Thesis, Department of Electrical Engineering, MIT. Also AD717336 available from National Technical Information Service.
- [8] Horn, B.K.P. (1974) "Determining Lightness from an Image," *Computer Graphics and Image Processing*, Vol. 3, No. 1, pp. 277-299, December.
- [9] Horn, B.K.P. (1975) "Obtaining Shape from Shading Information," Chapter 4 in *The Psychology of Computer Vision*, P.H. Winston (ed.), McGraw-Hill Book Company, New York, NY, pp. 115-155. Also in *Shape from Shading*, MIT Press, 1989.
- [10] Horn, B.K.P. (1977) "Understanding Image Intensities (sic)," *Artificial Intelligence*, Vol. 8, No. 2, pp. 201-231, April. Also (1987) in *Readings in Computer Vision*, Fischler, M.A. & O. Firschein (eds.), Morgan Kaufmann Publishers, Los Altos, CA, pp. 45-60.
- [11] Horn, B.K.P. (1986) *Robot Vision* (particularly Chapters 10 & 11, pp. 202-277), MIT Press, Cambridge, MA & McGraw-Hill Book Company, New York, NY.
- [12] Horn, B.K.P. (1990) "Height and Gradient from Shading," *International Journal of Computer Vision*, Vol. 5, No. 1, pp. 37-75, August. Also Memo 1105, MIT Artificial Intelligence Laboratory.
- [13] **Shape from Shading**. Eds. B.K.P. Horn and M.J. Brooks. MIT Press. Boston. 1989.
- [14] Horn, B.K.P. & R.W. Sjoberg (1979) "Calculating the Reflectance Map," *Applied Optics*, Vol. 18, No. 11, pp. 1770-1779, June. Also (1978) *Proceedings of the DARPA Image Understanding Workshop*, Carnegie-Mellon University, Pittsburgh, PA, November 14-15, pp. 115-126. Also in *Shape from Shading*, MIT Press, 1989.
- [15] Horn, B.K.P., Szeliski, R. and Yuille, A.L. (1991) "Impossible Shaded Images," Harvard Robotics Laboratory Technical Report 91-15, November.
- [16] Koenderink, J.J. & A.J. van Doorn (1980) "Photometric Invariants Related to Solid Shape," *Optica Acta*, Vol. 27, No. 7, pp. 981-996. Also in *Shape from Shading*, MIT Press, 1989.
- [17] Oliensis, J. (1989) "Existence and Uniqueness in Shape from Shading," unpublished report, Computer and Information Science, University of Massachusetts at Amherst.
- [18] Rindfleisch, T. (1966) "Photometric Method for Lunar Topography," *Photogrammetric Engineering*, Vol. 32, No. 2, pp. 262-277, March.

- [19] Saxberg, B.V.H. (1988) "A Modern Differential Geometric Approach to Shape from Shading," Ph.D. Thesis, Department of Electrical Engineering and Computer Science, MIT.
- [20] Yuille, A.L. (1989) "Zero crossings on lines of curvature," *Computer Vision, Graphics and Image Processing*, Vol. 45, pp. 68-87.
- [21] Zheng, Q. and Chellappa, R. (1991) "Estimation of Illuminant Direction, Albedo, and Shape from Shading", *IEEE Transactions on Pattern Analysis and Machine*, Vol. 13, No. 7, pp. 680-702, May.