

## Exercise 03

Lecturer: Mohsen Ghaffari

Teaching Assistant: Manuela Fischer

## 1 Deterministic Spanner Construction

Devise a deterministic algorithm that constructs a 2-additive spanner with  $\tilde{O}(n^{3/2})$  edges.

## 2 Monotone Submodular Maximization

Consider a set  $\mathcal{U}$  of  $n$  elements that we can buy and a function  $f : 2^{\mathcal{U}} \rightarrow \mathbb{R}^+$ , where for each subset  $S \subseteq \mathcal{U}$ , the value  $f(S)$  determines our profit if we buy exactly the elements of set  $S$ .

We assume two properties about this profit function: (A) Function  $f$  is monotone in the sense that  $f(S) \leq f(T)$  for any two sets  $S, T$  such that  $S \subseteq T$ , and (B) Function  $f$  is submodular in the sense that  $f(S \cup i) - f(S) \geq f(T \cup i) - f(T)$  for any  $i \in \mathcal{U}$  and any two sets  $S, T$  such that  $S \subseteq T$ . In simple words, the submodularity means that the marginal gain that we have by adding  $i$  to our purchase set diminishes as we move from one purchase set  $S$  to a superset of it  $T$ . That is, roughly speaking, the more that we already have in the purchase set, the less extra gain by adding an element to it.

Devise an algorithm that purchases a set  $S$  of (approximately) maximum profit, subject to the constraint that  $|S| \leq k$ , for some given value  $k \in \{1, 2, 3, \dots, n\}$ . What approximation factor do you get?

## 3 Connected Dominating Set

Given an  $n$ -node graph  $G = (V, E)$ , a set  $S \subset V$  of vertices is called a *Connected Dominating Set (CDS)* if the following two properties are satisfied: (1) each node  $v \in V$  is either in  $S$  or has a neighbor in  $S$ , (2) the subgraph  $G[S]$  induced by  $S$ —i.e., the one made of  $S$ -vertices and all edges whose both endpoints are in  $S$ —is connected. Devise an approximation algorithm for finding a minimum-cardinality CDS.

## 4 Max-weight Matroid Base (Williamson-Shmoys 2.12)

A matroid  $(\mathcal{E}, \mathcal{I})$  is defined by a ground set  $\mathcal{E}$  of elements and a collection  $\mathcal{I} = \{S_1, S_2, \dots, S_\ell\}$  of *independent* subsets  $S_i \subseteq \mathcal{E}$ , subject to the following conditions:

1. For any two subsets  $S$  and  $S'$  such that  $S \subseteq S'$ , if  $S'$  is independent, then so is  $S$ , i.e.,  $(S' \in \mathcal{I}) \Rightarrow (S \in \mathcal{I})$ .
2. For any two independent sets  $S$  and  $T$  such that  $|S| < |T|$ , there exists an element  $e \in T \setminus S$  such that  $(S \cup \{e\}) \in \mathcal{I}$ .

We call an independent set  $S$  a *base* if there is no  $T \in \mathcal{I}$  such that  $S \subsetneq T$ . That is,  $S$  is in some sense maximal with regard to independence.

Suppose that each element  $e \in \mathcal{E}$  has a weight  $w_e \geq 0$ . Devise an algorithm that finds a maximum-weight base of the matroid.

## 5 Walking on the Hypercube

Consider the  $d$ -dimensional hypercube, which has vertex set  $\{0, 1\}^d$  and where every two vertices that differ in exactly one coordinate are connected by an edge. Suppose that each edge has a random length drawn from an exponential distribution with mean 1, and the lengths of different edges are independent. Devise an algorithm that finds a walk from the vertex  $(0, 0, \dots, 0)$  to the vertex  $(1, 1, \dots, 1)$  with expected length  $O(\log d)$ .

## 6 $k$ -Center

Consider a set of  $n$  points  $P = \{p_1, p_2, \dots, p_n\}$  and a distance metric  $d : P \times P \rightarrow [0, \infty)$  where  $d(p_i, p_j)$  indicates the distance between the two points  $p_i, p_j \in P$ . For a set  $S \subseteq P$ , and an arbitrary point  $p' \in P$ , the distance of  $p'$  to  $S$  is equal to  $\min_{p'' \in S} d(p', p'')$ . The objective is to find a set  $S \subseteq P$  of  $k$  points, called centers, such that the maximum distance of any point  $p' \in P$  to the set  $S$  is minimized. Devise a 2-approximation algorithm for this problem.