

# Solving Algebraic First Order Differential Equations

Aubrey Jaffer

December 6, 1990

The derivative of any algebraic expression is algebraic. First solve the problem of finding antiderivatives where the solution is a rational expression. Work backwards from the form of the solution to completely characterize those derivatives which can lead to the algebraic solution.

## 1 Rational Function Differentiation

Let

$$y = \prod_{i=1}^k p_i(x)^{n_i}$$

be a rational function of  $x$  where the polynomials  $p_i(x)$  are squarefree and mutually relatively prime.

The derivative of  $y$  is

$$y' = x' \sum_{i=1}^k n_i p_i(x)^{n_i-1} p_i'(x) \prod_{j \neq i} p_j(x)^{n_j} \quad (1)$$

**Lemma 1** *The expression  $\sum_{i=1}^k n_i p_i'(x) \prod_{j \neq i} p_j(x)$  has no factors in common with  $p_i(x)$ .*

Assume that the expression has a common factor  $p_h(x)$ . Then

$$p_h(x) \text{ divides } \sum_{i=1}^k n_i p_i'(x) \prod_{j \neq i} p_j(x)$$

Now,  $p_h(x)$  divides all terms for  $i \neq h$  and since it divides the whole sum,  $p_h(x)$  must divide the remaining term  $n_h p_h'(x) \prod_{j \neq h} p_j(x)$ . But, from the above conditions,  $p_h(x)$  does not divide  $p_h'(x)$  [ $p_h(x)$  is squarefree] and  $p_h(x)$  does not divide  $p_j(x)$  for  $j \neq h$  [relatively prime condition].

## 2 Rational Function Integration

Now

$$y' = x' \left( \prod_{i=1}^k p_i(x)^{n_i-1} \right) \sum_{i=1}^k n_i p_i'(x) \prod_{j \neq i} p_j(x) \quad (2)$$

There are no common factors between the sum and product terms of equation 2 because of the relatively prime condition of equation 1 and because of Lemma 1. Hence, this equation cannot be reduced and is canonical.

Split equation 2 into factors with positive and negative exponents and renumber  $i$  to be negative when  $n_i$  is negative, giving

$$y' \prod_{-i} p_i(x)^{-n_i+1} = x' \left( \prod_{+i} p_i(x)^{n_i-1} \right) \sum_i n_i p_i'(x) \prod_{j \neq i} p_j(x) \quad (3)$$

Now to integrate equation 3 note that exponents  $-n_i+1 > 1$  because  $n_i < 0$ . Hence  $\prod_{-i} p_i(x)^{-n_i+1}$  can be factored (easily in fact by squarefree factorization). Now segregate the terms in the sum of equation 2 as well.

$$\begin{aligned} & \sum_i n_i p_i'(x) \prod_{j \neq i} p_j(x) = \\ & \sum_{-i} n_i p_i'(x) \prod_{-j \neq -i} p_j(x) \prod_{+k} p_k(x) + \sum_{+i} n_i p_i'(x) \prod_{+j \neq +i} p_j(x) \prod_{-k} p_k(x) \end{aligned}$$

Substituting into equation 3 yields

$$\begin{aligned} & y' \prod_{-i} p_i(x)^{-n_i+1} = \\ & x' \left( \prod_{+i} p_i(x)^{n_i} \right) \sum_{-i} n_i p_i'(x) \prod_{-j \neq -i} p_j(x) + \\ & x' \left( \prod_{+i} p_i(x)^{n_i-1} \right) \sum_{+i} n_i p_i'(x) \prod_{+j \neq +i} p_j(x) \prod_{-k} p_k(x) \end{aligned} \quad (4)$$

The right side of this equation is now grouped into four polynomial terms  $AB' + A'B$  where

$$\begin{aligned} A &= \prod_{+i} p_i(x)^{n_i} \\ B' &= \sum_{-i} n_i p_i'(x) \prod_{-j \neq -i} p_j(x) \\ A' &= \left( \prod_{+i} p_i(x)^{n_i-1} \right) \sum_{+i} n_i p_i'(x) \prod_{+j \neq +i} p_j(x) \\ B &= \prod_{-k} p_k(x) \end{aligned}$$

$A$  is the original numerator and  $A'$  it's derivative.  $B$  and  $B'$  can be derived from the squarefree factorization of the denominator of the integrand.  $A$  and  $A'$  can be recovered by a kind of long division of the right side of equation 4 by  $B$  and  $B'$  simultaneously. In addition to subtracting a term times  $B'$  subtract the term's derivative times  $B$ .

### 3 A First Order Differential Equation

Starting with equation 1 multiply through by  $\prod_{i=1}^k p_i(x)$  and replace  $\prod_{i=1}^k p_i(x)^{n_i}$  on the right side by  $y$ .

$$y' \prod_{i=1}^k p_i(x) = x'y \sum_{i=1}^k n_i p_i'(x) \prod_{j \neq i} p_j(x) \quad (5)$$

By Lemma 1 this cannot be simplified because the two sides have no factor in common. Hence, this form is canonical.

Therefore, given an equation of form  $y'q(x) = x'yr(x)$ , if it can be put into the form of equation 5, it can be solved as in equation 1. In order to do this we need to factor  $q(x)$ . This factoring can be seen as the same complexity as the partial fraction decomposition in Risch's algorithm.

Once we have factored  $q(x)$ , we need to find a set of  $n_i$  so that

$$\sum_{i=1}^k n_i p_i'(x) \prod_{j \neq i} p_j(x) = r(x)$$

. Now in order for this solution to be unique we need to show that the terms  $p_i'(x) \prod_{j \neq i} p_j(x)$  are linearly independent and hence form the basis for a vector space. Let's assume that they were not independent.

Suppose there existed a set of integers  $m_i$  such that

$$\sum_{i=1}^k m_i p_i'(x) \prod_{j \neq i} p_j(x) = 0$$

and there exists some  $m_i \neq 0$ . If only one  $m_i \neq 0$  then  $p_i'(x) \prod_{j \neq i} p_j(x) = 0$ . Since  $p_j(x) \neq 0$  then  $p_i'(x) = 0$ . But then  $p_i(x)$  would not be a polynomial in  $x$ . So then

$$-m_i p_i' \prod_{j \neq i} p_j(x) = \sum_{h \neq i} m_h p_h'(x) \prod_{j \neq h} p_j(x) \quad (6)$$

Now,  $p_i(x)$  divides every term on the right side of equation 6 so  $p_i(x)$  must also divide  $-m_i p_i'(x) \prod_{j \neq i} p_j(x)$ . But, because of squarefree,  $p_i(x)$  does not divide  $p_i'(x)$  and  $p_i(x)$  does not divide  $p_j(x)$  when  $j \neq i$ . Hence, there exists a unique set of coefficients satisfying equation 5.