Today

- Amplification of error
- BPP in $P/_{\text{poly}}$.
- BPP in PH.

Clarification on Games

Few lectures back .... we said some wrong things.

- Game is in PSPACE only if there is an a priori polynomial upper bound on its running time.
- Go: # of pieces on board increase all the time.
- Geography: Path length bounded by size of Atlas.
- Chess: No "a priori" upper bound - hence not known to be in PSPACE.

Last lecture

- Introduced randomness.
- Defined many classes (BPP, RP, RL etc.)
- Showed Poly. Ident. Testing in RP.
- Claimed USTCON in RL.
- Next on agenda: completeness and soundness.

RP Amplification

Suppose M accepts language L with completeness $c(n) = 1/n^2$ (and $s(n) = 0$). How to amplify completeness?

Amplification: Run machine n^4 times on independent random strings $y_1, \ldots, y_{n^4}$, and accept if one of the y_i's accepts.

$$\Pr_{\mathbf{y}}[\exists i \text{ s.t. } M(x, y_i)\text{accepts}] \geq 1-(1-1/n^2)^{n^4} \geq 1-$$

Thus completeness $1/\text{poly}(n)$ vs. $1 - \exp(n)$ are equivalent.

BPP amplification

- How to use the above idea for BPP?
- Natural idea:
  - Repeat N times.
  - Accept if # acceptances more than $(c+s)N/2$.
- Analysis?
  - Use "tail inequalities".
  - "Chernoff bound".

Chernoff bounds

Suppose $X_1, \ldots, X_N$ are independent identically distributed random variables in the interval $[0,1]$ with $\mathbf{E}[X_i] = \mu$.

Then

$$\Pr[|\frac{1}{N}\sum_i X_i - \mu| \geq \lambda] \leq e^{-\lambda^2 N/2}.$$

Consequence

Let $X_i = 1$ if $M(x, y_i)$ accepts and 0 o.w.

Applying Chernoff bounds, we see that if $N \sim m/(c-s)^2$ then amplification increases completeness to $1 - \exp(-m)$ and decreases soundness to $\exp(-m)$.

Next: Use this to show BPP in $\mathrm{P}/_{\mathrm{poly}}$.

Consequence: BPP in $\mathrm{P}/_{\mathrm{poly}}$

Say $L \in \mathrm{BPP}$. Assume w.l.o.g. that M is a two input machine recognizing L with $c(n) \geq 1 - 4^{-n}$ and $s(n) \leq 1 - 4^{-n}$. (Notice we get this by amplification.)

Say M uses m-bit random strings.

Claim: Exists $r \in \{0,1\}^m$ such that for every x, $M(x, r) = L(x)$.

Proof: Say $y \in \{0,1\}^m$ is BAD for x if $M(x, y) \neq L(x)$.

For any $x \in \{0,1\}^n$ there are at most 2^{m-2n} y's that are BAD for x.

Taking the union of all BAD sets, there are at most 2^{m-n} strings that are BAD for some x.

Since $2^m > 2^{m-n}$ there exists at least one y which is not BAD for any x. Setting $r \leftarrow y$ gives the Claim.

Thm: BPP $\subseteq$ P/$_{\mathrm{poly}}$.

Proof: P/$_{\mathrm{poly}}$ machine is M from the argument above. For every n, advice string is the $r \in \{0,1\}^m$ from the claim.

Next: BPP in PH

Note note quite trivial. How to have a bounded round interaction to comvince $x \in L$?

Consider following game: Deniss & I are all powerful players. I want to convince you (the audience) that $x \in L$ and Deniss claims otherwise. How can we prove our claims?

Draw picture here.

Most strings are good (M(x,y) $=$ accept); or very few are good. How to convince you?

Idea 1: I'll divide space into two equal parts with all bad strings in one part and a bijection pi between the two parts. I claim every string

or its map under bijection is good! If Deniss wants, he can challenge me!

If Deniss finds a string y where neither M(x,y) nor M(x,pi(y)) accept - he wins.

Else I win.

Seems convincing. I can win if bad set is smaller than 1/2. I can't win if bad set more than 1/2.

Problem: How do I give the bijection?

Bijections have to simple: So we'll stick $\pi_r : y \mapsto y \oplus r$.

In this space of bijections the proof doesn't go through. But the idea is starting to emanate.

Debate for membership in BPP

Theorem: If x in L there exist $r_1, \ldots, r_{2m} \in \{0,l\}^m$ such that the y's are covered; i.e., for every y there exists an $i \in [2m]$ such that $M(x, \pi_{r_i}(y))$ accepts.

If x not in L, then for any $r_1, \ldots, r_{2m} \in \{0,l\}^m$ there is an uncovered y.

Assuming theorem: Debate: I announce $r_1, \ldots, r_{2m}$. Deniss challenges with a y. You compute $M(x, y \oplus r_1) \vee \cdots \vee M(x, y \oplus r_{2m})$. If true, I win ($x \in L$) else Deniss wins ($x \notin L$) - you decide!

Proof of theorem

If x in L

$$\Pr_r[M(x, y \oplus r)] \geq 1 - 2^{-n} \geq 1/2.$$

$$\Pr_{r_1,\cdots,r_{2m}}[\exists i \in [2m] \text{ s.t. } M(x, y \oplus r_i)] \geq 1 - 2^{-2m}.$$

$$\Pr_{r_1,\cdots,r_{2m}}[\forall y \in \{0,1\}^m, \exists i \in [2m] \text{ s.t. } M(x, y \oplus r_i)]$$

Yields first part.

Proof of theorem (second part)

x not in L. Say I pick best possible $r_1, \ldots, r_{2m}$ below.

$$\Pr_y[M(x, y \oplus r_i)] \leq 1/100m.$$

$$\Pr_y[\exists i \in [2m] \text{ s.t. } M(x, y \oplus r_i)] \leq 1/50.$$

QED!

Power of the prover

If I am right - I just need to pick $r_1, \ldots, r_{2m}$ at random!

If Deniss is right, he just needs to pick y at random.

So we just need randomness to simulate randomness!

Hmm.... that didn't sound so impressive - I should have said ...

So we just need one-sided randomness to simulate two-sided randomness! You'll figure out what I mean in problem set!

Current issues in randomness

- Reducing randomness
  - Algorithm specific: Limited independence, Epsilon-bias.
  - Generically, during amplification: "Recycling".
- Using imperfect randomness: Extractors.
- Derandomization: Pseudorandomness, hardness versus randomness.