

CRESTED PRODUCTS OF MARKOV CHAINS

BY DANIELE D'ANGELI AND ALFREDO DONNO

Université de Genève

In this work we define two kinds of crested product for reversible Markov chains, which naturally appear as a generalization of the case of crossed and nested product, as in association schemes theory, even if we do a construction that seems to be more general and simple. Although the crossed and nested product are inspired by the study of Gelfand pairs associated with the direct and the wreath product of two groups, the crested products are a more general construction, independent from the Gelfand pairs theory, for which a complete spectral theory is developed. Moreover, the k -step transition probability is given. It is remarkable that these Markov chains describe some classical models (Ehrenfest diffusion model, Bernoulli–Laplace diffusion model, exclusion model) and give some generalization of them.

As a particular case of nested product, one gets the classical Insect Markov chain on the ultrametric space. Finally, in the context of the second crested product, we present a generalization of this Markov chain to the case of many insects and give the corresponding spectral decomposition.

1. Introduction. The starting point for this work is the paper [4] on Gelfand pairs (for a general theory and applications see [7]) by Ceccherini-Silberstein, Scarabotti and Tolli. In that paper, particular cases of homogeneous spaces are presented. Given two finite groups G and F , together with two subgroups $K \leq G$ and $H \leq F$, let $X = G/K$ and $Y = F/H$ be the corresponding homogeneous spaces. The Gelfand pairs associated with the direct product $G \times F$ and the wreath product $F \wr G$ are studied. A complete spectral decomposition, together with the associated spherical functions, is given for the space $L(X \times Y)$.

One can ask if it is possible to get the same spectral decomposition without using the group structure, but considering the product (in some sense)

Received February 2007; revised February 2008.

AMS 2000 subject classifications. 60J10, 15A69, 05E30, 05C25, 43A85.

Key words and phrases. Reversible Markov chain, crested product, nested product, crossed product, spectral theory, association schemes, Gelfand pairs.

<p>This is an electronic reprint of the original article published by the Institute of Mathematical Statistics in <i>The Annals of Applied Probability</i>, 2009, Vol. 19, No. 1, 414–453. This reprint differs from the original in pagination and typographic detail.</p>

of two reversible Markov chains P and Q defined on two finite sets X and Y , respectively.

The analogue of direct product and wreath product is developed in Sections 5 and 6, respectively. In particular, in Section 6, the Markov chain on the ultrametric space, studied by Figà-Talamanca in [11] and that we call “Insect Markov chain,” is revisited from a different point of view. See also [6].

An equivalent approach to Gelfand pairs product comes from the association schemes theory (see [2]). In particular, the direct and the wreath product of groups correspond to the crossed and nested product of association schemes. In [3] Bailey and Cameron introduced a more general product for association schemes called the crested product.

Inspired by this definition, in this paper two new products of Markov chains are introduced. We will refer to them as the *first* and the *second* crested product, respectively. Actually, the definition does not seem to be the exact analogue, in terms of Markov chains, of the crested product for association schemes, but it looks even more interesting. In fact, they extend the crossed and the nested product and a complete spectral theory is presented in Sections 4 and 7.

In particular, in Section 7, we show that the second crested product is a generalization of a Markov chain that we will call multi-insect, which is an analogue of the so-called nonbinary Johnson scheme, already studied in [4], that corresponds to considering h insects living in different subtrees and moving only one per each step in such a way that their distance is preserved. We give a generalization of the results of [4] in Section 7 with a more elementary and direct proof. This problem has been suggested by Scarabotti. As an example, the case of the bi-insect is studied.

2. Basic examples and motivations. In this section we present some examples, classical and new, of finite diffusion processes that one can treat using methods from harmonic analysis. In the next sections, we will study them by introducing special products of finite Markov chains and performing a complete spectral analysis of the associated Markov operator.

2.1. Classical models.

2.1.1. *The Ehrenfest diffusion model.* The classical Ehrenfest diffusion model consists of two urns numbered 0, 1 and n balls numbered $1, \dots, n$. A configuration is given by a placement of the balls into the urns. Therefore there are 2^n configurations and each of them can be identified with the subset A of $\{1, \dots, n\}$ corresponding to the set of balls contained, for instance, in the urn 0. Note that there is no ordering inside the urns. The initial configuration corresponds to the situation when all balls are inside the urn 0.

Then, at each step, a ball is randomly chosen (with probability $1/n$) and it is moved to the other urn. Denoting by Q_n the set of all subsets of $\{1, \dots, n\}$, we can describe this random process as follows. If we are in a configuration $A \in Q_n$, then, at the next step, we are in a new configuration $B \in Q_n$ of the form $A \sqcup \{j\}$ for some $j \notin A$ or $A \setminus \{j'\}$ for some $j' \in A$ and each of these configurations is chosen with probability $1/n$.

In [5], the authors give a generalization of this model to the case of m urns and n balls.

2.1.2. *The Bernoulli–Laplace diffusion model.* The Bernoulli–Laplace diffusion model consists of two urns numbered 0,1 and $2n$ balls, numbered $1, \dots, 2n$. A configuration is a placement of the balls into the two urns, n balls each. Therefore there are $\binom{2n}{n}$ configurations; each of them can be identified with an n -subset A of $\{1, \dots, 2n\}$ corresponding to the set of balls contained in the urn 0. The initial configuration corresponds to the situation when the balls contained in the urn 0 are $1, 2, \dots, n$. Denoting by Ω_n the set of all n -subsets of $\{1, 2, \dots, 2n\}$, we can describe this random process as follows. If we are in a configuration $A \in \Omega_n$, then at the next step we are in a new configuration $B \in \Omega_n$ of the form $A \sqcup \{i\} \setminus \{j\}$, for some $i \notin A$ and $j \in A$ and each of these configurations is chosen with probability $1/n^2$.

2.1.3. *The exclusion model.* Let (X, E) be an undirected connected graph with n vertices. Fix $h \leq n$ and choose an h -subset A of X . We put a particle on each vertex of A . Then the exclusion process is defined as a Markov chain on the set Ω_h of h -subsets of X . Starting from the state A , we pick a particle on some vertex of A with probability proportional to the degree of the vertex that it occupies, pick a neighboring site of this vertex at random and move the particle to the neighboring site provided this site is unoccupied. If the site is occupied by another particle, the chain stays at A . See [8] and [12] for more examples and details.

2.1.4. *The Insect Markov chain.* This Markov chain, in its classical version introduced by Figà-Talamanca [11], is performed on the (finite) ultrametric space given by the n th level of a q -ary rooted tree. At the starting instant the insect stays in a fixed vertex of the n th level. It performs a simple random walk on the tree. The next state of the Markov chain is given when the insect reaches again the n th level. The probability transition between two vertices only depends on their ultrametric distance and so this Markov chain is invariant under the action of the full automorphisms group of the rooted tree.

2.2. Generalizations.

2.2.1. *The first crested product.* The Ehrenfest model, generalized to the case of n balls and m urns, is described by the *crossed product* of n Markov chains defined on a space of cardinality m (the space of the urns), as will be shown in Section 5. Actually, the definition of crossed product that we give admits that the choice of the i th ball is weighted by a not necessarily homogeneous probability distribution p_i^0 .

One can ask what happens if a hierarchy in the set of balls is introduced. So suppose to have n ordered balls, numbered by $1, \dots, n$, and m urns.

At each step, one ball (say the i th one) is randomly chosen and it is moved into another urn following a transition probability P_i defined on the space of the urns. For all $j > i$, the ball j is also randomly moved to another urn.

This model is described by the *nested product* of n Markov chains defined on a space of cardinality m (the space of the urns), as will be shown in Section 6, where we suppose that each ball is chosen according with a probability distribution p_i^0 .

The Insect Markov chain described in Section 6.3 is obtained from this model for a particular value of p_i^0 and forcing the transition probability P_i to be uniform, as shown in Proposition 6.4.

The *first crested product* introduced in Section 4 is a generalization of both crossed and nested product and it describes the following diffusion model.

The (C, N) -Ehrenfest model. We have n balls numbered by $1, \dots, n$ and m urns. Suppose that we have a partition of the set of the balls in two subsets $C \amalg N$. The balls in C are white, the balls in N are black. At each step, we choose a ball i according with a probability distribution p_i^0 : if it is white, then we move it to another urn following a transition probability P_i and all the other balls are not moved. If it is black, we move it by P_i and then each ball (white or black) numbered by j , for $j > i$, is moved uniformly to a new urn.

The card players model. The previous model can also be described in the following way. Suppose that n card players sit at the same edge of a table. Each of them has a deck of cards consisting of m cards. Suppose that the set of players is partitioned in two subsets: blond hair and brown hair. At the starting moment, each player has a card. A player is randomly chosen, so he puts his card inside the deck and then chooses another card. If he is blond, the others keep their own card; if he is brown, then all the players (blond or brown) sitting at his right side also must change their card.

2.2.2. *The second crested product.*

The bi-insect. This is a generalization of the Insect Markov chain. In this case we have two insects that, at the starting time, live in the n th level of a tree, following the condition that their (ultrametric) distance is maximal, which is equivalent to requiring that they live in two different subtrees of depth $n - 1$. At each step, with probability p_0 they do not leave their own subtrees; with probability $1 - p_0$ one of them can leave its subtree choosing another subtree not occupied by the other insect. In the first case, we randomly choose one insect and change its position according with the Insect Markov chain on the corresponding subtree. In the second case, the moved insect randomly chooses its new position in the new subtree that it has occupied. This yields a Markov chain on the space of all possible configurations of two insects at maximal distance in a tree of depth n ; this Markov chain will be studied in Section 7.2.

The multi-insect. An immediate generalization of the previous construction can be performed considering h insects living in the n th level of the tree and having maximal distance from each other. At each step, with probability p_0 only one insect moves to another site staying in the same subtree; with probability $1 - p_0$ only one insect occupies a new subtree of depth $n - 1$ and randomly chooses its position. This situation will be referred to as the *multi-insect* in Section 7.

Moreover, the multi-insect can be also regarded as an exclusion process on the ultrametric space with the constraint that the h insects keep maximal distance from each other. At each step an insect moves, as in the exclusion process a particle does.

The aim of this work is also to diagonalize a more general Markov chain, that we call the *second crested product*, described in Section 7 and that corresponds to the following model.

The crested Bernoulli–Laplace model. Suppose that we have two urns numbered 0, 1 and n balls: the urn 0 contains h balls and the urn 1 contains the remaining $n - h$ balls. The h balls in the urn 0 are endowed with a label that can go from 1 to s (note that two different balls can have the same label). Moreover, we can use a (nonhomogeneous) coin. At each step, we flip the coin: if we get heads, then we randomly choose a ball i in the urn 0 and a ball j in the urn 1 and exchange them. Before putting j in the urn 0 we attach a label (randomly chosen in $\{1, \dots, s\}$) on it.

If we get tails, then we randomly choose a ball i in the urn 0 and we change or not its label according with a transition probability Q on the set $\{1, \dots, s\}$.

Observe that, if we get heads, then we perform on the set of the balls an analogue of the Bernoulli–Laplace diffusion model, where the urns contain h and $n - h$ balls, respectively.

Although in Section 7.1 the spectral analysis of the second crested product is given, in Section 7.2 we give explicit computations only in the case of the bi-insect, which seems to be easier to handle.

2.3. *Motivations.* The classical models described in the previous sections are well known in literature and they have been studied by many authors using approaches deriving from Gelfand pairs theory, representations theory, associations schemes, distance regular graphs and orthogonal polynomials.

In the Ehrenfest model, generalized to the case of m urns and n balls, the set of all configurations can be identified with the Hamming scheme $X_{n,m} = \{0, \dots, m-1\}^n$ consisting of all the functions defined on the set $\{1, \dots, n\}$ taking values in $\{0, \dots, m-1\}$. The group that naturally acts on $X_{n,m}$ is the wreath product $S_m \wr S_n$. The stabilizer of a fixed function in $X_{n,m}$ is isomorphic to the subgroup $S_{m-1} \wr S_n$. In [5], for example, the authors show that $(S_m \wr S_n, S_{m-1} \wr S_n)$ is a Gelfand pair, they give the decomposition of the space $L(X_{n,m})$ of the complex functions defined on $X_{n,m}$ into irreducible submodules and provide an expression for the respective spherical functions by using Krawtchouk polynomials. These submodules coincide with the eigenspaces of the Markov operator on $L(X_{n,m})$ describing the Ehrenfest model. A different approach to the Hamming scheme is given in [2] by using association schemes theory.

Following the idea of Diaconis, it is possible to study the rate of convergence to the stationary distribution of finite Markov chains by using Gelfand pairs theory. In particular, given a Markov chain which is invariant under the action of a group G , its transition operator can be expressed via a Fourier series where the classical exponentials are replaced by the irreducible representations of G . In [10] the authors apply this argument to the generalized Bernoulli–Laplace diffusion model, consisting of two urns containing h and $n-h$ balls, respectively. All the states of this Markov chain constitute the so-called Johnson scheme associated with the Gelfand pair $(S_n, S_h \times S_{n-h})$. The corresponding decomposition into irreducible subrepresentations (and so into eigenspaces for the associated Markov operator) is given, for instance, in [5], where the spherical functions are described in terms of Hahn polynomials. Like the Hamming scheme, also the Johnson scheme can be presented as an association scheme (see [2]).

Finally, the Insect Markov chain can be studied using a Gelfand pair associated with the automorphisms group of the rooted tree (see [6] and [11]). The Fourier analysis that one can develop allows to give a complete spectral analysis and to investigate some ergodic properties.

The approach proposed in this paper is more elementary. We define and study several Markov chains, generalizing the classical models described above, by using only techniques and arguments coming from linear and multilinear algebra. This allows to give directly the spectral analysis of the

associated Markov operator, without invoking the theory of groups and their representations. Our construction shows what really are the structures to the base of the analysis of these models and suggests the way to approach more general cases.

In the case of the first crested product we consider the Cartesian product $X = X_1 \times \cdots \times X_n$ of n finite spaces and a Markov operator P on $L(X)$ given by a convex combination of Markov operators on $L(X_i)$ depending on a partition of the set $\{1, \dots, n\}$. The eigenspaces of the operator P are expressed as certain combinations of the tensor products of the eigenspaces of the Markov operators on every space $L(X_i)$. Their form is easier in the case of the crossed product when none of the spaces X_i is distinguished from the other ones (see Section 5). On the other hand, it becomes more complicated if one introduces a hierarchy. The extremal case gives the nested product (see Section 6), where the space X can be identified with the n th level of a rooted tree.

In the second crested product we consider the Cartesian product of two finite spaces X and Y , where we are given the eigenspaces of a Markov operator Q defined on $L(Y)$. Set Θ_h the space of the functions with domain an h -subset of X and values in Y . We define a Markov operator P [see (6)] on $L(\Theta_h)$ and we investigate its spectral decomposition by using two differential operators D and D^* (see Definition 7.1) and the known spectral analysis of Q .

Although the computational aspect can be more or less complicated depending on the singular cases, the methods used do not require anything more than linear or multilinear algebra.

3. Preliminaries. The following topics about finite Markov chains can be found in [6].

Consider a finite set X , with $|X| = m$. Let P be a stochastic matrix of size m whose rows and columns are indexed by the elements of X , so that

$$\sum_{x \in X} p(x_0, x) = 1,$$

for every $x_0 \in X$. Consider the Markov chain on X with transition matrix P .

DEFINITION 3.1. The Markov chain P is reversible if there exists a strict probability measure π on X such that

$$\pi(x)p(x, y) = \pi(y)p(y, x),$$

for all $x, y \in X$.

We will say that P and π are in detailed balance. For a complete treatment about these and related topics see [1].

Define on $L(X) = \{f : X \rightarrow \mathbb{C}\}$ a scalar product in the following way:

$$\langle f_1, f_2 \rangle_\pi = \sum_{x \in X} f_1(x) \overline{f_2(x)} \pi(x),$$

for all $f_1, f_2 \in L(X)$ and the linear operator $P : L(X) \rightarrow L(X)$ by

$$(Pf)(x) = \sum_{y \in X} p(x, y) f(y).$$

It is easy to verify that π and P are in detailed balance if and only if P is self-adjoint with respect to the scalar product $\langle \cdot, \cdot \rangle_\pi$. Under these hypotheses, it is known that the matrix P can be diagonalized over the reals. Moreover, 1 is always an eigenvalue of P and for any eigenvalue λ one has $|\lambda| \leq 1$.

Let λ_z be the eigenvalues of the matrix P , for every $z \in X$, with $\lambda_{z_0} = 1$. Then there exists an invertible unitary real matrix $U = (u(x, y))_{x, y \in X}$ such that $PU = U\Delta$, where $\Delta = (\lambda_x \delta_x(y))_{x, y \in X}$ is the diagonal matrix whose entries are the eigenvalues of P . This equation gives, for all $x, z \in X$,

$$(1) \quad \sum_{y \in X} p(x, y) u(y, z) = u(x, z) \lambda_z.$$

Moreover, we have $U^T D U = I$, where $D = (\pi(x) \delta_x(y))_{x, y \in X}$ is the diagonal matrix of coefficients of π . This second equation gives, for all $y, z \in X$,

$$(2) \quad \sum_{x \in X} u(x, y) u(x, z) \pi(x) = \delta_y(z).$$

Hence, the first equation tells us that each column of U is an eigenvector of P ; the second one tells us that these columns are orthogonal with respect to the product $\langle \cdot, \cdot \rangle_\pi$.

PROPOSITION 3.2. *The k th step transition probability is given by*

$$(3) \quad p^{(k)}(x, y) = \pi(y) \sum_{z \in X} u(x, z) \lambda_z^k u(y, z),$$

for all $x, y \in X$.

PROOF. The proof is a consequence of (1) and (2). In fact, the matrix $U^T D$ is the inverse of U , so that $U U^T D = I$. In formulæ, we have

$$\sum_{y \in X} u(x, y) u(z, y) = \frac{1}{\pi(z)} \Delta_z(x).$$

From the equation $PU = U\Delta$ we get $P = U\Delta U^T D$, which gives

$$p(x, y) = \pi(y) \sum_{z \in X} u(x, z) \lambda_z u(y, z).$$

Iterating this argument, we get

$$P^k = U \Delta^k U^T D,$$

which is the assertion. \square

Recall that there exists a correspondence between reversible Markov chains and weighted graphs.

DEFINITION 3.3. A weight on a graph $\mathcal{G} = (X, E)$ is a function $w : X \times X \rightarrow [0, +\infty)$ such that:

- (1) $w(x, y) = w(y, x)$;
- (2) $w(x, y) > 0$ if and only if $x \sim y$.

If \mathcal{G} is a weighted graph, it is possible to associate with w a stochastic matrix $P = (P(x, y))_{x, y \in X}$ on X by setting

$$p(x, y) = \frac{w(x, y)}{W(x)},$$

with $W(x) = \sum_{z \in X} w(x, z)$. The corresponding Markov chain is called the random walk on \mathcal{G} . It is easy to prove that the matrix P is in detailed balance with the distribution π defined, for every $x \in X$, as

$$\pi(x) = \frac{W(x)}{W},$$

with $W = \sum_{z \in X} W(z)$. Moreover, π is strictly positive if X does not contain isolated vertices. The inverse construction can be done. So, if we have a transition matrix P on X which is in detailed balance with the probability π , then we can define a weight w as $w(x, y) = \pi(x)p(x, y)$. This definition guarantees the symmetry of w and, by setting $E = \{\{x, y\} : w(x, y) > 0\}$, we get a weighted graph.

There are some important relations between the weighted graph associated with a transition matrix P and its spectrum. In fact, it is easy to prove that the multiplicity of the eigenvalue 1 of P equals the number of connected components of \mathcal{G} . Moreover, the following propositions hold.

PROPOSITION 3.4. *Let $\mathcal{G} = (X, E, w)$ be a finite connected weighted graph and denote P the corresponding transition matrix. Then the following are equivalent:*

- (1) \mathcal{G} is bipartite;
- (2) the spectrum $\sigma(P)$ is symmetric;
- (3) $-1 \in \sigma(P)$.

DEFINITION 3.5. Let P be a stochastic matrix. P is ergodic if there exists $n_0 \in \mathbb{N}$ such that

$$p^{(n_0)}(x, y) > 0 \quad \text{for all } x, y \in X.$$

PROPOSITION 3.6. Let $\mathcal{G} = (X, E)$ be a finite graph. Then the following conditions are equivalent:

- (1) \mathcal{G} is connected and not bipartite;
- (2) for every weight function on X , the associated transition matrix P is ergodic.

So we can conclude that a reversible transition matrix P is ergodic if and only if the eigenvalue 1 has multiplicity 1 and -1 is not an eigenvalue.

In what follows we always suppose that the eigenvalue 1 has multiplicity 1, so that the graph associated with the probability P is connected. This is equivalent to requiring that the probability P is irreducible, according with the following definition.

DEFINITION 3.7. A stochastic matrix P on a set X is irreducible if, for every $x_1, x_2 \in X$, there exists $n = n(x_1, x_2)$ such that $p^{(n)}(x_1, x_2) > 0$.

4. The first crested product. In this section we introduce a particular product of Markov chains defined on different sets. This idea is inspired by the definition of crested product for association schemes given in [3].

Let X_i be a finite set, with $|X_i| = m_i$, for every $i = 1, \dots, n$, so that we can identify X_i with the set $\{0, 1, \dots, m_i - 1\}$. Let P_i be an irreducible Markov chain on X_i and let p_i be the transition probability associated with P_i . Moreover, assume that p_i is in detailed balance with the strict probability measure σ_i on X_i , that is,

$$\sigma_i(x)p_i(x, y) = \sigma_i(y)p_i(y, x),$$

for all $x, y \in X_i$.

Consider the product $X_1 \times \dots \times X_n$. Let $\{1, \dots, n\} = C \amalg N$ be a partition of the set $\{1, \dots, n\}$ and let $p_1^0, p_2^0, \dots, p_n^0$ be a probability distribution on $\{1, \dots, n\}$, that is, $p_i^0 > 0$ for every $i = 1, \dots, n$ and $\sum_{i=1}^n p_i^0 = 1$.

DEFINITION 4.1. The first crested product of Markov chains P_i 's with respect to the partition $\{1, \dots, n\} = C \amalg N$ is the Markov chain on the product $X_1 \times \dots \times X_n$ whose transition matrix is

$$\begin{aligned} P &= \sum_{i \in C} p_i^0 (I_1 \otimes \dots \otimes I_{i-1} \otimes P_i \otimes I_{i+1} \otimes \dots \otimes I_n) \\ &\quad + \sum_{i \in N} p_i^0 (I_1 \otimes \dots \otimes I_{i-1} \otimes P_i \otimes J_{i+1} \otimes \dots \otimes J_n), \end{aligned}$$

where I_i denotes the identity matrix of size m_i and J_i denotes the uniform matrix on X_i , that is,

$$J_i = \frac{1}{m_i} \begin{pmatrix} 1 & 1 & \cdots & 1 \\ 1 & \ddots & & \vdots \\ \vdots & & \ddots & \vdots \\ 1 & \cdots & \cdots & 1 \end{pmatrix}.$$

In other words, we choose an index i in $\{1, \dots, n\}$ following the distribution p_1^0, \dots, p_n^0 . If $i \in C$, then P acts on the i th coordinate by the matrix P_i and fixes the remaining coordinates; if $i \in N$, then P fixes the coordinates $\{1, \dots, i-1\}$, acts on the i th coordinate by the matrix P_i and changes uniformly the remaining ones.

For all $(x_1, \dots, x_n), (y_1, \dots, y_n) \in X_1 \times \cdots \times X_n$, the transition probability p associated with P is given by

$$\begin{aligned} & p((x_1, \dots, x_n), (y_1, \dots, y_n)) \\ &= \sum_{i \in C} p_i^0 (\delta_1(x_1, y_1) \cdots \delta_{i-1}(x_{i-1}, y_{i-1}) \\ & \quad \times p_i(x_i, y_i) \delta_{i+1}(x_{i+1}, y_{i+1}) \cdots \delta_n(x_n, y_n)) \\ & \quad + \sum_{i \in N} p_i^0 \left(\frac{\delta_1(x_1, y_1) \cdots \delta_{i-1}(x_{i-1}, y_{i-1}) p_i(x_i, y_i)}{\prod_{j=i+1}^n m_j} \right), \end{aligned}$$

where δ_i is defined by

$$\delta_i(x_i, y_i) = \begin{cases} 1, & \text{if } x_i = y_i, \\ 0, & \text{otherwise.} \end{cases}$$

We want to study the spectral theory of the operator P . We recall that the following isomorphism holds:

$$L(X_1 \times \cdots \times X_n) \cong \bigotimes_{i=1}^n L(X_i),$$

with $(f_1 \otimes \cdots \otimes f_n)(x_1, \dots, x_n) := f_1(x_1) f_2(x_2) \cdots f_n(x_n)$.

Assume that, for every $i = 1, \dots, n$, the following spectral decomposition holds:

$$L(X_i) = \bigoplus_{j_i=0}^{r_i} V_{j_i}^i,$$

that is, $V_{j_i}^i$ is an eigenspace for P_i with associated eigenvalue λ_{j_i} and whose dimension is m_{j_i} .

Now set $N = \{i_1, \dots, i_l\}$ and $C = \{c_1, \dots, c_h\}$, with $h + l = n$ and such that $i_1 < \cdots < i_l$ and $c_1 < \cdots < c_h$.

THEOREM 4.2. *The probability P defined above is reversible if and only if P_k is symmetric for every $k > i_1$. If this is the case, P is in detailed balance with the strict probability measure π on $X_1 \times \cdots \times X_n$ given by*

$$\pi(x_1, \dots, x_n) = \frac{\sigma_1(x_1)\sigma_2(x_2) \cdots \sigma_{i_1}(x_{i_1})}{m_{i_1+1} \cdots m_n}.$$

PROOF. Consider the elements $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ belonging to $X_1 \times \cdots \times X_n$. First, we want to prove that the condition $\sigma_k = \frac{1}{m_k}$, for every $k > i_1$, is sufficient. Let $k \in \{1, \dots, n\}$ such that $x_i = y_i$ for every $i = 1, \dots, k-1$ and $x_k \neq y_k$. Suppose $k < i_1$. Then we have

$$p(x, y) = p_k^0(p_k(x_k, y_k)\delta_{k+1}(x_{k+1}, y_{k+1}) \cdots \delta_n(x_n, y_n)).$$

If $x_i = y_i$ for every $i = k+1, \dots, n$, we get

$$\begin{aligned} \pi(x)p(x, y) &= \sigma_1(x_1) \cdots \sigma_k(x_k) \cdots \sigma_{i_1}(x_{i_1}) p_k^0 \frac{p_k(x_k, y_k)}{m_{i_1+1} \cdots m_n} \\ &= \sigma_1(y_1) \cdots \sigma_k(y_k) \cdots \sigma_{i_1}(y_{i_1}) p_k^0 \frac{p_k(y_k, x_k)}{m_{i_1+1} \cdots m_n} \\ &= \pi(y)p(y, x), \end{aligned}$$

since $\sigma_k(x_k)p_k(x_k, y_k) = \sigma_k(y_k)p_k(y_k, x_k)$. If the condition $x_i = y_i$ is not satisfied for every $i = k+1, \dots, n$, then the equality $\pi(x)p(x, y) = \pi(y)p(y, x) = 0$ easily follows.

If $k = i_1$, then we get

$$p(x, y) = p_{i_1}^0 \left(p_{i_1}(x_{i_1}, y_{i_1}) \frac{1}{m_{i_1+1} \cdots m_n} \right)$$

and so

$$\begin{aligned} \pi(x)p(x, y) &= \sigma_1(x_1) \cdots \sigma_{i_1}(x_{i_1}) p_{i_1}^0 \frac{p_{i_1}(x_{i_1}, y_{i_1})}{m_{i_1+1}^2 \cdots m_n^2} \\ &= \sigma_1(y_1) \cdots \sigma_{i_1}(y_{i_1}) p_{i_1}^0 \frac{p_{i_1}(y_{i_1}, x_{i_1})}{m_{i_1+1}^2 \cdots m_n^2} \\ &= \pi(y)p(y, x), \end{aligned}$$

since $\sigma_{i_1}(x_{i_1})p_{i_1}(x_{i_1}, y_{i_1}) = \sigma_{i_1}(y_{i_1})p_{i_1}(y_{i_1}, x_{i_1})$.

In the case $k > i_1$, we have

$$p(x, y) = \sum_{i \in N, i \leq k} p_i^0 \frac{p_i(x_i, y_i)}{m_{i+1} \cdots m_n}$$

and so

$$\begin{aligned}\pi(x)p(x, y) &= \frac{\sigma_1(x_1) \cdots \sigma_{i_1}(x_{i_1})}{m_{i_1+1} \cdots m_n} \sum_{i \in N, i \leq k} p_i^0 \frac{p_i(x_i, y_i)}{m_{i+1} \cdots m_n} \\ &= \frac{\sigma_1(y_1) \cdots \sigma_{i_1}(y_{i_1})}{m_{i_1+1} \cdots m_n} \sum_{i \in N, i \leq k} p_i^0 \frac{p_i(y_i, x_i)}{m_{i+1} \cdots m_n} \\ &= \pi(y)p(y, x).\end{aligned}$$

In fact, the terms corresponding to an index $i < k$ satisfy $p_i(x_i, y_i) = p_i(y_i, x_i)$ since $x_i = y_i$; the term corresponding to the index k satisfies $p_k(x_k, y_k) = p_k(y_k, x_k)$ since the equality

$$p_k(x_k, y_k) = p_k(y_k, x_k)$$

holds by hypothesis.

Now we want to prove that the condition $\sigma_k = \frac{1}{m_k}$, for every $k > i_1$, is also necessary. Suppose that the equality $\pi(x)p(x, y) = \pi(y)p(y, x)$ holds. By the hypothesis of irreducibility we can consider two elements $x^0, y^0 \in X_1 \times \cdots \times X_n$ such that $x_{i_1}^0 \neq y_{i_1}^0$ and with the property that $p_{i_1}(x_{i_1}^0, y_{i_1}^0) \neq 0$. Now we have

$$\begin{aligned}\pi(x^0)p(x^0, y^0) &= \pi(y^0)p(y^0, x^0) \quad \Leftrightarrow \\ \pi(x^0)p_{i_1}(x_{i_1}^0, y_{i_1}^0) &= \pi(y^0)p_{i_1}(y_{i_1}^0, x_{i_1}^0).\end{aligned}$$

This gives

$$\frac{\pi(x^0)}{\pi(y^0)} = \frac{p_{i_1}(y_{i_1}^0, x_{i_1}^0)}{p_{i_1}(x_{i_1}^0, y_{i_1}^0)} = \frac{\sigma_{i_1}(x_{i_1}^0)}{\sigma_{i_1}(y_{i_1}^0)}.$$

Consider now the element $x = (x_1^0, \dots, x_{i_1}^0, y_{i_1+1}^0, \dots, y_n^0)$. The equality $\pi(x)p(x, y^0) = \pi(y^0)p(y^0, x)$ implies

$$\frac{\pi(x)}{\pi(y^0)} = \frac{p_{i_1}(y_{i_1}^0, x_{i_1}^0)}{p_{i_1}(x_{i_1}^0, y_{i_1}^0)} = \frac{\sigma_{i_1}(x_{i_1}^0)}{\sigma_{i_1}(y_{i_1}^0)}.$$

So we get $\pi(x^0) = \pi(x)$, that is, the probability π does not depend on the coordinates $i_1 + 1, \dots, n$. Set now $x' = (x_1^0, \dots, x_{i_1}^0, \dots, x_{k-1}^0, x_k, \dots, x_n)$. The equality $\pi(x^0)p(x^0, x') = \pi(x')p(x', x^0)$ gives

$$\pi(x^0) \left(\sum_{j \in N, j \leq k} p_j^0(p_j(x_j^0, x'_j)) \right) = \pi(x') \left(\sum_{j \in N, j \leq k} p_j^0(p_j(x'_j, x_j^0)) \right).$$

Since the probability π does not depend on the coordinates $i_1 + 1, \dots, n$, we get $p_k(x_k^0, x'_k) = p_k(x'_k, x_k^0)$. This implies $\sigma_k(x'_k) = \sigma_k(x_k^0)$ and so the hypothesis of irreducibility guarantees that σ_k is uniform on X_k . This completes the proof. \square

THEOREM 4.3. *The eigenspaces of the operator P are given by:*

- $W^1 \otimes \dots \otimes W^{k-1} \otimes V_{j_k}^k \otimes V_0^{k+1} \otimes V_0^{k+2} \otimes \dots \otimes V_0^n$, with $j_k \neq 0$, for $k \in \{i_1 + 1, \dots, n\}$ and where

$$W^i = \begin{cases} L(X_i), & \text{if } i \in N, \\ V_{j_i}^i, & j_i = 0, \dots, r_i, \quad \text{if } i \in C, \end{cases}$$

with eigenvalue

$$\sum_{i \in C: i < k} p_i^0 \lambda_{j_i} + p_k^0 \lambda_{j_k} + \sum_{i > k} p_i^0.$$

- $V_{j_1}^1 \otimes \dots \otimes V_{j_{i_1-1}}^{i_1-1} \otimes V_{j_{i_1}}^{i_1} \otimes V_0^{i_1+1} \otimes \dots \otimes V_0^n$, with $j_t = 0, \dots, r_t$, for every $t = 1, \dots, i_1$, with eigenvalue

$$\sum_{i=1}^{i_1} p_i^0 \lambda_{j_i} + \sum_{i=i_1+1}^n p_i^0.$$

PROOF. Fix an index $k \in \{i_1 + 1, i_1 + 2, \dots, n\}$ and consider the function φ in the space

$$W^1 \otimes \dots \otimes W^{k-1} \otimes V_{j_k}^k \otimes V_0^{k+1} \otimes V_0^{k+2} \otimes \dots \otimes V_0^n,$$

with $j_k \neq 0$ and

$$W^i = \begin{cases} L(X_i), & \text{if } i \in N, \\ V_{j_i}^i, & j_i = 0, \dots, r_i, \quad \text{if } i \in C, \end{cases}$$

so that $\varphi = \varphi_1 \otimes \dots \otimes \varphi_{k-1} \otimes \varphi_k \otimes \varphi_{k+1} \otimes \dots \otimes \varphi_n$ with $\varphi_i \in W^i$ for $i = 1, \dots, k-1$, $\varphi_k \in V_{j_k}^k$ and $\varphi_l \in V_0^l$ for $l = k+1, \dots, n$. Set $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$; then

$$\begin{aligned} (P\varphi)(x) &= \sum_y p(x, y) \varphi(y) \\ &= \sum_y \left(\sum_{i \in C} p_i^0 \delta_1(x_1, y_1) \cdots \delta_{i-1}(x_{i-1}, y_{i-1}) \right. \\ &\quad \times p_i(x_i, y_i) \delta_{i+1}(x_{i+1}, y_{i+1}) \cdots \delta_n(x_n, y_n) \\ &\quad \left. + \sum_{i \in N} p_i^0 \delta_1(x_1, y_1) \cdots \delta_{i-1}(x_{i-1}, y_{i-1}) p_i(x_i, y_i) \frac{1}{m_{i+1}} \cdots \frac{1}{m_n} \right) \\ &\quad \times \varphi_1(y_1) \cdots \varphi_{k-1}(y_{k-1}) \varphi_k(y_k) \varphi_{k+1}(y_{k+1}) \cdots \varphi_n(y_n) \\ &= \sum_{i \in C, i \leq k} \left(\sum_{y_i} p_i^0 p_i(x_i, y_i) \varphi_i(y_i) \right) \end{aligned}$$

$$\begin{aligned}
& \times \varphi_1(x_1) \cdots \varphi_{i-1}(x_{i-1}) \varphi_{i+1}(x_{i+1}) \cdots \varphi_n(x_n) \\
& + \sum_{i \in C, i > k} \left(\sum_{y_i} p_i^0 p_i(x_i, y_i) \varphi_i(y_i) \right) \\
& \quad \times \varphi_1(x_1) \cdots \varphi_{i-1}(x_{i-1}) \varphi_{i+1}(x_{i+1}) \cdots \varphi_n(x_n) \\
& + \sum_{i \in N, i > k} \left(\sum_{y_i, \dots, y_n} p_i^0 p_i(x_i, y_i) \frac{1}{m_{i+1}} \cdots \frac{1}{m_n} \varphi_i(y_i) \cdots \varphi_n(y_n) \right) \\
& \quad \times \varphi_1(x_1) \cdots \varphi_{i-1}(x_{i-1}) \\
& + \chi_N(k) \sum_{y_k, \dots, y_n} p_k^0 p_k(x_k, y_k) \frac{1}{m_{k+1}} \cdots \frac{1}{m_n} \\
& \quad \times \varphi_1(x_1) \cdots \varphi_{k-1}(x_{k-1}) \varphi_k(y_k) \cdots \varphi_n(y_n) \\
& = \sum_{i \in C, i \leq k} p_i^0 \lambda_{j_i} \varphi(x) + \sum_{i \in C, i > k} p_i^0 \cdot 1 \cdot \varphi(x) \\
& \quad + \sum_{i \in N, i > k} \left(\sum_{y_i} p_i^0 p_i(x_i, y_i) \varphi_i(y_i) \right) \\
& \quad \times \varphi_1(x_1) \cdots \varphi_{i-1}(x_{i-1}) \varphi_{i+1}(x_{i+1}) \cdots \varphi_n(x_n) \\
& \quad + \chi_N(k) \sum_{y_k} p_k^0 p_k(x_k, y_k) \\
& \quad \times \varphi_1(x_1) \cdots \varphi_{k-1}(x_{k-1}) \varphi_k(y_k) \varphi_{k+1}(x_{k+1}) \cdots \varphi_n(x_n) \\
& = \sum_{i \in C, i \leq k} p_i^0 \lambda_{j_i} \varphi(x) + \sum_{i \in C, i > k} p_i^0 \varphi(x) \\
& \quad + \sum_{i \in N, i > k} p_i^0 \varphi(x) + \chi_N(k) p_k^0 \lambda_{j_k} \varphi(x) \\
& = \left(\sum_{i \in C, i < k} p_i^0 \lambda_{j_i} + p_k^0 \lambda_{j_k} + \sum_{i > k} p_i^0 \right) \varphi(x),
\end{aligned}$$

where χ_N is the characteristic function of N . Note that in this case the addends corresponding to the indices $i < k$, $i \in N$, are equal to 0 since we have supposed $j_k \neq 0$.

Consider now the function φ in the space

$$V_{j_1}^1 \otimes \cdots \otimes V_{j_{i_1-1}}^{i_1-1} \otimes V_{j_{i_1}}^{i_1} \otimes V_0^{i_1+1} \otimes \cdots \otimes V_0^n,$$

with $j_t = 0, \dots, r_t$, for every $t = 1, \dots, i_1$. In this case we have

$$(P\varphi)(x) = \sum_y p(x, y) \varphi(y)$$

$$\begin{aligned}
&= \sum_{i \in C, i < i_1} \left(\sum_{y_i} p_i^0 p_i(x_i, y_i) \varphi_i(y_i) \right) \\
&\quad \times \varphi_1(x_1) \cdots \varphi_{i-1}(x_{i-1}) \varphi_{i+1}(x_{i+1}) \cdots \varphi_n(x_n) \\
&+ \sum_{i \in C, i > i_1} \left(\sum_{y_i} p_i^0 p_i(x_i, y_i) \varphi_i(y_i) \right) \\
&\quad \times \varphi_1(x_1) \cdots \varphi_{i-1}(x_{i-1}) \varphi_{i+1}(x_{i+1}) \cdots \varphi_n(x_n) \\
&+ \sum_{i \in N, i > i_1} \left(\sum_{y_i, \dots, y_n} p_i^0 p_i(x_i, y_i) \frac{1}{m_{i+1}} \cdots \frac{1}{m_n} \varphi_i(y_i) \cdots \varphi_n(x_n) \right) \\
&\quad \times \varphi_1(x_1) \cdots \varphi_{i-1}(x_{i-1}) \\
&+ \sum_{y_{i_1}, \dots, y_n} \left(p_{i_1}^0 p_{i_1}(x_{i_1}, y_{i_1}) \frac{1}{m_{i_1+1}} \cdots \frac{1}{m_n} \varphi_{i_1}(y_{i_1}) \cdots \varphi_n(x_n) \right) \\
&\quad \times \varphi_1(x_1) \cdots \varphi_{i_1-1}(x_{i_1-1}) \\
&= \sum_{i \in C, i < i_1} p_i^0 \lambda_{j_i} \varphi(x) + \sum_{i \in C, i > i_1} p_i^0 \varphi(x) \\
&\quad + \sum_{i \in N, i > i_1} p_i^0 \varphi(x) + p_{i_1}^0 \lambda_{j_{i_1}} \varphi(x) \\
&= \left(\sum_{i=1}^{i_1} p_i^0 \lambda_{j_i} + \sum_{i=i_1+1}^n p_i^0 \right) \varphi(x).
\end{aligned}$$

Observe that, by computing the sum of the dimensions of these eigenspaces, we get

$$\sum_{k=i_1+1}^n m_1 \cdots m_{k-1} (m_k - 1) + m_1 m_2 \cdots m_{i_1} = m_1 m_2 \cdots m_n,$$

which is just the dimension of the space $X_1 \times \cdots \times X_n$. \square

REMARK 4.4. The expression of the eigenvalues of P given in the previous theorem tells us that if P_i is ergodic for every $i = 1, \dots, n$, then also P is ergodic, since the eigenvalue 1 is obtained with multiplicity 1 and the eigenvalue -1 can never be obtained.

We can now give the matrices U, D and Δ associated with P . For every i , let U_i, D_i and Δ_i be the matrices of eigenvectors, of the coefficients of σ_i and of eigenvalues for the probability P_i , respectively. The expression of the matrix U , whose columns are an orthonormal basis of eigenvectors for P , easily follows from Theorem 4.3. In order to get the diagonal matrix

D , whose entries are the coefficients of π , it suffices to consider the tensor product of the corresponding matrices associated with the probability P_i , for every $i = 1, \dots, n$, as it follows from Theorem 4.2. Finally, to get the matrix Δ of eigenvalues of P it suffices to replace, in the expression of the matrix P , the matrix P_i by Δ_i and the matrix J_i by the corresponding diagonal matrix J_i^{diag} , which has the eigenvalue 1 with multiplicity 1 and the eigenvalue 0 with multiplicity $m_i - 1$. So we have the following proposition.

PROPOSITION 4.5. *Let P be the crested product of the Markov chains P_i , with $i = 1, \dots, n$. Then we have:*

- $U = \sum_{k=i_1+1}^n M_1 \otimes \dots \otimes M_{k-1} \otimes (U_k - A_k) \otimes A_{k+1} \otimes \dots \otimes A_n + U_1 \otimes U_2 \otimes \dots \otimes U_{i_1} \otimes A_{i_1+1} \otimes \dots \otimes A_n$, with

$$M_i = \begin{cases} I_i^{\sigma_i\text{-norm}}, & \text{if } i \in N, \\ U_i, & \text{if } i \in C, \end{cases}$$

where

$$I_i^{\sigma_i\text{-norm}} = \begin{pmatrix} \frac{1}{\sqrt{\sigma_i(0)}} & & & \\ & \frac{1}{\sqrt{\sigma_i(1)}} & & \\ & & \ddots & \\ & & & \frac{1}{\sqrt{\sigma_i(m_i-1)}} \end{pmatrix}.$$

By A_i we denote the matrix of size m_i whose entries on the first column are all 1 and the remaining ones are 0:

- $D = \bigotimes_{i=1}^n D_i$.
- $\Delta = \sum_{i \in C} p_i^0 (I_1 \otimes \dots \otimes I_{i-1} \otimes \Delta_i \otimes I_{i+1} \otimes \dots \otimes I_n) + \sum_{i \in N} p_i^0 (I_1 \otimes \dots \otimes I_{i-1} \otimes \Delta_i \otimes J_{i+1}^{\text{diag}} \otimes \dots \otimes J_n^{\text{diag}})$.

Observe that another matrix U' of eigenvectors for P is given by $U' = \bigotimes_{i=1}^n U_i$. The matrix U that we have given above seems to be more useful whenever one wants to compute the k th step transition probability $p^{(k)}(0, x)$ using the formula (3), since it contains a greater number of zeros in the first row with respect to U' and so a small number of terms in the sum are nonzero.

Suppose $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$ elements in $X = X_1 \times \dots \times X_n$. From (3) and Proposition 4.5, we have

$$\begin{aligned} & p^{(k)}(x, y) \\ &= \pi(y) \left[\sum_{z \in X} \left(\sum_{r=i_1+1}^n m_1(x_1, z_1) \cdots m_{r-1}(x_{r-1}, z_{r-1}) (u_r - a_r)(x_r, z_r) \right) \right] \end{aligned}$$

$$\begin{aligned}
& \times a_{r+1}(x_{r+1}, z_{r+1}) \cdots a_n(x_n, z_n) \\
& + u_1(x_1, z_1) \cdots u_{i_1}(x_{i_1}, z_{i_1}) \\
& \quad \times a_{i_1+1}(x_{i_1+1}, z_{i_1+1}) \cdots a_n(x_n, z_n) \Big) \lambda_z^k \\
& \times \left(\sum_{r=i_1+1}^n m_1(y_1, z_1) \cdots m_{r-1}(y_{r-1}, z_{r-1}) (u_r - a_r)(y_r, z_r) \right. \\
& \quad \times a_{r+1}(y_{r+1}, z_{r+1}) \cdots a_n(y_n, z_n) \\
& \quad + u_1(y_1, z_1) \cdots u_{i_1}(y_{i_1}, z_{i_1}) \\
& \quad \left. \times a_{i_1+1}(y_{i_1+1}, z_{i_1+1}) \cdots a_n(y_n, z_n) \right),
\end{aligned}$$

where m_i, u_i, a_i are the probabilities associated with the matrices M_i, U_i, A_i occurring in Proposition 4.5.

5. The crossed product. The crossed product of the Markov chains P_i 's can be obtained as a particular case of the crested product, by setting $C = \{1, \dots, n\}$ (so that $N = \emptyset$) and it is also called the *direct* product. The analogous case for product of groups has been studied in [9].

In this case, we get the following transition probability:

$$p((x_1, \dots, x_n), (y_1, \dots, y_n)) = \sum_{i=1}^n p_i^0 \delta(x_1, y_1) \cdots p_i(x_i, y_i) \cdots \delta(x_n, y_n).$$

This corresponds to choosing the i th coordinate with probability p_i^0 and to changing it according with the probability transition P_i . So we get

$$p((x_1, \dots, x_n), (y_1, \dots, y_n)) = \begin{cases} p_i^0 p_i(x_i, y_i), & \text{if } x_j = y_j \text{ for all } j \neq i, \\ 0, & \text{otherwise.} \end{cases}$$

So, for $X_1 = \dots = X_n =: X$ and $p_0^1 = \dots = p_n^0 = \frac{1}{n}$, the probability p defines an analogue of the Ehrenfest model, where n is the number of balls and $|X| = m$ is the number of urns. In order to obtain a new configuration, we choose a ball with probability $1/n$ (let it be the i th ball in the urn x_i) and with probability $p_i(x_i, y_i)$ we put it in the urn y_i .

As a consequence of Theorem 4.2, we get that if P_i is in detailed balance with π_i , then P is in detailed balance with the strict probability measure π on $X_1 \times \dots \times X_n$ defined as

$$\pi(x_1, \dots, x_n) = \pi_1(x_1) \pi_2(x_2) \cdots \pi_n(x_n).$$

The matrix P associated with the probability p is given by

$$(4) \quad P = \sum_{i=1}^n p_i^0 (I_1 \otimes \cdots \otimes P_i \otimes \cdots \otimes I_n).$$

The following proposition studies the spectral theory of the operator P and it is a straightforward consequence of Theorem 4.3.

PROPOSITION 5.1. *Let $\varphi_0^i = 1_{X_i}, \varphi_1^i, \dots, \varphi_{m_i-1}^i$ be the eigenfunctions of P_i associated with the eigenvalues $\lambda_0^i = 1, \lambda_1^i, \dots, \lambda_{m_i-1}^i$, respectively. Then the eigenvalues of P are the $m_1 m_2 \cdots m_n$ numbers*

$$\lambda_I = \sum_{k=1}^n p_k^0 \lambda_{i_k}^k,$$

with $I = (i_1, \dots, i_n) \in \{0, \dots, m_1 - 1\} \times \cdots \times \{0, \dots, m_n - 1\}$. The corresponding eigenfunctions are defined as

$$\varphi_I((x_1, \dots, x_n)) = \varphi_{i_1}^1(x_1) \cdots \varphi_{i_n}^n(x_n).$$

As a consequence of Proposition 4.5, in order to get the matrices U, D and Δ associated with P , it suffices to consider the tensor product of the corresponding matrices associated with the probability P_i , for every $i = 1, \dots, n$. For every i , let U_i, D_i and Δ_i be the matrices of eigenvectors, of the coefficients of π_i and of eigenvalues for the probability P_i , respectively. We have the following corollary.

COROLLARY 5.2. *Let P be the probability defined in (4). Then we have*

$$\begin{aligned} PU &= U\Delta, \\ U^T DU &= I, \end{aligned}$$

where $U = \bigotimes_{i=1}^n U_i$, $\Delta = \bigotimes_{i=1}^n \Delta_i$ and $D = \bigotimes_{i=1}^n D_i$.

In particular, we can express the k th step transition probability matrix as

$$P^k = \left(\bigotimes_{i=1}^n U_i \right) \left(\bigotimes_{i=1}^n \Delta_i \right)^k \left(\bigotimes_{i=1}^n U_i \right)^T \left(\bigotimes_{i=1}^n D_i \right).$$

Let $x = (x_1, \dots, x_n)$ and $y = (y_1, \dots, y_n)$. Then we get

$$\begin{aligned} p^{(k)}(x, y) &= \pi(y) \sum_I \varphi_I(x) \lambda_I^k \varphi_I(y) \\ &= \pi_1(y_1) \cdots \pi_n(y_n) \\ &\quad \times \sum_I \varphi_{i_1}^1(x_1) \cdots \varphi_{i_n}^n(x_n) (p_1^0 \lambda_{i_1}^1 + \cdots + p_n^0 \lambda_{i_n}^n)^k \varphi_{i_1}^1(y_1) \cdots \varphi_{i_n}^n(y_n), \end{aligned}$$

with $I = (i_1, \dots, i_n)$.

As we said in Remark 4.4, if the matrix P_i is ergodic for every $i = 1, \dots, n$, then also the matrix P is ergodic, since the eigenvalue 1 can be obtained only by choosing $I = 0^n$ and the eigenvalue -1 can never be obtained.

REMARK 5.3. Put $n = 1$ and set $X = \{0, 1, \dots, m-1\}$. Consider the action of the symmetric group S_m on X . The stabilizer of a fixed element $x_0 = 0$ is isomorphic to the symmetric group S_{m-1} . It is well known (see [5]) that (S_m, S_{m-1}) is a Gelfand pair and the following decomposition of $L(X)$ into irreducible representations holds:

$$L(X) = V_0 \oplus V_1,$$

where $V_0 \cong \mathbb{C}$ is the space of constant functions on X and $V_1 = \{f : X \rightarrow \mathbb{C} : \sum_{i=0}^{m-1} f(i) = 0\}$. So we have $\dim V_0 = 1$ and $\dim V_1 = m - 1$.

Analogously, one can consider the action of the wreath product $S_m \wr S_n$ on $X^n = X \times \dots \times X$, defined in the natural way, and then one can study the decomposition of $L(X^n)$. We have

$$L(X^n) \cong L(X)^{\otimes n} \cong \bigoplus_{j=0}^n W_j,$$

with

$$W_j = \bigoplus_{w(i_1, \dots, i_n) = j} V_{i_1} \otimes V_{i_2} \otimes \dots \otimes V_{i_n},$$

where $w(i_1, \dots, i_n) = \#\{k : i_k = 1\}$. So we have $\dim W_j = \binom{n}{j} (m-1)^j$.

If we define on X the uniform transition probability, that is, $p_u(x, y) = \frac{1}{m}$ for all $x, y \in X$, then the matrix P_u is the matrix J of size m .

The eigenvalues of this matrix are 1 (with multiplicity 1) and 0 (with multiplicity $m-1$). The corresponding eigenspaces in $L(X)$ are, respectively, $V_0 \cong \mathbb{C}$ and $V_1 = \{f : X \rightarrow \mathbb{C} : \sum_{i=0}^{m-1} f(i) = 0\}$.

This means that, by choosing the uniform transition probability on X , one gets again the results obtained by considering the Gelfand pair (S_m, S_{m-1}) .

Also in the case of X^n we can find again the results obtained (see [5]) by considering the Gelfand pair $(S_m \wr S_n, S_{m-1} \wr S_n)$. For $P_u = J$ we have $\lambda_0 = 1$, $\lambda_1 = \dots = \lambda_{m-1} = 0$. Consider now the transition probability on X^n defined in (4), with $p_1^0 = \dots = p_n^0 = \frac{1}{n}$. The eigenfunctions φ_I associated with the eigenvalue $\frac{1}{n}(n-j)$, with $j = 0, \dots, n$, are in number of $\binom{n}{j} (m-1)^j$. Moreover

$$\sum_{j=0}^n \binom{n}{j} (m-1)^j = \sum_{j=0}^n \binom{n}{j} (m-1)^j 1^{n-j} = m^n = \dim L(X^n).$$

For every $j = 0, \dots, n$, these functions belong to W_j and they are a basis for W_j . So W_j is the eigenspace associated with the eigenvalue $\frac{1}{n}(n-j)$.

More generally, consider the case of any reversible transition probability p on X . Let $\lambda_0 = 1$, $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of P and $V_0 \cong \mathbb{C}$, V_1, \dots, V_k the corresponding eigenspaces. We get

$$L(X^n) \cong (V_0 \oplus V_1 \oplus \dots \oplus V_k)^{\otimes n}.$$

The eigenfunctions φ_I associated with the eigenvalue $\frac{1}{n} \sum_{i=0}^k r_i \lambda_i$, with $\sum_{i=0}^k r_i = n$, are

$$\binom{r_0 + r_1 + \cdots + r_k}{r_0, \dots, r_k} \prod_{i=0}^k (\dim V_i)^{r_i}$$

and the corresponding eigenspaces are the tensor products in which r_i copies of V_i , for $i = 0, 1, \dots, k$, appear. Moreover, the number of different eigenspaces is equal to the number of integer solutions of the equation

$$r_0 + r_1 + \cdots + r_k = n, \quad r_i \geq 0,$$

so it is $\binom{k+n}{n}$.

The definition of multinomial coefficient as $\binom{r_0+r_1+\cdots+r_k}{r_0, \dots, r_k} = \frac{(r_0+\cdots+r_k)!}{r_0!r_1!\cdots r_k!}$ guarantees that

$$\begin{aligned} \sum_{r_0+\cdots+r_k=n} \binom{n}{r_0, \dots, r_k} (\dim V_0)^{r_0} \cdots (\dim V_k)^{r_k} &= (\dim V_0 + \cdots + \dim V_k)^n \\ &= m^n, \end{aligned}$$

as we wanted.

6. The nested product.

6.1. *General properties.* The nested product of the Markov chains P_i 's can be obtained as a particular case of the crested product, by setting $N = \{1, \dots, n\}$ (so that $C = \emptyset$). The term *nested* comes from the association schemes theory (see [2]).

Consider the product

$$X_1 \times \cdots \times X_n$$

and let P_i be a transition probability on X_i . We assume that p_i is in detailed balance with the strict probability measure π_i , for all $i = 1, \dots, n$.

The formula of crested product becomes, in this case,

$$(5) \quad P = \sum_{i=1}^n p_i^0 (I_1 \otimes \cdots \otimes P_i \otimes J_{i+1} \otimes J_{i+2} \otimes \cdots \otimes J_n).$$

Theorem 4.2 tells us that P is reversible if and only if P_k is symmetric, for every $k > 1$, that is, $\pi_i = \frac{1}{m_i}$ for every $i = 2, \dots, n$. In this case, P is in detailed balance with the strict probability measure π on $X_1 \times \cdots \times X_n$ given by

$$\pi(x_1, \dots, x_n) = \frac{\pi_1(x_1)}{\prod_{i=2}^n m_i}.$$

So let us assume π_i to be uniform for every $i = 2, \dots, n$. The transition probability associated with P is

$$\begin{aligned} p((x_1, \dots, x_n), (y_1, \dots, y_n)) \\ = \frac{p_1^0 p_1(x_1, y_1)}{m_2 m_3 \cdots m_n} + \sum_{j=2}^{n-1} \frac{\delta((x_1, \dots, x_{j-1}), (y_1, \dots, y_{j-1})) p_j^0 p_j(x_j, y_j)}{m_{j+1} \cdots m_n} \\ + \delta((x_1, \dots, x_{n-1}), (y_1, \dots, y_{n-1})) p_n^0 p_n(x_n, y_n). \end{aligned}$$

As we did in the case of the crossed product, we want to study the spectral theory of the operator P defined in (5).

Let

$$L(X_i) = \bigoplus_{k_i=0}^{r_i} W_{k_i}^i$$

be the spectral decomposition of $L(X_i)$, for all $i = 1, \dots, n$ and let $\lambda_0^i = 1$, $\lambda_1^i, \dots, \lambda_{r_i}^i$ be the distinct eigenvalues of P_i associated with these eigenspaces. From Theorem 4.3 we get the following proposition.

PROPOSITION 6.1. *The eigenspaces of $L(X_1 \times \cdots \times X_n)$ are:*

- $L(X_1) \otimes \cdots \otimes L(X_{n-1}) \otimes W_{k_n}^n$, of eigenvalue $p_n^0 \lambda_{k_n}^n$, for $k_n = 1, \dots, r_n$, with multiplicity $m_1 \cdots m_{n-1} \dim(W_{k_n}^n)$;
- $L(X_1) \otimes \cdots \otimes L(X_j) \otimes W_{k_{j+1}}^{j+1} \otimes W_0^{j+2} \otimes \cdots \otimes W_0^n$, of eigenvalue $p_{j+1}^0 \lambda_{k_{j+1}}^{j+1} + p_{j+2}^0 + \cdots + p_n^0$, with $k_{j+1} = 1, \dots, r_{j+1}$ and for $j = 1, \dots, n-2$, with multiplicity $m_1 \cdots m_j \dim(W_{k_{j+1}}^{j+1})$;
- $W_{k_1}^1 \otimes W_0^2 \otimes \cdots \otimes W_0^n$, of eigenvalue $p_1^0 \lambda_{k_1}^1 + p_2^0 + \cdots + p_n^0$, for $k_1 = 0, 1, \dots, r_1$, with multiplicity $\dim(W_{k_1}^1)$.

Moreover, as in the general case, one can verify that, under the hypothesis that the operators P_i are ergodic, for $i = 1, \dots, n$, then also the operator P is ergodic.

The application of Proposition 4.5 to the case of the nested product yields the following corollary.

COROLLARY 6.2. *Let P be the nested product of the probabilities P_i , with $i = 1, \dots, n$. Then we have:*

- $U = U_1 \otimes A_2 \otimes \cdots \otimes A_n + \sum_{k=2}^n I_1^{\sigma_1\text{-norm}} \otimes \cdots \otimes I_{k-1}^{\sigma_{k-1}\text{-norm}} \otimes (U_k - A_k) \otimes A_{k+1} \otimes \cdots \otimes A_n$;

- $D = \bigotimes_{i=1}^n D_i$;
- $\Delta = \sum_{i=1}^n p_i^0 (I_1 \otimes \cdots \otimes I_{i-1} \otimes \Delta_i \otimes J_{i+1}^{\text{diag}} \otimes \cdots \otimes J_n^{\text{diag}})$.

6.2. *k-steps transition probability.* The formula that describes the transition probability after k steps in the case of the nested product can be simplified using the base of eigenvectors given in Corollary 6.2 and supposing that the starting point is $0 = (0, \dots, 0)$.

From the general formula, with the usual notation, we get

$$\begin{aligned}
& p^{(k)}(0, y) \\
&= \pi(y) \left[\sum_{z \in X} \left(\sum_{r=2}^n \delta_{\sigma_1}(0, z_1) \cdots \delta_{\sigma_{r-1}}(0, z_{r-1}) (u_r - a_r)(0, z_r) \right. \right. \\
&\quad \times a_{r+1}(0, z_{r+1}) \cdots a_n(0, z_n) \\
&\quad \left. \left. + u_1(0, z_1) a_2(0, z_2) \cdots a_n(0, z_n) \right) \lambda_z^k \right. \\
&\quad \times \left(\sum_{r=2}^n \delta_{\sigma_1}(y_1, z_1) \cdots \delta_{\sigma_{r-1}}(y_{r-1}, z_{r-1}) (u_r - a_r)(y_r, z_r) \right. \\
&\quad \times a_{r+1}(y_{r+1}, z_{r+1}) \cdots a_n(y_n, z_n) \\
&\quad \left. \left. + u_1(y_1, z_1) a_2(y_2, z_2) \cdots a_n(y_n, z_n) \right) \right] \\
&= \pi(y) \left[1 + \sum_{j=2}^n \sum_{\substack{z_j \neq 0 \\ z_i=0, i \neq j}} \left(\sum_{r=j}^n \frac{1}{\sqrt{\sigma_1(0)} \cdots \sqrt{\sigma_{r-1}(0)}} (u_r - a_r)(0, z_r) \right) \right. \\
&\quad \times \left(p_r^0 \lambda_{z_r}^r + \sum_{m>r} p_m^0 \right)^k \\
&\quad \times \left(\sum_{r=j}^n \delta_{\sigma_1}(y_1, 0) \delta_{\sigma_2}(y_2, z_2) \cdots \right. \\
&\quad \times \delta_{\sigma_{r-1}}(y_{r-1}, z_{r-1}) (u_r - a_r)(y_r, z_r) \\
&\quad \left. \times a_{r+1}(y_{r+1}, z_{r+1}) \cdots a_n(y_n, z_n) \right) \\
&\quad \left. + \sum_{\substack{z_1 \neq 0 \\ z_i=0, i>1}} u_1(0, z_1) \left(p_1^0 \lambda_{z_1}^1 + \sum_{m=2}^n p_m^0 \right)^k u_1(y_1, z_1) \right].
\end{aligned}$$

Observe that in this case the sum consists of no more than

$$|X_1| + \sum_{i=2}^n (|X_i| - 1) = \sum_{i=1}^n |X_i| - n + 1$$

nonzero terms.

EXAMPLE 6.3. We want to express the k th step transition probability in the case $n = 2$. So consider the product $X \times Y$, with $X = \{0, 1, \dots, m\}$ and $Y = \{0, 1, \dots, n\}$. Let

$$L(X) = \bigoplus_{j=0}^r V_j \quad \text{and} \quad L(Y) = \bigoplus_{i=0}^s W_i$$

be the spectral decomposition of the spaces $L(X)$ and $L(Y)$, respectively. Let $\lambda_0 = 1, \lambda_1, \dots, \lambda_r$ and $\mu_0 = 1, \mu_1, \dots, \mu_s$ be the distinct eigenvalues of P_X and P_Y , respectively. Then the eigenspaces of $L(X \times Y)$ are $L(X) \otimes W_i$, for $i = 1, \dots, s$, with dimension $(m+1) \dim W_i$ and associated eigenvalue $p_Y^0 \mu_i$, and $V_j \otimes W_0$, for $j = 0, \dots, r$, with dimension $\dim V_j$ and associated eigenvalue $p_X^0 \lambda_j + p_Y^0$.

The expression of U becomes

$$U = I_X^{\sigma_X - \text{norm}} \otimes (U_Y - A_Y) + U_X \otimes A_Y.$$

In particular, let $\{v^0, v_1^1, \dots, v_{\dim(V_1)}^1, \dots, v_1^r, \dots, v_{\dim(V_r)}^r\}$ and $\{w^0, w_1^1, \dots, w_{\dim(W_1)}^1, \dots, w_1^s, \dots, w_{\dim(W_s)}^s\}$ be the eigenvectors of P_X and P_Y , respectively, that is, they represent the columns of the matrices U_X and U_Y .

Then, the columns of the matrix U corresponding to the elements $(i, 0) \in \{0, \dots, m\} \times \{0, \dots, n\}$ are the eigenvectors $v^i \otimes (1, \dots, 1)$ with eigenvalue $p_X^0 \lambda_i + p_Y^0$. On the other hand, the columns corresponding to the elements $(i, j) \in \{0, \dots, m\} \times \{0, \dots, n\}$, with $j = 1, \dots, n$, are the eigenvectors

$$\left(0, \dots, 0, \underbrace{\frac{1}{\sqrt{\sigma_X(i)}}}_{\text{ith place}}, 0, \dots, 0 \right) \otimes w^j$$

whose eigenvalue is $p_Y^0 \mu_j$. As a consequence, only $m+1+n$ of these eigenvectors can be nonzero in the first coordinate, so the probability $p^{(k)}((0, 0), (x, y))$ can be expressed as a sum of $m+1+n$ nonzero terms; moreover, these terms become $m+1$ if $x \neq 0$. We have

$$\begin{aligned} & p^{(k)}((0, 0), (x, y)) \\ &= \pi((x, y)) \left(\sum_{i=0}^m v^i(0) v^i(x) (p_X^0 \lambda_i + p_Y^0)^k \right. \\ & \quad \left. + \frac{1}{\sqrt{\sigma_X(0) \sigma_X(x)}} \sum_{j=1}^n w^j(0) \delta_0(x) w^j(y) (p_Y^0 \mu_j)^k \right) \end{aligned}$$

$$\begin{aligned}
&= \frac{\sigma_X(x)}{n+1} \left[\sum_{i=0}^r \left(\sum_{a=1}^{\dim(V_i)} v_a^i(0) v_a^i(x) \right) (p_X^0 \lambda_i + p_Y^0)^k \right. \\
&\quad \left. + \sum_{j=1}^s \left(\frac{1}{\sqrt{\sigma_X(0) \sigma_X(x)}} \sum_{b=1}^{\dim(W_j)} w_b^j(0) \delta_0(x) w_b^j(y) \right) (p_Y^0 \mu_j)^k \right].
\end{aligned}$$

6.3. *The insect.* It is clear that the product $X_1 \times \cdots \times X_n$ can be regarded as the rooted tree T of depth n , such that the root has degree m_1 , each vertex of the first level has m_2 children and in general each vertex of the i th level of the tree has m_{i+1} children, for every $i = 1, \dots, n-1$. We denote the i th level of the tree by L_i . In this way, every vertex $x \in L_i$ can be regarded as a word $x = x_1 \cdots x_i$, where $x_j \in \{0, 1, \dots, m_j - 1\}$.

We want to show that the nested product of Markov chains is the generalization of the ‘‘insect problem’’ studied by Figà-Talamanca in [11].

Let us imagine that an insect lives in a leaf $x \in L_n$ and that it performs a simple random walk on the graph T starting from x .

Then there exists a probability distribution μ_x on L_n such that, for every $y \in L_n$, $\mu_x(y)$ is the probability that y is the first point in L_n visited by the insect in the random walk. If we put $\bar{p}(x, y) = \mu_x(y)$, then we get a stochastic matrix $\bar{P} = (\bar{p}(x, y))_{x, y \in L_n}$. Since the random walk is $\text{Aut}(T)$ -invariant, we can suppose that the random walk starts at the leftmost vertex, that we will call $x_0 = (0, \dots, 0)$. We recall that $\text{Aut}(T)$ is the group of all automorphisms of T , given by the iterated wreath product $S_{m_n} \wr S_{m_{n-1}} \wr \cdots \wr S_{m_1}$. We want to study this Markov chain defined on L_n .

Set $\xi_n = \emptyset$ and $\xi_i = 00 \dots 0$ ($n-i$ times). For $j \geq 0$, let α_j be the probability that the insect reaches ξ_{j+1} given that ξ_j is reached at least once. This definition implies $\alpha_0 = 1$ and $\alpha_1 = \frac{1}{m_{n+1}}$. In fact, with probability 1, the insect reaches the vertex ξ_1 at the first step and, starting from ξ_1 , with probability $\frac{1}{m_{n+1}}$ it reaches ξ_2 , while with probability $\frac{m_n}{m_{n+1}}$ it returns to L_n . Finally, we have $\alpha_n = 0$. For $1 < j < n$, there is the following recursive relation:

$$\alpha_j = \frac{1}{m_{n+1-j} + 1} + \alpha_{j-1} \alpha_j \frac{m_{n+1-j}}{m_{n+1-j} + 1}.$$

In fact, starting at ξ_j , with probability $\frac{1}{m_{n+1-j} + 1}$ the insect reaches in one step ξ_{j+1} , otherwise with probability $\frac{m_{n+1-j}}{m_{n+1-j} + 1}$ it reaches ξ_{j-1} or one of its brothers; then, with probability α_{j-1} it reaches again ξ_j and one starts the recursive argument.

The solution, for $1 \leq j \leq n-1$, is given by

$$\alpha_j = \frac{1 + m_n + m_n m_{n-1} + m_n m_{n-1} m_{n-2} + \cdots + m_n m_{n-1} m_{n-2} \cdots m_{n-j+2}}{1 + m_n + m_n m_{n-1} + m_n m_{n-1} m_{n-2} + \cdots + m_n m_{n-1} m_{n-2} \cdots m_{n-j+1}}$$

$$= 1 - \frac{m_n m_{n-1} m_{n-2} \cdots m_{n-j+1}}{1 + m_n + m_n m_{n-1} + m_n m_{n-1} m_{n-2} + \cdots + m_n m_{n-1} m_{n-2} \cdots m_{n-j+1}}.$$

Moreover, we have

$$\begin{aligned} \bar{p}(x_0, x_0) &= \frac{1}{m_n}(1 - \alpha_1) + \frac{1}{m_n m_{n-1}} \alpha_1 (1 - \alpha_2) + \cdots \\ &\quad + \frac{1}{m_n m_{n-1} \cdots m_2} \alpha_1 \alpha_2 \cdots \alpha_{n-2} (1 - \alpha_{n-1}) + \frac{1}{m_n \cdots m_1} \alpha_1 \cdots \alpha_{n-1}. \end{aligned}$$

Indeed the j th summand is the probability of returning back to x_0 if the corresponding random walk in T reaches ξ_j but not ξ_{j+1} . It is not difficult to compute $\bar{p}(x_0, x)$, where x is a point at distance j from x_0 . For $j = 1$, we clearly have $\bar{p}(x_0, x_0) = \bar{p}(x_0, x)$. We observe that, for $j > 1$, to reach x one is forced to first reach ξ_j , so that we have

$$\begin{aligned} \bar{p}(x_0, x) &= \frac{1}{m_n \cdots m_{n-j+1}} \alpha_1 \alpha_2 \cdots \alpha_{j-1} (1 - \alpha_j) + \cdots \\ &\quad + \frac{1}{m_n \cdots m_2} \alpha_1 \alpha_2 \cdots \alpha_{n-2} (1 - \alpha_{n-1}) + \frac{1}{m_n \cdots m_1} \alpha_1 \alpha_2 \cdots \alpha_{n-1}. \end{aligned}$$

Since the random walk is invariant with respect to the action of $\text{Aut}(T)$, which acts isometrically on the tree, we get the same formula for any pair of vertices $x, y \in L_n$ such that $d(x, y) = j$.

PROPOSITION 6.4. *The stochastic matrix*

$$\begin{aligned} &p((x_1, \dots, x_n), (y_1, \dots, y_n)) \\ &= \frac{p_1^0 p_1(x_1, y_1)}{m_2 m_3 \cdots m_n} + \sum_{j=2}^{n-1} \frac{\delta((x_1, \dots, x_{j-1}), (y_1, \dots, y_{j-1})) p_j^0 p_j(x_j, y_j)}{m_{j+1} \cdots m_n} \\ &\quad + \delta((x_1, \dots, x_{n-1}), (y_1, \dots, y_{n-1})) p_n^0 p_n(x_n, y_n), \end{aligned}$$

defined in (5), gives rise to the Insect Markov chain on L_n , regarded as $X_1 \times \cdots \times X_n$, choosing $p_i^0 = \alpha_1 \alpha_2 \cdots \alpha_{n-i} (1 - \alpha_{n-i+1})$ for $i = 1, \dots, n-1$ and $p_n^0 = 1 - \alpha_1$ and the transitions probabilities p_j 's to be uniform for all $j = 1, \dots, n$.

PROOF. Set, for every $i = 1, \dots, n-1$,

$$p_i^0 = \alpha_1 \alpha_2 \cdots \alpha_{n-i} (1 - \alpha_{n-i+1})$$

and $p_n^0 = 1 - \alpha_1$. Moreover, assume that the probability p_i on X_i is uniform, that is,

$$P_i = J_i.$$

If $d(x_0, x) = n$, then we get

$$p(x_0, x) = \frac{\alpha_1 \alpha_2 \cdots \alpha_{n-1}}{m_1 m_2 \cdots m_n}.$$

If $d(x_0, x) = j > 1$, that is, $x_i^0 = x_i$ for all $i = 1, \dots, n - j$, then

$$p(x_0, x) = \frac{\alpha_1 \alpha_2 \cdots \alpha_{n-1}}{m_1 m_2 \cdots m_n} + \sum_{i=1}^{n-j} \frac{\alpha_1 \cdots \alpha_{n-i-1} (1 - \alpha_{n-i})}{m_n \cdots m_{i+2} m_{i+1}}.$$

Finally, if $x = x_0$, we get

$$p(x_0, x_0) = \frac{\alpha_1 \alpha_2 \cdots \alpha_{n-1}}{m_1 m_2 \cdots m_n} + \sum_{i=1}^{n-2} \frac{\alpha_1 \cdots \alpha_{n-i-1} (1 - \alpha_{n-i})}{m_n \cdots m_{i+2} m_{i+1}} + \frac{(1 - \alpha_1)}{m_n}.$$

This completes the proof. \square

The decomposition of the space $L(L_n) = L(X_1 \times \cdots \times X_n)$ under the action of $Aut(T)$ is known (see [6]). Denote $Z_0 \cong \mathbb{C}$ the trivial representation and, for every $j = 1, \dots, n$, define the following subspace:

$$Z_j = \left\{ f \in L(L_n) : f = f(x_1, \dots, x_j), \sum_{i=0}^{m_j-1} f(x_1, \dots, x_{j-1}, i) \equiv 0 \right\}$$

of dimension $m_1 \cdots m_{j-1} (m_j - 1)$. In virtue of the correspondence between $Aut(T)$ -invariant operators and bi- $Stab_{Aut(T)}(0^n)$ -invariant functions, the corresponding eigenvalues are given by the spherical Fourier transform of the convolver that represents \overline{P} , namely

$$\lambda_j = \sum_{x \in L_n} \overline{P}(x_0, x) \phi_j(x),$$

where ϕ_j is the j th spherical function, for all $j = 0, 1, \dots, n$. It is easy to verify that one gets:

- $\lambda_0 = 1$;
- $\lambda_j = 1 - \alpha_1 \alpha_2 \cdots \alpha_{n-j}$, for every $j = 1, \dots, n - 1$;
- $\lambda_n = 0$.

In particular, if we set

$$p_i^0 = \alpha_1 \alpha_2 \cdots \alpha_{n-i} (1 - \alpha_{n-i+1})$$

for every $i = 1, \dots, n - 1$, with $p_n^0 = 1 - \alpha_1$ and $P_i = J_i$ for every $i = 1, \dots, n$, the eigenspaces given for $L(X_1 \times \cdots \times X_n)$ in Proposition 6.1 are exactly the Z_j 's with the corresponding eigenvalues.

Let us prove that the eigenvalues that we have obtained in Proposition 6.1 coincide with the eigenvalues corresponding to the eigenspaces Z_0, Z_1, \dots, Z_n .

We want to get these eigenvalues by using the formulas given in Proposition 6.1 for the eigenvalues of the nested product P by setting $P_i = J_i$, then $p_i^0 = \alpha_1 \alpha_2 \cdots \alpha_{n-i} (1 - \alpha_{n-i+1})$ for $i = 1, \dots, n-1$ and $p_n^0 = 1 - \alpha_1$. First of all, we observe that the eigenvalues of the operator P_i are 1, with multiplicity 1 and 0, with multiplicity $m_i - 1$. So we get $L(X_i) = W_0^i \oplus W_1^i$, with $\dim(W_1^i) = m_i - 1$, for all $i = 1, \dots, n$. Following the formulas that we have given, the eigenspaces of P are:

- $L(X_1) \otimes L(X_2) \otimes \cdots \otimes L(X_{n-1}) \otimes W_1^n$;
- $L(X_1) \otimes L(X_2) \otimes \cdots \otimes L(X_{n-j-1}) \otimes W_1^{n-j} \otimes W_0^{n-j+1} \otimes \cdots \otimes W_0^n$, for every $j = 1, \dots, n-1$;
- $W_0^1 \otimes W_0^2 \otimes \cdots \otimes W_0^n$.

The corresponding eigenvalues are:

- $p_n^0 \lambda_1^n = 0$;
- $\sum_{i=n-j+1}^n p_i^0$, for every $j = 1, \dots, n-1$;
- $\sum_{i=1}^n p_i^0 = 1$.

We need to prove that, for every $j = 1, \dots, n-1$, the eigenvalue $\sum_{i=n-j+1}^n p_i^0$ is equal to $1 - \alpha_1 \alpha_2 \cdots \alpha_j$. We prove the assertion by induction on j .

If $j = 1$, we have $p_n^0 = 1 - \alpha_1$. Now suppose the assertion to be true for j and show that it holds also for $j+1$. We get

$$\begin{aligned} \sum_{i=n-j}^n p_i^0 &= \sum_{i=n-j+1}^n p_i^0 + p_{n-j}^0 = 1 - \alpha_1 \alpha_2 \cdots \alpha_j + \alpha_1 \cdots \alpha_j (1 - \alpha_{j+1}) \\ &= 1 - \alpha_1 \cdots \alpha_j \alpha_{j+1}. \end{aligned}$$

7. The second crested product. In this section we define a different kind of product of two spaces X and Y , that we will call the second crested product. In fact it contains, as particular cases, the crossed product and the nested product described in Sections 5 and 6, respectively. We will study a Markov chain P on the set Θ_k of functions from X to Y whose domains are k -subsets of X , giving the spectrum and the relative eigenspaces.

7.1. General theory. Let X be a finite set of cardinality n , say $X = \{1, 2, \dots, n\}$. For every $k = 1, \dots, n$, denote by Ω_k the set of k -subsets of X , so that $|\Omega_k| = \binom{n}{k}$.

Now let Y be a finite set and let Q be a transition matrix on Y , which is in detailed balance with the strict probability τ . Let $\lambda_0 = 1, \lambda_1, \dots, \lambda_m$ be the distinct eigenvalues of Q and denote by W_j the corresponding eigenspaces, for every $j = 0, 1, \dots, m$, so that the following spectral decomposition holds:

$$L(Y) = \bigoplus_{j=0}^m W_j.$$

Moreover, assume that the dimension of the eigenspace associated with the eigenvalue 1 is 1 and set $\dim(W_j) = m_j$, for every $j = 1, \dots, m$.

Recall that the eigenspace W_0 is generated by the vector

$$\underbrace{(1, \dots, 1)}_{|Y| \text{ times}}$$

and that W_j is orthogonal to W_0 with respect to the scalar product $\langle \cdot, \cdot \rangle_\tau$, for every $j = 1, \dots, m$.

For every $k = 1, \dots, n$, consider the space

$$\Theta_k = \{(A, \theta) : A \in \Omega_k \text{ and } \theta \in Y^A\},$$

that is, the space of functions θ whose domain $A = \text{dom}(\theta)$ is a k -subset of X and which take values in Y .

The set $\Theta = \coprod_{k=0}^n \Theta_k$ is a poset with respect to the relation \subseteq defined in the following way:

$$\varphi \subseteq \chi \text{ if } \text{dom}(\varphi) \subseteq \text{dom}(\chi) \text{ and } \varphi = \chi|_{\text{dom}(\varphi)}.$$

The Markov chain P on Θ_k that we are going to define can be regarded as follows. Let $0 < p_0 < 1$ a real number. Then, starting from a function $\theta \in \Theta_k$, with probability p_0 we can reach a function $\varphi \in \Theta_k$ having the same domain as θ and that can differ from θ at most in one image, according with the probability Q on Y .

On the other hand, with probability $1 - p_0$ we can reach in one step a function $\varphi \in \Theta_k$ whose domain intersects the domain of θ in $k - 1$ elements (on which the functions coincide), and in such a way that the image of the k th element of the domain of φ is uniformly chosen.

Note that P defines a Markovian operator on the space $L(\Theta_k)$ of all complex functions defined on Θ_k .

When Y is the ultrametric space, the Markov chain P represents the so-called multi-insect, which generalizes the Insect Markov chain already studied. In particular if $|X| = n$, we consider k insects living in k different subtrees and moving only one per each step in such a way that their distance is preserved, giving rise to a Markov chain on the space of all possible configurations of k insects having this property.

In fact each element in Θ_k can be regarded as a configuration of k insects and vice versa. For example, let $\theta \in \Theta_k$ be a function such that $\text{dom}(\theta) = \{x_1, \dots, x_k\}$ and $\theta(x_i) = y_i$, with $x_i \in X$ and $y_i \in Y$ for all $i = 1, \dots, k$. Then the corresponding configuration of k insects has an insect at each leaf (x_i, y_i) . They live in all different subtrees since $x_i \neq x_j$ for $i \neq j$.

We observe that the cardinality of this space is $\binom{n}{k}|Y|^k$. This space can be regarded as the variety of subtrees (see [4]) of branch indices $(k, 1)$ in the rooted tree $(n, |Y|)$.

If $\theta, \varphi \in \Theta_k$, with domains A and B respectively, then define the matrix Δ , indexed by Θ_k , whose entries are

$$\Delta_{\theta, \varphi} = \begin{cases} 1, & \text{if } |A \cap B| = k - 1 \text{ and } \theta|_{A \cap B} = \varphi|_{A \cap B}, \\ 0, & \text{otherwise.} \end{cases}$$

Observe that the matrix Δ is symmetric.

The operator P can be expressed in terms of the operator associated with Δ and of another operator M as

$$(6) \quad P = p_0 M + (1 - p_0) \frac{\Delta}{\text{norm}(\Delta)},$$

where M describes the situation in which the domain is not changed and only one of the images of the function $\theta \in \Theta_k$ is changed according with the probability Q on Y . An analytic expression for M will be presented below. On the other hand, Δ describes the situation in which we pass from a function whose domain is A to a function whose domain is $A \sqcup \{i\} \setminus \{j\}$, with $i \notin A$ and $j \in A$, and we choose uniformly the image in Y of the element i . So the action of Δ on Ω_k is an analogue of the Bernoulli–Laplace diffusion model. By $\text{norm}(\Delta)$ we indicate the number of nonzero entries in each row of the matrix associated with Δ .

It is easy to check that M is in detailed balance with the strict probability measure defined as

$$\tau_M(\theta) = \frac{1}{\binom{n}{k}} \prod_{i \in A} \tau(\theta(i)),$$

where $\theta \in \Theta_k$ and $\text{dom}(\theta) = A$. On the other hand, it follows from the definition of the Markov chain Δ that the weighted graph associated with Δ is connected. From this and from the fact that the nonzero entries of Δ are all equal to 1, we can deduce that Δ is reversible and in detailed balance with a uniform probability measure. This forces τ_M to be uniform and so we have to assume that τ is uniform on Y and the matrix Q is symmetric.

In this way, P is in detailed balance with the uniform probability measure π such that $\pi(\theta) = \frac{1}{\binom{n}{k} |Y|^k}$, for every $\theta \in \Theta_k$. This choice of τ guarantees that, if f is any function in W_j , with $j = 1, \dots, m$, then $\sum_{y \in Y} f(y) = 0$.

The spectral theory of the operator M has been studied in Section 5. In fact, it corresponds to choosing, with probability $\frac{1}{k}$, only one element of the domain and to changing the corresponding image with respect to the probability Q on Y , fixing the remaining ones. So we focus our attention to investigate the spectral theory of the operator Δ .

Let us introduce two differential operators.

DEFINITION 7.1. (1) For every $k = 2, \dots, n$ the operator $D_k : L(\Theta_k) \longrightarrow L(\Theta_{k-1})$ is defined by

$$(D_k F)(\varphi) = \sum_{\theta \in \Theta_k : \varphi \subseteq \theta} F(\theta),$$

for every $F \in L(\Theta_k)$ and $\varphi \in \Theta_{k-1}$.

(2) For $k = 1, \dots, n$ the operator $D_k^* : L(\Theta_{k-1}) \longrightarrow L(\Theta_k)$ is defined by

$$(D_k^* F)(\theta) = \sum_{\varphi \in \Theta_{k-1} : \varphi \subseteq \theta} F(\varphi),$$

for every $F \in L(\Theta_{k-1})$ and $\theta \in \Theta_k$.

Observe that the operator D_k^* is the adjoint of D_k . The following decomposition holds:

$$L(\Theta_k) = L\left(\prod_{A \in \Omega_k} Y^A\right) = \bigoplus_{A \in \Omega_k} L(Y^A).$$

In order to get a basis for the space $L(Y^A)$, for every $A \in \Omega_k$, we introduce some special functions that we will call fundamental functions.

DEFINITION 7.2. Suppose that $A \in \Omega_k$ and that $F^j \in L(Y)$ for every $j \in A$. Suppose also that each F^j belongs to an eigenspace of Q and set $a_i = |\{j \in A : F^j \in W_i\}|$. Then the tensor product $\bigotimes_{j \in A} F^j$ will be called a fundamental function of type $\underline{a} = (a_0, a_1, \dots, a_m)$ in $L(Y^A)$.

In other words, we have

$$\left(\bigotimes_{j \in A} F^j\right)(\theta) = \prod_{j \in A} F^j(\theta(j)),$$

for every $\theta \in Y^A$. We also set $\ell(\underline{a}) = a_1 + \dots + a_m = k - a_0$.

The introduction of the fundamental functions allows to give a useful expression for the operators M and Δ .

If $F \in L(Y^A) \subseteq L(\Theta_k)$ is the fundamental function $F = \bigotimes_{j \in A} F^j$, with $|A| = k$ and $F^j : Y \longrightarrow \mathbb{C}$, then $MF = \frac{1}{k} \sum_{j \in A} [(\bigotimes_{i \in A, i \neq j} F^i) \otimes QF^j]$. So, if $\theta \in \Theta_k$ and $\text{dom}(\theta) = A$, we get

$$(MF)(\theta) = \frac{1}{k} \sum_{j \in A} \left[\prod_{i \in A, i \neq j} F^i(\theta(i)) \left(\sum_{y \in Y} q(\theta(j), y) F^j(y) \right) \right].$$

Analogously one has $(\Delta F)(\theta) = \sum_{\varphi} F(\varphi)$, where the sum is over all $\varphi \in \Theta_k$ such that $\text{dom}(\varphi) \cap \text{dom}(\theta) = k - 1$ and $\varphi \equiv \theta$ on $\text{dom}(\varphi) \cap \text{dom}(\theta)$. If

$A = (\text{dom}(\theta) \cap A) \sqcup \{i\}$ (we denote by \sqcup the disjoint union), then

$$\left(\Delta \left(\bigotimes_{j \in A} F^j \right) \right) (\theta) = \sum_{\varphi} \bigotimes_{j \in A} F^j(\varphi) = \prod_{j \in \text{dom}(\varphi) \cap A} F^j(\theta(j)) \left(\sum_{y \in Y} F^i(y) \right).$$

Denote $P_{k, \underline{a}, A}$ the subspace of $L(Y^A)$ spanned by the fundamental functions of type \underline{a} and

$$P_{k, \underline{a}} = \bigoplus_{A \in \Omega_k} P_{k, \underline{a}, A}.$$

LEMMA 7.3. D_k maps $P_{k, \underline{a}}$ to $P_{k-1, \underline{a}'}$, where $\underline{a}' = (a_0 - 1, a_1, \dots, a_m)$. Conversely D_k^* maps $P_{k-1, \underline{a}'}$ to $P_{k, \underline{a}}$.

PROOF. Let F be a fundamental function of type \underline{a} in $L(Y^A)$ and let $B \subset A$ such that $A = B \sqcup \{i\}$. Then for every $\varphi \in Y^B$, we have

$$\begin{aligned} (D_k F)(\varphi) &= \sum_{\theta \in Y^A: \varphi \subseteq \theta} F(\theta) \\ &= \sum_{\theta \in Y^A: \varphi \subseteq \theta} \prod_{j \in A} F^j(\theta(j)) \\ &= \left(\sum_{y \in Y} F^i(y) \right) \prod_{j \in B} F^j(\varphi(j)). \end{aligned}$$

The value of $\sum_{y \in Y} F^i(y)$ is zero if $F^i \in W_j$ for $j = 1, \dots, m$ and so $D_k F \equiv 0$ if $a_0 = 0$. If $F^i \in W_0$, then $D_k F \in P_{k-1, \underline{a}'}$.

Analogously, let $F \in P_{k-1, \underline{a}', B}$ with $B \in \Omega_{k-1}$. Then for every $\theta \in Y^A$, $A = B \sqcup \{i\}$, one has

$$\begin{aligned} (D_k^* F)(\theta) &= \sum_{\varphi \in Y^B: \varphi \subseteq \theta} F(\varphi) \\ &= \prod_{j \in B} F^j(\varphi(j)) \\ &= F^i(\theta(i)) \prod_{j \in B} F^j(\theta(j)), \end{aligned}$$

where $F^i \equiv 1$ on Y (and so $F^i \in W_0$). \square

The restriction of D_k to $P_{k, \underline{a}}$ will be denoted by $D_{k, \underline{a}}$ and the restriction of D_k^* to $P_{k-1, \underline{a}'}$ by $D_{k, \underline{a}}^*$.

The study of the compositions of the operators $D_{k,\underline{a}}$ and $D_{k,\underline{a}}^*$ plays a central role. In fact it will be shown that the eigenspaces of these operators are also eigenspaces for Δ . Consider, for example, $D_{k+1}D_{k+1}^*$ applied to a function $F \in L(\Theta_k)$ and calculated on $\theta \in \Theta_k$. The functions $\varphi \in \Theta_{k+1}$ such that $\varphi \supseteq \theta$ are in number of $|Y|(n-k)$. Each of them covers $k+1$ functions in Θ_k ; one of them is the function θ , the other ones are functions in Θ_k whose domains differ by the domain of θ of an element and coincide on their intersection. These functions are in number of $|Y|(n-k)k$ and they correspond to functions that one can reach starting from θ in the Markov chain described by Δ . From this it follows that $norm(\Delta) = |Y|(n-k)k$.

LEMMA 7.4. *Let $F \in P_{k,\underline{a},A}$, with $A \in \Omega_k$. Then*

$$D_{k,\underline{a}}^*D_{k,\underline{a}} = |Y|(k - \ell(\underline{a}))I + Q_{k,\underline{a}},$$

where $Q_{k,\underline{a}}$ is defined by setting

$$(7) \quad (Q_{k,\underline{a}}F)(\theta) = \begin{cases} 0, & \text{if } F^i \notin W_0, \\ |Y|F(\bar{\theta}), & \text{if } F^i \in W_0, \end{cases}$$

for every $\theta \in \Theta_k$ such that $|dom(\theta) \cap A| = k-1$ and $A \setminus dom(\theta) = \{i\}$. We denote by $\bar{\theta}$ the function in Θ_k whose domain is A and such that $\bar{\theta}|_{A \setminus \{i\}} = \theta$ and $\bar{\theta}(i) = \theta(i_0)$, where $dom(\theta) \setminus A = \{i_0\}$.

PROOF. Take $F \in P_{k,\underline{a},A}$ and $\theta \in \Theta_k$. We have

$$\begin{aligned} (D_{k,\underline{a}}^*D_{k,\underline{a}}F)(\theta) &= \sum_{\varphi \in \Theta_{k-1}: \varphi \subseteq \theta} (D_{k,\underline{a}}F)(\varphi) \\ &= \sum_{\varphi \in \Theta_{k-1}: \varphi \subseteq \theta} \sum_{\substack{\omega \in \Theta_k: \omega \supseteq \varphi, \\ dom(\omega) = A}} F(\omega). \end{aligned}$$

If $dom(\theta) = A$, then we get

$$\begin{aligned} (D_{k,\underline{a}}^*D_{k,\underline{a}}F)(\theta) &= \sum_{j \in A} \left(\sum_{y \in Y} F^j(y) \right) \prod_{t \in A \setminus \{j\}} F^t(\theta(t)) \\ &= |Y|(k - \ell(\underline{a})) \prod_{t \in A} F^t(\theta(t)) \\ &= |Y|(k - \ell(\underline{a}))F(\theta), \end{aligned}$$

where the second equality follows from the fact that $\sum_{y \in Y} F^j(y) = |Y|$ if $F^j \in W_0$ and $\sum_{y \in Y} F^j(y) = 0$ whenever $F^j \notin W_0$.

On the other hand, if $|dom(\theta) \cap A| = k - 1$, with $A \setminus dom(\theta) = \{i\}$, then

$$\begin{aligned} (D_{k,\underline{a}}^* D_{k,\underline{a}} F)(\theta) &= \left(\sum_{y \in Y} F^i(y) \right) \prod_{j \in A \setminus \{i\}} F^j(\theta(j)) \\ &= \begin{cases} 0, & \text{if } F^i \notin W_0, \\ |Y|F(\bar{\theta}), & \text{if } F^i \in W_0, \end{cases} \end{aligned}$$

which is just the definition of $Q_{k,\underline{a}}$. \square

LEMMA 7.5. *Let $F \in P_{k,\underline{a}',A}$, with $A \in \Omega_k$. Then*

$$D_{k+1,\underline{a}} D_{k+1,\underline{a}}^* = |Y|(n-k)I + Q_{k,\underline{a}},$$

where $Q_{k,\underline{a}}$ is defined as in (7).

PROOF. Take $F \in P_{k,\underline{a}',A}$ and $\theta \in \Theta_k$. We have

$$\begin{aligned} (D_{k+1,\underline{a}} D_{k+1,\underline{a}}^* F)(\theta) &= \sum_{\varphi \in \Theta_{k+1}: \theta \subseteq \varphi} (D_{k+1,\underline{a}}^* F)(\varphi) \\ &= \sum_{\varphi \in \Theta_{k+1}: \theta \subseteq \varphi} \sum_{\substack{\omega \in \Theta_k: \omega \supseteq \varphi, \\ dom(\omega) = A}} F(\omega). \end{aligned}$$

If $dom(\theta) = A$, then we get

$$\begin{aligned} (D_{k+1,\underline{a}} D_{k+1,\underline{a}}^* F)(\theta) &= \sum_{j \in A^c} \sum_{y \in Y} F(\theta) \\ &= |Y|(n-k)F(\theta). \end{aligned}$$

On the other hand, if $|dom(\theta) \cap A| = k - 1$, with $A \setminus dom(\theta) = \{i\}$, then

$$\begin{aligned} (D_{k+1,\underline{a}} D_{k+1,\underline{a}}^* F)(\theta) &= \left(\sum_{y \in Y} F^i(y) \right) \prod_{j \in A \setminus \{i\}} F^j(\theta(j)) \\ &= \begin{cases} 0, & \text{if } F^i \notin W_0, \\ |Y|F(\bar{\theta}), & \text{if } F^i \in W_0, \end{cases} \\ &= (Q_{k,\underline{a}} F)(\theta). \end{aligned}$$

This completes the proof. \square

The following corollary easily follows.

COROLLARY 7.6. *Let $F \in P_{k,\underline{a}',A}$, with $A \in \Omega_k$. Then*

$$D_{k+1,\underline{a}} D_{k+1,\underline{a}}^* - D_{k,\underline{a}'}^* D_{k,\underline{a}'} = |Y|(n + \ell(\underline{a}) - 2k)I.$$

Consider now the operator $D_{k,\underline{a}}: P_{k,\underline{a}} \longrightarrow P_{k-1,\underline{a}'}$.

DEFINITION 7.7. For $0 \leq \ell(\underline{a}) \leq k \leq n$, set

$$P_{k,\underline{a},k} = \text{Ker}(D_{k,\underline{a}})$$

and inductively, for $k \leq h \leq n$, set

$$P_{h,\underline{a},k} = D_{h,\underline{a}}^* P_{h-1,\underline{a}',k}.$$

These spaces have a fundamental importance because they exactly constitute the eigenspaces of the operator Δ . This will be a consequence of the following proposition.

PROPOSITION 7.8. $P_{h,\underline{a}',k}$ is an eigenspace for the operator $D_{h+1,\underline{a}} D_{h+1,\underline{a}}^*$ and the corresponding eigenvalue is $|Y|(n + \ell(\underline{a}) - k - h)(h - k + 1)$.

PROOF. We prove the assertion by induction on h . If $h = k$, from the last corollary we get $D_{k+1,\underline{a}} D_{k+1,\underline{a}}^* |P_{k,\underline{a}',k} = |Y|(n + \ell(\underline{a}) - 2k)I$, since $D_{k,\underline{a}} P_{k,\underline{a}',k} = 0$ by definition of $P_{k,\underline{a}',k}$.

Now suppose the lemma to be true for $k \leq t \leq h$ and recall that, by definition, we have $P_{h+1,\underline{a}',k} = D_{h+1,\underline{a}'}^* P_{h,\underline{a}'',k}$. Moreover, Corollary 7.6 gives

$$D_{h+2,\underline{a}} D_{h+2,\underline{a}}^* - D_{h+1,\underline{a}'}^* D_{h+1,\underline{a}'} = |Y|(n + \ell(\underline{a}) - 2(h + 1))I.$$

So we get

$$\begin{aligned} D_{h+2,\underline{a}} D_{h+2,\underline{a}}^* |P_{h+1,\underline{a}',k} &= D_{h+1,\underline{a}'}^* |D_{h+1,\underline{a}'} D_{h+1,\underline{a}'}^* P_{h,\underline{a}'',k} \\ &\quad + |Y|(n + \ell(\underline{a}) - 2(h + 1))P_{h+1,\underline{a}',k} \\ &= |Y|(n + \ell(\underline{a}) - k - h)(h - k + 1)D_{h+1,\underline{a}'}^* P_{h,\underline{a}'',k} \\ &\quad + |Y|(n + \ell(\underline{a}) - 2(h + 1))P_{h+1,\underline{a}',k} \\ &= |Y|(n + \ell(\underline{a}) - k - h - 1)(h - k + 2)P_{h+1,\underline{a}',k}, \end{aligned}$$

where the second equality follows from the inductive hypothesis and the third one from an easy computation. This completes the proof. \square

COROLLARY 7.9. $P_{h,\underline{a}',k}$ is an eigenspace for Δ of eigenvalue $|Y|(n + \ell(\underline{a}) - k - h)(h - k + 1) - |Y|(n - h)$.

PROOF. It suffices to observe that the operator $Q_{h,\underline{a}}$ defined in (7) coincides with the operator Δ on the space $P_{h,\underline{a}}$ and then the assertion follows from Lemma 7.5 and Proposition 7.8. \square

In particular, after normalizing the matrix Δ we obtain $\frac{\Delta}{\text{norm}(\Delta)}$ and the corresponding eigenvalue is $\frac{1}{|Y|^{(n-h)h}}(|Y|(n + \ell(\underline{a}) - k - h)(h - k + 1) - |Y|(n - h))$.

The following lemma holds.

LEMMA 7.10. *Given $\ell(\underline{a})$ and h , then, for $\ell(\underline{a}) \leq k \leq \min\{h, \frac{n+\ell(\underline{a})}{2}\}$, the spaces $P_{h,\underline{a}',k}$ are mutually orthogonal.*

PROOF. Each $P_{h,\underline{a}',k}$ is an eigenspace for the self-adjoint operator $D_{h+1,\underline{a}} \times D_{h+1,\underline{a}}^*$. Since the eigenvalue $|Y|(n + \ell(\underline{a}) - k - h)(h - k + 1)$ is a strictly decreasing function of k for $k \leq \frac{n+\ell(\underline{a})}{2}$, then to different values of k correspond different eigenvalues. This proves the assertion. \square

Recall that, if $\underline{a} = (a_0, a_1, \dots, a_m)$, we set $\underline{a}' = (a_0 - 1, a_1, \dots, a_m)$ and, inductively, $\underline{a}^{h+1} = \underline{a}^h - (1, 0, \dots, 0)$.

PROPOSITION 7.11. *Let F be a function in $P_{k,\underline{a}^{h-k},k}$. Then, for $\ell(\underline{a}) \leq k \leq \frac{n+\ell(\underline{a})}{2}$ and $k \leq h \leq n + \ell(\underline{a}) - k$, we have*

$$\|D_{h,\underline{a}}^* D_{h-1,\underline{a}'}^* \cdots D_{k+1,\underline{a}^{h-k-1}}^* F\|^2 = \frac{(n + \ell(\underline{a}) - 2k)!(h - k)!}{(n + \ell(\underline{a}) - k - h)!} |Y|^{h-k} \|F\|^2.$$

In particular, $D_{h,\underline{a}}^* D_{h-1,\underline{a}'}^* \cdots D_{k+1,\underline{a}^{h-k-1}}^*$ is an isomorphism of $P_{k,\underline{a}^{h-k},k}$ onto $P_{h,\underline{a},k}$.

PROOF. We prove the assertion by induction on h . For $h = k + 1$ and $F \in P_{k,\underline{a}',k}$, we have

$$\begin{aligned} \|D_{k+1,\underline{a}}^* F\|^2 &= \langle D_{k+1,\underline{a}}^* F, D_{k+1,\underline{a}}^* F \rangle \\ &= \langle D_{k+1,\underline{a}} D_{k+1,\underline{a}}^* F, F \rangle \\ &= |Y|(n + \ell(\underline{a}) - 2k) \|F\|^2 \end{aligned}$$

by Proposition 7.8, so the assertion is true. For $h > k + 1$, applying Proposition 7.8 to $D_{h,\underline{a}} D_{h,\underline{a}}^*$, we get

$$\begin{aligned} \|D_{h,\underline{a}}^* D_{h-1,\underline{a}'}^* \cdots D_{k+1,\underline{a}^{h-k-1}}^* F\|^2 &= \langle D_{h,\underline{a}} D_{h,\underline{a}}^* D_{h-1,\underline{a}'}^* \cdots D_{k+1,\underline{a}^{h-k-1}}^* F, D_{h-1,\underline{a}'}^* \cdots D_{k+1,\underline{a}^{h-k-1}}^* F \rangle \\ &= |Y|(n + \ell(\underline{a}) - k - h + 1)(h - k) \|D_{h-1,\underline{a}'}^* \cdots D_{k+1,\underline{a}^{h-k-1}}^* F\|^2. \end{aligned}$$

Now the proposition follows by induction. \square

PROPOSITION 7.12. *Assume $\ell(\underline{a}) \leq h \leq \frac{n+\ell(\underline{a})}{2}$. Then:*

- (1) $P_{h,\underline{a}} = \bigoplus_{k=\ell(\underline{a})}^{\min\{h, n+\ell(\underline{a})-h\}} P_{h,\underline{a},k}$;
(2) $D_{h+1,\underline{a}}^* : P_{h,\underline{a}'} \longrightarrow P_{h+1,\underline{a}}$ is an injective map.

PROOF. We prove the assertion by induction on h .

Assume that (1) and (2) are true for $\ell(\underline{a}) - 1 \leq h \leq t \leq \frac{n+\ell(\underline{a})-1}{2}$. For $h = \ell(\underline{a}) - 1$ we have $P_{\ell(\underline{a})-1,\underline{a}} = 0$ and so the proposition trivially holds.

Since the operator $D_{h,\underline{a}}^*$ is the adjoint of $D_{h,\underline{a}}$ we have the following decomposition:

$$\begin{aligned} P_{h,\underline{a}} &= \text{Ker}(D_{h,\underline{a}}) \oplus D_{h,\underline{a}}^* P_{h-1,\underline{a}'} \\ &= P_{h,\underline{a},h} \oplus D_{h,\underline{a}}^* P_{h-1,\underline{a}'}. \end{aligned}$$

In particular

$$P_{t+1,\underline{a}} = P_{t+1,\underline{a},t+1} \oplus D_{t+1,\underline{a}}^* P_{t,\underline{a}'}.$$

By induction

$$P_{t,\underline{a}'} = \bigoplus_{k=\ell(\underline{a})}^t P_{t,\underline{a}',k}$$

and so

$$\begin{aligned} P_{t+1,\underline{a}} &= P_{t+1,\underline{a},t+1} \oplus D_{t+1,\underline{a}}^* \left(\bigoplus_{k=\ell(\underline{a})}^t P_{t,\underline{a}',k} \right) \\ &= \bigoplus_{k=\ell(\underline{a})}^{t+1} P_{t+1,\underline{a},k}. \end{aligned}$$

This proves (1), while (2) follows from (1) and Proposition 7.11. \square

COROLLARY 7.13. *The dimension of the spaces $P_{h,\underline{a},k}$ that appear in the decomposition of $P_{h,\underline{a}}$ is*

$$\frac{n + \ell(\underline{a}) + 1 - 2k}{n - k + 1} \binom{n}{k} \binom{k}{\ell(\underline{a})} \binom{\ell(\underline{a})}{a_1, \dots, a_m} \prod_{j=1}^m (\dim(W_j))^{a_j}.$$

PROOF. From the previous proposition it follows that

$$\dim(P_{t+1,\underline{a},t+1}) = \dim(P_{t+1,\underline{a}}) - \dim(P_{t,\underline{a}'}).$$

Now

$$\dim(P_{t+1,\underline{a}}) = \binom{n}{t+1} \binom{t+1}{a_0, a_1, \dots, a_m} \prod_{j=1}^m (\dim(W_j))^{a_j}.$$

In fact, $\binom{n}{t+1}$ represents the number of $(t+1)$ -subsets in X and $\binom{t+1}{a_0, a_1, \dots, a_m} \times \prod_{j=1}^m (\dim(W_j))^{a_j}$ represents the number of possible choices in the fundamental function $F = \prod_{r \in A} F^r$ of a_i functions belonging to the eigenspace W_i of $L(Y)$. Thus

$$\begin{aligned} \dim(P_{t+1, \underline{a}, t+1}) &= \binom{n}{t+1} \binom{t+1}{a_0, a_1, \dots, a_m} \prod_{j=1}^m (\dim(W_j))^{a_j} \\ &\quad - \binom{n}{t} \binom{t}{a_0-1, a_1, \dots, a_m} \prod_{j=1}^m (\dim(W_j))^{a_j} \\ &= \frac{n-t-a_0}{n-t} \binom{n}{t+1} \binom{t+1}{a_0, a_1, \dots, a_m} \prod_{j=1}^m (\dim(W_j))^{a_j}. \end{aligned}$$

Since, by Proposition 7.11, $\dim(P_{h, \underline{a}, k}) = \dim(P_{k, \underline{a}^{h-k}, k})$ one can obtain the result replacing t by $k-1$ and \underline{a} by \underline{a}^{h-k} . \square

We want to find now the eigenvector of $\frac{\Delta}{\text{norm}(\Delta)}$ associated with the eigenvalue 1. Consider in $P_{1, (1, 0, \dots, 0)}$ the function

$$f = \sum_{i=1}^n f_i,$$

where f_i is the fundamental function of type $(1, 0, \dots, 0)$ whose domain is $\{i\}$. Set

$$\langle f \rangle = P_{1, (1, 0, \dots, 0), 0} =: D_{1, (1, 0, \dots, 0)}^* P_{0, (0, \dots, 0), 0}.$$

So the element $F_0 = D_{h, (h, 0, \dots, 0)}^* \cdots D_{3, (3, 0, \dots, 0)}^* D_{2, (2, 0, \dots, 0)}^* f$ is the generator of the space $P_{h, (h, 0, \dots, 0), 0}$, which has dimension 1. Corollary 7.9 implies that $P_{h, (h, 0, \dots, 0), 0}$ is an eigenspace for $\frac{\Delta}{\text{norm}(\Delta)}$ and the corresponding eigenvalue is 1. Moreover, the connectedness of the graph associated with Δ implies that this is the unique (up to constant) eigenvector of eigenvalue 1. We denote by $P_{1, (1, 0, \dots, 0), 1}$ the orthogonal subspace to $P_{1, (1, 0, \dots, 0), 0}$ in $P_{1, (1, 0, \dots, 0)}$. It has dimension $n-1$.

Observe that the definition of fundamental functions is strictly linked to the spectral theory of the operator Q and so of the operator M restricted to each domain. In fact, if F is a fundamental function in $P_{h, \underline{a}, A}$, with $\underline{a} = (a_0, a_1, \dots, a_m)$ and $A \in \Omega_h$, then it is an eigenvector for the operator M and the corresponding eigenvalue is $\frac{1}{h} \sum_{j=0}^m a_j \lambda_j$. So the set of the eigenvalues of M is given by $\binom{n}{h}$ copies of these values. In particular, the eigenspace $P_{h, \underline{a}, k}$ of $\frac{\Delta}{\text{norm}(\Delta)}$ is also an eigenspace for M and an eigenvector in this space has eigenvalue $\frac{1}{h} \sum_{j=0}^m a_j \lambda_j$. So, by Corollary 7.9 and definition (6) of P , we get the following theorem.

THEOREM 7.14. $P_{h,\underline{a},k}$ is an eigenspace for P with eigenvalue

$$p_0 \cdot \frac{1}{h} \sum_{j=0}^m a_j \lambda_j + (1 - p_0) \frac{(n + \ell(\underline{a}) - k - h)(h - k + 1) - (n - h)}{h(n - h)}.$$

REMARK 7.15. It is easy to check that the operator M is not ergodic. In fact its associated graph contains $\binom{n}{h}$ connected components and so the multiplicity of the eigenvalue 1 for M is $\binom{n}{h}$.

On the other hand we already observed that the operator $\frac{\Delta}{\text{norm}(\Delta)}$ has the eigenvalue 1 with multiplicity 1. To conclude that it is ergodic it suffices to show that -1 is not an eigenvalue, that is, the associated graph is not bipartite. In fact consider $\theta \in \Theta_h$ with domain $\{i_1, \dots, i_h\}$ and $\theta(i_j) = y_j$, for every $j = 1, \dots, h$. By definition of Δ we can connect θ with φ , whose domain is $\{i_1, \dots, i_{h-1}, i_t\}$, $i_h \neq i_t$ and such that $\varphi(i_j) = y_j = \theta(i_j)$ for all $j = 1, \dots, h-1$ and $\varphi(i_t) = y_t$. Moreover θ can also be connected with ϱ whose domain is $\{i_1, \dots, i_{h-2}, i_h, i_t\}$ and such that $\varrho(i_j) = y_j = \theta(i_j)$ for all $j = 1, \dots, h-2, h$ and $\varrho(i_t) = y_t = \varphi(i_t)$. On the other hand φ and ϱ are connected as well and this proves that the graph is not bipartite.

From Proposition 3.4 we can deduce the ergodicity for the operator P , since the multiplicity of the eigenvalue 1 is 1 and the eigenvalue -1 does not appear in the spectrum of P .

REMARK 7.16. The second crested product reduces to the crossed product if $k = n$ and to the nested product if $k = 1$.

In fact, if $k = n$, the domain of a function $\theta \in \Theta_n$ cannot be changed and θ can be identified with the n -tuple $(y_1, \dots, y_n) \in Y^n$ of its images. The operator P becomes

$$P = \frac{1}{n} \sum_{i=1}^n I_1 \otimes \dots \otimes I_{i-1} \otimes Q \otimes I_{i+1} \otimes \dots \otimes I_n,$$

which is the crossed product on the space Y^n .

If $k = 1$, then Δ has the following expression:

$$\Delta = \begin{pmatrix} 0 & 1 & \dots & \dots & 1 \\ 1 & 0 & 1 & & \vdots \\ \vdots & 1 & \ddots & & \vdots \\ \vdots & & & \ddots & 1 \\ 1 & \dots & \dots & 1 & 0 \end{pmatrix}$$

and $\text{norm}(\Delta) = n - 1$. So we get

$$P = p_0(I_X \otimes Q) + (1 - p_0) \left(\frac{\Delta}{\text{norm}(\Delta)} \otimes J_Y \right),$$

which is just the nested product of X and Y , with $P_X = \frac{\Delta}{\text{norm}(\Delta)}$ and $P_Y = Q$.

7.2. *Bi-insect.* In what follows, we take Y as a homogeneous rooted tree of degree q and depth $m - 1$ and we give an explicit description of the spectrum of the operator $P = p_0M + (1 - p_0)\frac{\Delta}{\text{norm}(\Delta)}$ acting on the space $L(\Theta_2)$. Therefore we are considering functions in Θ_2 such that the image of each element of the domain is an insect. In other words, our aim is to diagonalize the bi-insect Markov chain defined in Section 2.2.2.

Suppose X to be a set of cardinality n and let $m \geq 3$. Recall that we have the decomposition

$$L(Y) = \bigoplus_{j=0}^{m-1} W_j,$$

where $W_0 \cong \mathbb{C}$ and

$$W_j = \left\{ f \in L(L_{m-1}) : f = f(x_1, \dots, x_j), \sum_{i=0}^{q-1} f(x_1, \dots, x_{j-1}, i) \equiv 0 \right\},$$

for every $j = 1, \dots, m - 1$. Observe that $\dim(W_j) = q^{j-1}(q - 1)$.

The eigenspaces relative to the operator $\Delta/\text{norm}(\Delta)$ are the subspaces of the form $P_{2,(a_0,a_1,\dots,a_{m-1}),k}$, with $k = 0, 1, 2$. The corresponding eigenvalue is

$$\frac{1}{q^{m-1}(n-2)2} [q^{m-1}(n + \ell(\underline{a}) - k - 2)(2 - k + 1) - q^{m-1}(n - 2)].$$

So, by dependence of $\ell(\underline{a})$, we get the following eigenspaces:

- $P_{2,(a_0,a_1,\dots,a_{m-1}),2} \begin{cases} a_0 = 0, & \text{with eigenvalue } \lambda = 0, \\ a_0 = 1, & \text{with eigenvalue } \lambda = -\frac{1}{2(n-2)}, \\ a_0 = 2, & \text{with eigenvalue } \lambda = -\frac{1}{n-2}, \end{cases}$
- $P_{2,(a_0,a_1,\dots,a_{m-1}),1} \begin{cases} a_0 = 1, & \text{with eigenvalue } \lambda = \frac{1}{2}, \\ a_0 = 2, & \text{with eigenvalue } \lambda = \frac{n-4}{2(n-2)}, \end{cases}$
- $P_{2,(2,0,\dots,0),0} \Rightarrow a_0 = 2$ with eigenvalue $\lambda = 1$.

Now we describe the eigenvalues of these eigenspaces with respect to the operator M and to join the results.

If F is a fundamental function of type $(a_0, a_1, \dots, a_{m-1})$, then it has eigenvalue $\frac{1}{2} \sum_{j=0}^{m-1} a_j \lambda_j$, where $\lambda_j = 1 - \frac{q-1}{q^{m-j-1}}$ is the eigenvalue of the eigenspace W_j , of dimension $q^{j-1}(q - 1)$, occurring in the spectral decomposition of $L(Y)$. From this we can fill the following table in which we give the eigenspaces, together with the corresponding eigenvalue and dimension.

- $P_{2,(a_0,a_1,\dots,a_{m-1}),2}$. We have three different cases:

(1) if $a_0 = 0$, the corresponding eigenspace is

$$P_{2,(0,\dots,0, \underbrace{1}_{i\text{th place}}, 0,\dots,0, \underbrace{1}_{j\text{th place}}, 0,\dots,0),2}$$

of dimension $n(n-1)(q-1)^2q^{i-1}q^{j-1}$, with eigenvalue $\frac{p_0}{2}(\lambda_i + \lambda_j)$;

(2) if $a_0 = 1$, the corresponding eigenspace is

$$P_{2,(1,\dots,0, \underbrace{1}_{i\text{th place}}, 0,\dots,0),2}$$

of dimension $n(n-2)(q-1)q^{i-1}$, with eigenvalue $p_0\frac{1+\lambda_i}{2} + (1-p_0)\frac{-1}{2(n-2)}$;

(3) if $a_0 = 2$, the corresponding eigenspace is $P_{2,(2,0,\dots,0),2}$ of dimension $\frac{n(n-3)}{2}$ with eigenvalue $p_0 + (1-p_0)\frac{-1}{n-2}$.

- $P_{2,(a_0,a_1,\dots,a_{m-1}),1}$. We have two different cases:

(1) if $a_0 = 1$, the corresponding eigenspace is

$$P_{2,(1,\dots,0, \underbrace{1}_{i\text{th place}}, 0,\dots,0),1}$$

of dimension $n(q-1)q^{i-1}$, with eigenvalue $p_0\frac{1+\lambda_i}{2} + \frac{1-p_0}{2}$;

(2) if $a_0 = 2$, the corresponding eigenspace is $P_{2,(2,0,\dots,0),1}$ of dimension $n-1$, with eigenvalue $p_0 + (1-p_0)\frac{n-4}{2(n-2)}$.

- $P_{2,(2,0,\dots,0),0}$. In this case, the dimension of the eigenspace is 1 with eigenvalue 1.

REFERENCES

- [1] ALDOUS, D. and FILL, J. (2009). Reversible Markov chains and random walk on graphs. Monograph in preparation. Available at <http://www.stat.berkeley.edu/~aldous/RWG/book.html>.
- [2] BAILEY, R. A. (2004). *Association Schemes: Designed Experiments, Algebra and Combinatorics*. Cambridge Studies in Advanced Mathematics **84**. Cambridge Univ. Press, Cambridge. [MRMR2047311](#)
- [3] BAILEY, R. A. and CAMERON, P. J. (2005). Crested products of association schemes. *J. Lond. Math. Soc. (2)* **72** 1–24. [MRMR2145725](#)
- [4] CECCHERINI-SILBERSTEIN, T., SCARABOTTI, F. and TOLLI, F. (2006). Trees, wreath products and finite Gelfand pairs. *Adv. Math.* **206** 503–537. [MRMR2263713](#)
- [5] CECCHERINI-SILBERSTEIN, T., SCARABOTTI, F. and TOLLI, F. (2007). Finite Gelfand pairs and their applications to probability and statistics. *J. Math. Sci. (N. Y.)* **141** 1182–1229.
- [6] CECCHERINI-SILBERSTEIN, T., SCARABOTTI, F. and TOLLI, F. (2008). *Harmonic Analysis on Finite Groups: Representation Theory, Gelfand Pairs and Markov Chains*. Cambridge Studies in Advanced Mathematics **108**. Cambridge Univ. Press, Cambridge. [MRMR2389056](#)

- [7] DIACONIS, P. (1988). *Group Representations in Probability and Statistics. Lecture Notes—Monograph Series* **11**. IMS, Hayward, CA. [MRMR964069](#)
- [8] DIACONIS, P. and SALOFF-COSTE, L. (1993). Comparison theorems for reversible Markov chains. *Ann. Appl. Probab.* **3** 696–730. [MRMR1233621](#)
- [9] DIACONIS, P. and SALOFF-COSTE, L. (1993). Comparison techniques for random walk on finite groups. *Ann. Probab.* **21** 2131–2156. [MRMR1245303](#)
- [10] DIACONIS, P. and SHAHSHAHANI, M. (1987). Time to reach stationarity in the Bernoulli–Laplace diffusion model. *SIAM J. Math. Anal.* **18** 208–218. [MRMR871832](#)
- [11] FIGÀ-TALAMANCA, A. (2001). An application of Gelfand pairs to a problem of diffusion in compact ultrametric spaces. In *Topics in Probability and Lie Groups: Boundary Theory. CRM Proceedings & Lecture Notes* **28** 51–67. Amer. Math. Soc., Providence, RI. [MRMR1832434](#)
- [12] LIGGETT, T. M. (1985). *Interacting Particle Systems. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]* **276**. Springer, New York. [MRMR776231](#)

SECTION DE MATHÉMATIQUES
UNIVERSITÉ DE GENÈVE
2-4, RUE DU LIÈVRE, CASE POSTALE 64
1211 GENÈVE 4
SUISSE
E-MAIL: Daniele.Dangeli@math.unige.ch
Alfredo.Donno@math.unige.ch