

# LECTURE 10

## LECTURE OUTLINE


- Min Common/Max Crossing Th. III
- Nonlinear Farkas Lemma/Linear Constraints
- Linear Programming Duality
- Convex Programming Duality
- Optimality Conditions

**Reading:** Sections 4.5, 5.1, 5.2, 5.3.1, 5.3.2

**Recall the MC/MC Theorem II:** If  $-\infty < w^*$   
and

$0 \in \text{ri}(D) = \{u \mid \text{there exists } w \in \mathfrak{R} \text{ with } (u, w) \in \overline{M}\}$

then  $q^* = w^*$  and there exists  $\mu$  s. t.  $q(\mu) = q^*$ .




## MC/MC TH. III - POLYHEDRAL

- Consider the MC/MC problems, and assume that  $-\infty < w^*$  and:

- (1)  $\bar{M}$  is a “horizontal translation” of  $\tilde{M}$  by  $-P$ ,

$$\bar{M} = \tilde{M} - \{(u, 0) \mid u \in P\},$$

where  $P$ : polyhedral and  $\tilde{M}$ : convex.



- (2) We have  $\text{ri}(\tilde{D}) \cap P \neq \emptyset$ , where

$$\tilde{D} = \{u \mid \text{there exists } w \in \Re \text{ with } (u, w) \in \tilde{M}\}$$


Then  $q^* = w^*$ , there is a max crossing solution, and all max crossing solutions  $\bar{\mu}$  satisfy  $\bar{\mu}'d \leq 0$  for all  $d \in R_P$ .

- **Comparison with Th. II:** Since  $D = \tilde{D} - P$ , the condition  $0 \in \text{ri}(D)$  of Theorem II is

$$\text{ri}(\tilde{D}) \cap \text{ri}(P) \neq \emptyset$$

## PROOF OF MC/MC TH. III

- Consider the *disjoint* convex sets  $C_1 = \{(u, v) \mid v > w \text{ for some } (u, w) \in \tilde{M}\}$  and  $C_2 = \{(u, w^*) \mid u \in P\}$  [ $u \in P$  and  $(u, w) \in \tilde{M}$  with  $w^* > w$  contradicts the definition of  $w^*$ ]



- Since  $C_2$  is polyhedral, there exists a separating hyperplane not containing  $C_1$ , i.e., a  $(\bar{\mu}, \beta) \neq (0, 0)$  such that

$$\beta w^* + \bar{\mu}' z \leq \beta v + \bar{\mu}' x, \quad \forall (x, v) \in C_1, \quad \forall z \in P$$

$$\inf_{(x,v) \in C_1} \{\beta v + \bar{\mu}' x\} < \sup_{(x,v) \in C_1} \{\beta v + \bar{\mu}' x\}$$

Since  $(0, 1)$  is a direction of recession of  $C_1$ , we see that  $\beta \geq 0$ . Because of the relative interior point assumption,  $\beta \neq 0$ , so we may assume that  $\beta = 1$ .

## PROOF (CONTINUED)

- Hence,

$$w^* + \bar{\mu}'z \leq \inf_{(u,v) \in C_1} \{v + \bar{\mu}'u\}, \quad \forall z \in P,$$

so that

$$\begin{aligned} w^* &\leq \inf_{(u,v) \in C_1, z \in P} \{v + \bar{\mu}'(u - z)\} \\ &= \inf_{(u,v) \in \tilde{M} - P} \{v + \bar{\mu}'u\} \\ &= \inf_{(u,v) \in \overline{M}} \{v + \bar{\mu}'u\} \\ &= q(\bar{\mu}) \end{aligned}$$

Using  $q^* \leq w^*$  (weak duality), we have  $q(\bar{\mu}) = q^* = w^*$ .

Proof that all max crossing solutions  $\bar{\mu}$  satisfy  $\bar{\mu}'d \leq 0$  for all  $d \in R_P$ : follows from

$$q(\mu) = \inf_{(u,v) \in C_1, z \in P} \{v + \mu'(u - z)\}$$

so that  $q(\mu) = -\infty$  if  $\mu'd > 0$ . **Q.E.D.**

- Geometrical intuition: every  $(0, -d)$  with  $d \in R_P$ , is direction of recession of  $\overline{M}$ .

## MC/MC TH. III - A SPECIAL CASE


- Consider the MC/MC framework, and assume:

- For a convex function  $f : \mathfrak{R}^m \mapsto (-\infty, \infty]$ , an  $r \times m$  matrix  $A$ , and a vector  $b \in \mathfrak{R}^r$ :

$$\overline{M} = \{ (u, w) \mid \text{for some } (x, w) \in \text{epi}(f), Ax - b \leq u \}$$

so  $\overline{M} = \tilde{M} + \text{Positive Orthant}$ , where

$$\tilde{M} = \{ (Ax - b, w) \mid (x, w) \in \text{epi}(f) \}$$



- There is an  $\bar{x} \in \text{ri}(\text{dom}(f))$  s. t.  $A\bar{x} - b \leq 0$ .

Then  $q^* = w^*$  and there is a  $\mu \geq 0$  with  $q(\mu) = q^*$ .

- Also  $\overline{M} = M \approx \text{epi}(p)$ , where  $p(u) = \inf_{Ax - b \leq u} f(x)$ .
- We have  $w^* = p(0) = \inf_{Ax - b \leq 0} f(x)$ .

# NONL. FARKAS' L. - POLYHEDRAL ASSUM.

- Let  $X \subset \mathfrak{R}^n$  be convex, and  $f : X \mapsto \mathfrak{R}$  and  $g_j : \mathfrak{R}^n \mapsto \mathfrak{R}$ ,  $j = 1, \dots, r$ , be linear so  $g(x) = Ax - b$  for some  $A$  and  $b$ . Assume that


$$f(x) \geq 0, \quad \forall x \in X \text{ with } Ax - b \leq 0$$

Let

$$Q^* = \{ \mu \mid \mu \geq 0, f(x) + \mu'(Ax - b) \geq 0, \forall x \in X \}.$$

Assume that there exists a vector  $\bar{x} \in \text{ri}(X)$  such that  $A\bar{x} - b \leq 0$ . Then  $Q^*$  is nonempty.

**Proof:** As before, apply special case of MC/MC Th. III of preceding slide, using the fact  $w^* \geq 0$ , implied by the assumption.



## (LINEAR) FARKAS' LEMMA

- Let  $A$  be an  $m \times n$  matrix and  $c \in \mathfrak{R}^m$ . The system  $Ay = c, y \geq 0$  has a solution if and only if

$$A'x \leq 0 \quad \Rightarrow \quad c'x \leq 0. \quad (*)$$

- **Alternative/Equivalent Statement:** If  $P = \text{cone}\{a_1, \dots, a_n\}$ , where  $a_1, \dots, a_n$  are the columns of  $A$ , then  $P = (P^*)^*$  (Polar Cone Theorem).

**Proof:** If  $y \in \mathfrak{R}^n$  is such that  $Ay = c, y \geq 0$ , then  $y'A'x = c'x$  for all  $x \in \mathfrak{R}^m$ , which implies Eq. (\*).

Conversely, apply the Nonlinear Farkas' Lemma with  $f(x) = -c'x$ ,  $g(x) = A'x$ , and  $X = \mathfrak{R}^m$ . Condition (\*) implies the existence of  $\mu \geq 0$  such that

$$-c'x + \mu'A'x \geq 0, \quad \forall x \in \mathfrak{R}^m,$$

or equivalently

$$(A\mu - c)'x \geq 0, \quad \forall x \in \mathfrak{R}^m,$$

or  $A\mu = c$ .

# LINEAR PROGRAMMING DUALITY

- Consider the linear program

$$\begin{aligned} & \text{minimize } c'x \\ & \text{subject to } a'_j x \geq b_j, \quad j = 1, \dots, r, \end{aligned}$$

where  $c \in \mathfrak{R}^n$ ,  $a_j \in \mathfrak{R}^n$ , and  $b_j \in \mathfrak{R}$ ,  $j = 1, \dots, r$ .

- The dual problem is

$$\begin{aligned} & \text{maximize } b'\mu \\ & \text{subject to } \sum_{j=1}^r a_j \mu_j = c, \quad \mu \geq 0. \end{aligned}$$


- **Linear Programming Duality Theorem:**

- (a) If either  $f^*$  or  $q^*$  is finite, then  $f^* = q^*$  and both the primal and the dual problem have optimal solutions.
- (b) If  $f^* = -\infty$ , then  $q^* = -\infty$ .
- (c) If  $q^* = \infty$ , then  $f^* = \infty$ .

**Proof:** (b) and (c) follow from weak duality. For part (a): If  $f^*$  is finite, there is a primal optimal solution  $x^*$ , by existence of solutions of quadratic programs. Use Farkas' Lemma to construct a dual feasible  $\mu^*$  such that  $c'x^* = b'\mu^*$  (next slide).



## PROOF OF LP DUALITY (CONTINUED)



- Let  $x^*$  be a primal optimal solution, and let  $J = \{j \mid a'_j x^* = b_j\}$ . Then,  $c'y \geq 0$  for all  $y$  in the cone of “feasible directions”

$$D = \{y \mid a'_j y \geq 0, \forall j \in J\}$$

By Farkas' Lemma, for some scalars  $\mu_j^* \geq 0$ ,  $c$  can be expressed as

$$c = \sum_{j=1}^r \mu_j^* a_j, \quad \mu_j^* \geq 0, \forall j \in J, \quad \mu_j^* = 0, \forall j \notin J.$$

Taking inner product with  $x^*$ , we obtain  $c'x^* = b'\mu^*$ , which in view of  $q^* \leq f^*$ , shows that  $q^* = f^*$  and that  $\mu^*$  is optimal.

# LINEAR PROGRAMMING OPT. CONDITIONS

A pair of vectors  $(x^*, \mu^*)$  form a primal and dual optimal solution pair if and only if  $x^*$  is primal-feasible,  $\mu^*$  is dual-feasible, and

$$\mu_j^*(b_j - a_j'x^*) = 0, \quad \forall j = 1, \dots, r. \quad (*)$$

**Proof:** If  $x^*$  is primal-feasible and  $\mu^*$  is dual-feasible, then

$$\begin{aligned} b'\mu^* &= \sum_{j=1}^r b_j\mu_j^* + \left( c - \sum_{j=1}^r a_j\mu_j^* \right)' x^* \\ &= c'x^* + \sum_{j=1}^r \mu_j^*(b_j - a_j'x^*) \end{aligned} \quad (**)$$

So if Eq. (\*) holds, we have  $b'\mu^* = c'x^*$ , and weak duality implies that  $x^*$  is primal optimal and  $\mu^*$  is dual optimal.

Conversely, if  $(x^*, \mu^*)$  form a primal and dual optimal solution pair, then  $x^*$  is primal-feasible,  $\mu^*$  is dual-feasible, and by the duality theorem, we have  $b'\mu^* = c'x^*$ . From Eq. (\*\*), we obtain Eq. (\*).

# CONVEX PROGRAMMING

Consider the problem

$$\begin{aligned} & \text{minimize } f(x) \\ & \text{subject to } x \in X, g_j(x) \leq 0, j = 1, \dots, r, \end{aligned}$$

where  $X \subset \mathfrak{R}^n$  is convex, and  $f : X \mapsto \mathfrak{R}$  and  $g_j : X \mapsto \mathfrak{R}$  are convex. Assume  $f^*$ : finite.

- Recall the connection with the max crossing problem in the MC/MC framework where  $M = \text{epi}(p)$  with

$$p(u) = \inf_{x \in X, g(x) \leq u} f(x)$$

- Consider the Lagrangian function

$$L(x, \mu) = f(x) + \mu'g(x),$$

the dual function

$$q(\mu) = \begin{cases} \inf_{x \in X} L(x, \mu) & \text{if } \mu \geq 0, \\ -\infty & \text{otherwise} \end{cases}$$

and the dual problem of maximizing  $\inf_{x \in X} L(x, \mu)$  over  $\mu \geq 0$ .

# STRONG DUALITY THEOREM

• Assume that  $f^*$  is finite, and that one of the following two conditions holds:

(1) There exists  $\bar{x} \in X$  such that  $g(\bar{x}) < 0$ .

(2) The functions  $g_j, j = 1, \dots, r$ , are affine, and there exists  $\bar{x} \in \text{ri}(X)$  such that  $g(\bar{x}) \leq 0$ .

Then  $q^* = f^*$  and the set of optimal solutions of the dual problem is nonempty. Under condition (1) this set is also compact.

• **Proof:** Replace  $f(x)$  by  $f(x) - f^*$  so that  $f(x) - f^* \geq 0$  for all  $x \in X$  w/  $g(x) \leq 0$ . Apply Nonlinear Farkas' Lemma. Then, there exist  $\mu_j^* \geq 0$ , s.t.

$$f^* \leq f(x) + \sum_{j=1}^r \mu_j^* g_j(x), \quad \forall x \in X$$

• It follows that

$$f^* \leq \inf_{x \in X} \{f(x) + \mu^{*'} g(x)\} \leq \inf_{x \in X, g(x) \leq 0} f(x) = f^*.$$

Thus equality holds throughout, and we have

$$f^* = \inf_{x \in X} \left\{ f(x) + \sum_{j=1}^r \mu_j^* g_j(x) \right\} = q(\mu^*)$$

# QUADRATIC PROGRAMMING DUALITY

- Consider the quadratic program

$$\begin{aligned} & \text{minimize} \quad \frac{1}{2}x'Qx + c'x \\ & \text{subject to} \quad Ax \leq b, \end{aligned}$$

where  $Q$  is positive definite.

- If  $f^*$  is finite, then  $f^* = q^*$  and there exist both primal and dual optimal solutions, since the constraints are linear.

- Calculation of dual function:

$$q(\mu) = \inf_{x \in \mathbb{R}^n} \left\{ \frac{1}{2}x'Qx + c'x + \mu'(Ax - b) \right\}$$

The infimum is attained for  $x = -Q^{-1}(c + A'\mu)$ , and, after substitution and calculation,

$$q(\mu) = -\frac{1}{2}\mu'AQ^{-1}A'\mu - \mu'(b + AQ^{-1}c) - \frac{1}{2}c'Q^{-1}c$$

- The dual problem, after a sign change, is

$$\begin{aligned} & \text{minimize} \quad \frac{1}{2}\mu'P\mu + t'\mu \\ & \text{subject to} \quad \mu \geq 0, \end{aligned}$$

where  $P = AQ^{-1}A'$  and  $t = b + AQ^{-1}c$ .

## OPTIMALITY CONDITIONS

- We have  $q^* = f^*$ , and the vectors  $x^*$  and  $\mu^*$  are optimal solutions of the primal and dual problems, respectively, iff  $x^*$  is feasible,  $\mu^* \geq 0$ , and

$$x^* \in \arg \min_{x \in X} L(x, \mu^*), \quad \mu_j^* g_j(x^*) = 0, \quad \forall j. \quad (1)$$

**Proof:** If  $q^* = f^*$ , and  $x^*, \mu^*$  are optimal, then

$$\begin{aligned} f^* = q^* = q(\mu^*) &= \inf_{x \in X} L(x, \mu^*) \leq L(x^*, \mu^*) \\ &= f(x^*) + \sum_{j=1}^r \mu_j^* g_j(x^*) \leq f(x^*), \end{aligned}$$

where the last inequality follows from  $\mu_j^* \geq 0$  and  $g_j(x^*) \leq 0$  for all  $j$ . Hence equality holds throughout above, and (1) holds.

Conversely, if  $x^*, \mu^*$  are feasible, and (1) holds,

$$\begin{aligned} q(\mu^*) &= \inf_{x \in X} L(x, \mu^*) = L(x^*, \mu^*) \\ &= f(x^*) + \sum_{j=1}^r \mu_j^* g_j(x^*) = f(x^*), \end{aligned}$$

so  $q^* = f^*$ , and  $x^*, \mu^*$  are optimal. **Q.E.D.**

# QUADRATIC PROGRAMMING OPT. COND.

For the quadratic program

$$\begin{aligned} & \text{minimize } \frac{1}{2}x'Qx + c'x \\ & \text{subject to } Ax \leq b, \end{aligned}$$

where  $Q$  is positive definite,  $(x^*, \mu^*)$  is a primal and dual optimal solution pair if and only if:

- Primal and dual feasibility holds:

$$Ax^* \leq b, \quad \mu^* \geq 0$$

- Lagrangian optimality holds [ $x^*$  minimizes  $L(x, \mu^*)$  over  $x \in \mathbb{R}^n$ ]. This yields

$$x^* = -Q^{-1}(c + A'\mu^*)$$

- Complementary slackness holds [ $(Ax^* - b)'\mu^* = 0$ ]. It can be written as

$$\mu_j^* > 0 \quad \Rightarrow \quad a'_j x^* = b_j, \quad \forall j = 1, \dots, r,$$

where  $a'_j$  is the  $j$ th row of  $A$ , and  $b_j$  is the  $j$ th component of  $b$ .

# LINEAR EQUALITY CONSTRAINTS

- The problem is

$$\begin{aligned} & \text{minimize} && f(x) \\ & \text{subject to} && x \in X, \quad g(x) \leq 0, \quad Ax = b, \end{aligned}$$

where  $X$  is convex,  $g(x) = (g_1(x), \dots, g_r(x))'$ ,  $f : X \mapsto \mathfrak{R}$  and  $g_j : X \mapsto \mathfrak{R}$ ,  $j = 1, \dots, r$ , are convex.

- Convert the constraint  $Ax = b$  to  $Ax \leq b$  and  $-Ax \leq -b$ , with corresponding dual variables  $\lambda^+ \geq 0$  and  $\lambda^- \geq 0$ .
- The Lagrangian function is

$$f(x) + \mu'g(x) + (\lambda^+ - \lambda^-)'(Ax - b),$$

and by introducing a dual variable  $\lambda = \lambda^+ - \lambda^-$ , with no sign restriction, it can be written as

$$L(x, \mu, \lambda) = f(x) + \mu'g(x) + \lambda'(Ax - b).$$

- The dual problem is

$$\begin{aligned} & \text{maximize} && q(\mu, \lambda) \equiv \inf_{x \in X} L(x, \mu, \lambda) \\ & \text{subject to} && \mu \geq 0, \quad \lambda \in \mathfrak{R}^m. \end{aligned}$$



# DUALITY AND OPTIMALITY COND.

- **Pure equality constraints:**

- (a) Assume that  $f^*$ : finite and there exists  $\bar{x} \in \text{ri}(X)$  such that  $A\bar{x} = b$ . Then  $f^* = q^*$  and there exists a dual optimal solution.
- (b)  $f^* = q^*$ , and  $(x^*, \lambda^*)$  are a primal and dual optimal solution pair if and only if  $x^*$  is feasible, and

$$x^* \in \arg \min_{x \in X} L(x, \lambda^*)$$

**Note:** No complementary slackness for equality constraints.

- **Linear and nonlinear constraints:**

- (a) Assume  $f^*$ : finite, that there exists  $\bar{x} \in X$  such that  $A\bar{x} = b$  and  $g(\bar{x}) < 0$ , and that there exists  $\tilde{x} \in \text{ri}(X)$  such that  $A\tilde{x} = b$ . Then  $q^* = f^*$  and there exists a dual optimal solution.
- (b)  $f^* = q^*$ , and  $(x^*, \mu^*, \lambda^*)$  are a primal and dual optimal solution pair if and only if  $x^*$  is feasible,  $\mu^* \geq 0$ , and

$$x^* \in \arg \min_{x \in X} L(x, \mu^*, \lambda^*), \quad \mu_j^* g_j(x^*) = 0, \quad \forall j$$