

Faster Algorithms for Rectangular Matrix Multiplication

François Le Gall
Department of Computer Science
Graduate School of Information Science and Technology
The University of Tokyo
legall@is.s.u-tokyo.ac.jp

Abstract

Let α be the maximal value such that the product of an $n \times n^\alpha$ matrix by an $n^\alpha \times n$ matrix can be computed with $n^{2+o(1)}$ arithmetic operations. In this paper we show that $\alpha > 0.30298$, which improves the previous record $\alpha > 0.29462$ by Coppersmith (Journal of Complexity, 1997). More generally, we construct a new algorithm for multiplying an $n \times n^k$ matrix by an $n^k \times n$ matrix, for any value $k \neq 1$. The complexity of this algorithm is better than all known algorithms for rectangular matrix multiplication. In the case of square matrix multiplication (i.e., for $k = 1$), we recover exactly the complexity of the algorithm by Coppersmith and Winograd (Journal of Symbolic Computation, 1990).

These new upper bounds can be used to improve the time complexity of several known algorithms that rely on rectangular matrix multiplication. For example, we directly obtain a $O(n^{2.5302})$ -time algorithm for the all-pairs shortest paths problem over directed graphs with small integer weights, improving over the $O(n^{2.575})$ -time algorithm by Zwick (JACM 2002), and also improve the time complexity of sparse square matrix multiplication.

1 Introduction

Background. Matrix multiplication is one of the most fundamental problems in computer science and mathematics. Besides the fact that several computational problems in linear algebra can be reduced to the computation of the product of two matrices, the complexity of matrix multiplication also arises as a bottleneck in a multitude of other computational tasks (e.g., graph algorithms). The standard method for multiplying two $n \times n$ matrices uses $O(n^3)$ arithmetic operations. Strassen showed in 1969 that this trivial algorithm is not optimal, and gave a algorithm that uses only $O(n^{2.808})$ arithmetic operations. This has been the beginning of a long story of improvements that lead to the upper bound $O(n^{2.376})$ by Coppersmith and Winograd [10], which has been further improved to $O(n^{2.3727})$ very recently by Vassilevska Williams [27]. Note that all the above complexities refer to the number of arithmetic operations involved, but naturally the same upper bounds hold for the time complexity as well when each arithmetic operation can be done in negligible time (e.g., in $\text{poly}(\log n)$ time).

Finding the optimal value of the exponent of square matrix multiplication is naturally one of the most important open problems in algebraic complexity. It is widely believed that the product of two $n \times n$ matrices can be computed with $O(n^{2+\epsilon})$ arithmetic operations for any constant $\epsilon > 0$. Several conjectures, including conjectures about combinatorial structures [10] and about group theory [7, 6], would, if true, lead to this result (see also [1] for recent work on these conjectures). Another way to interpret this open problem is by considering the multiplication of an $n \times m$ matrix by an $m \times n$ matrix. Suppose that the matrices are defined over a field. For any $k > 0$, define the exponent of such a rectangular matrix multiplication as follows:

$$\omega(1, 1, k) = \inf\{\tau \in \mathbb{R} \mid C(n, n, \lfloor n^k \rfloor) = O(n^\tau)\},$$

where $C(n, n, \lfloor n^k \rfloor)$ denotes the minimum number of arithmetic operations needed to multiply an $n \times \lfloor n^k \rfloor$ matrix by an $\lfloor n^k \rfloor \times n$ matrix. Note that, while the value $\omega(1, 1, k)$ may depend on the field

under consideration, it is known that it can depend only on the characteristic of the field [23]. Define $\omega = \omega(1, 1, 1)$ and $\alpha = \sup\{k \mid \omega(1, 1, k) = 2\}$. The value ω represents the exponent of square matrix multiplication, and the value α essentially represents the largest value such that the product of an $n \times n^\alpha$ matrix by an $n^\alpha \times n$ matrix can be computed with $O(n^{2+\epsilon})$ arithmetic operations for any constant ϵ . Since $\omega = 2$ if and only if $\alpha = 1$, one possible strategy towards showing that $\omega = 2$ is to give lower bounds on α . Coppersmith [8] showed in 1982 that $\alpha > 0.172$. Then, based on the techniques developed in [10], Coppersmith [9] improved this lower bound to $\alpha > 0.29462$. This is the best lower bound on α known so far.

Excepting Coppersmith's works on the value α , there have been relatively few algorithms that focused specifically on rectangular matrix multiplication. Since it is well known (see, e.g., [17]) that multiplying an $n \times n$ matrix by an $n \times m$ matrix, or an $m \times n$ matrix by an $n \times n$ matrix, can be done with the same number of arithmetic operations as multiplying an $n \times m$ matrix by an $m \times n$ matrix, the value $\omega(1, 1, k)$ represents the exponent of all these three types of rectangular matrix multiplications. Note that, by decomposing the product into smaller matrix products, it is easy to obtain (see, e.g., [17]) the following upper bound:

$$\omega(1, 1, k) = \begin{cases} 2 & \text{if } 0 \leq k \leq \alpha \\ 2 + (\omega - 2) \frac{k - \alpha}{1 - \alpha} & \text{if } \alpha \leq k \leq 1. \end{cases} \quad (1)$$

Lotti and Romani [17] obtained nontrivial upper bounds on $\omega(1, 1, k)$ based on the seminal result by Coppersmith [8] and on early works on square matrix multiplication. Huang and Pan [13] showed how to apply ideas from [10] to the rectangular setting and obtained the upper bound $\omega(1, 1, 2) < 3.333954$, but this approach did not lead to any upper bound better than (1) for $k \leq 1$. Ke, Zeng, Han and Pan [16] further improved Huang and Pan's result to $\omega(1, 1, 2) < 3.2699$, by using again the approach from [10], and also reported the upper bounds $\omega(1, 1, 0.8) < 2.2356$ and $\omega(1, 1, 0.5356) < 2.0712$, which are better than those obtained by (1). Their approach, nevertheless, did not give any improvement for the value of α .

Besides the fact that a better understanding of $\omega(1, 1, k)$ gives insights into the nature of matrix multiplication and ultimately may help showing that $\omega = 2$, fast algorithms for multiplying an $n \times n^k$ matrix by an $n^k \times n$ with $k \neq 1$ have also a multitude of applications. Typical examples not directly related to linear algebra include the construction of fast algorithms for the all-pairs shortest paths problem [2, 19, 29, 33, 34], the dynamic computation of the transitive closure [12, 22], finding ancestors [11], detecting directed cycles [31], or computing the diameter of a graph [30]. Rectangular matrix multiplication has also been used in computational complexity [18, 28], and to speed-up sparse square matrix multiplication [3, 15, 32] or tasks in computational geometry [14, 15]. Obtaining new upper bounds on $\omega(1, 1, k)$ would thus reduce the asymptotic time complexity of algorithms in a wide range of areas. We nevertheless stress that such improvements are only of theoretical interest, since the huge constants involved in the complexity of fast matrix multiplication usually make these algorithms impractical.

Short description of the approach by Coppersmith and Winograd. The results [9, 13, 16, 24, 27] mentioned above are all obtained by extending the approach by Coppersmith and Winograd [10]. This approach is an illustration of a general methodology initiated in the 1970's based on the theory of bilinear and trilinear forms, through which most of the improvements for matrix multiplication have been obtained. Informally, the idea is to start with a *basic construction* (some small trilinear form), and then exploit general properties of matrix multiplication (in particular Schönhage's asymptotic sum inequality [23]) to derive an upper bound on the exponent ω from this construction. The main contributions of [10] consist of two parts: the discovery of new basic constructions and the introduction of strong techniques to analyze them. In their paper, Coppersmith and Winograd actually present three algorithms, based on three different basic constructions. The first basic construction (Section 6 in [10]) is the simplest of the three and leads to the upper bound $\omega < 2.40364$. The second basic construction (Section 7 in [10]), that we will refer in this paper as F_q (here $q \in \mathbb{N}$ is a parameter), leads to the upper bound

$\omega < 2.38719$. The third basic construction (Section 8 in [10]) is $F_q \otimes F_q$, the tensor product of two instances of F_q , and leads to the improved upper bound $\omega < 2.375477$.

In view of the last result, it was natural to ask if taking larger tensor powers of F_q as the basic construction leads to better bounds on ω . The case $r = 3$ was explicitly mentioned as an open problem in [10] but did not seem to lead to any improvement. Stothers [24] and Vassilevska Williams [27] succeeded in analyzing the fourth tensor product $F_q^{\otimes 4}$ and obtained a better upper bound on ω , the first improvement in more than twenty years. Vassilevska Williams further presented a general framework that enables a systematic analysis for higher tensor products of the basic construction, and used this framework to show that $\omega < 2.3727$, for the basic construction $F^{\otimes 8}$, the best upper bound obtained so far.

The algorithms for rectangular matrix multiplication [9, 13, 16] already mentioned use a similar approach. Huang and Pan [13] obtained their improvement on $\omega(1, 1, 2)$ by taking the easiest of the three constructions in [10] and carefully modifying the analysis to evaluate the complexity of rectangular matrix multiplication. Ke, Zeng, Han and Pan [16] obtained their improvements similarly, but by using the second basic construction from [10] (the construction F_q) instead, which led to better upper bounds. These approaches, while very natural, do not provide any nontrivial lower bounds on α : the upper bounds on $\omega(1, 1, k)$ obtained are strictly larger than 2 even for small values of k . In order to obtain the lower bound $\alpha > 0.29462$, Coppersmith [9] relied on a more complex approach: the basic construction considered is still F_q , but several instances for distinct values of q are combined together in a subtle way in order to keep the complexity of the resulting algorithm small enough (i.e., not larger than $n^{2+o(1)}$).

Statement of our results and discussion. In this paper we construct new algorithms for rectangular matrix multiplication, by taking the tensor power $F_q \otimes F_q$ as basic construction and analyzing this construction in the framework of rectangular matrix multiplication. We use these ideas to prove that $\omega(1, 1, k) = 2$ for any $k \leq 0.3029805$, as stated in the following theorem.

Theorem 1.1. *For any value $k < 0.3029805\dots$, the product of an $n \times n^k$ matrix by an $n^k \times n$ matrix can be computed with $O(n^{2+\epsilon})$ arithmetic operations for any constant $\epsilon > 0$.*

Theorem 1.1 shows that $\alpha > 0.30298$, which improves the previous record $\alpha > 0.29462$ by Coppersmith. More generally, in the present work we present an algorithm for multiplying an $n \times n^k$ matrix by an $n^k \times n$ matrix, for any value k . We show that the complexity of this algorithm can be expressed as a (nonlinear) optimization problem, and use this formulation to derive upper bounds on $\omega(1, 1, k)$. Table 1 shows the bounds we obtain for several values of k . The bounds obtained for $0 \leq k \leq 1$ are represented in Figure 1 as well.

The results of this paper can be seen as a generalization of Coppersmith-Winograd's approach to the rectangular setting. In the case of square matrix multiplication (i.e., for $k = 1$), we recover naturally the same upper bound $\omega(1, 1, 1) < 2.375477$ as the one obtained in [10]. Let us mention that we can, in a rather straightforward way, combine our results with the upper bound $\omega < 2.3727$ by Vassilevska Williams [27] to obtain slightly improved bounds for $k \approx 1$. The idea is, very similarly to how Equation (1) was obtained, to exploit the convexity of the function $\omega(1, 1, k)$. Concretely, for any fixed value $0 \leq k_0 < 1$, the inequality

$$\omega(1, 1, k) \leq \omega(1, 1, k_0) + (\omega - \omega(1, 1, k_0)) \frac{k - k_0}{1 - k_0}$$

holds for any k such that $k_0 \leq k \leq 1$. This enables us to combine an upper bound on $\omega(1, 1, k_0)$, for instance one of the values in Table 1, with the improved upper bound $\omega < 2.3727$ by Vassilevska Williams. Since the improvement is small and concerns only the case $k \approx 1$, we will not discuss it further.

For $k > 0.29462$ and $k \neq 1$, the complexity of our algorithms is better than all known algorithms for rectangular matrix multiplication, including the algorithms [13, 16] mentioned above. Moreover, for

k	upper bound on $\omega(1, 1, k)$	k	upper bound on $\omega(1, 1, k)$	k	upper bound on $\omega(1, 1, k)$
0.30298	2	0.60	2.096571	1.30	2.624703
0.31	2.000063	0.65	2.125676	1.40	2.711707
0.32	2.000371	0.70	2.156959	1.50	2.800116
0.33	2.000939	0.75	2.190087	1.75	3.025906
0.34	2.001771	0.80	2.224790	2.00	3.256689
0.35	2.002870	0.85	2.260830	2.25	3.490957
0.40	2.012175	0.90	2.298048	2.50	3.727808
0.45	2.027102	0.95	2.336306	3.00	4.207372
0.50	2.046681	1.00	2.375477	3.50	4.693151
0.5302	2.060396	1.10	2.456151	4.00	5.180715
0.55	2.070063	1.20	2.539392	5.00	6.166736

Table 1: Our upper bounds on the exponent of the multiplication of an $n \times n^k$ matrix by an $n^k \times n$ matrix.




Figure 1: Our upper bounds (in plain line) on $\omega(1, 1, k)$, for $0 \leq k \leq 1$. The dashed line represents the upper bounds on $\omega(1, 1, k)$ obtained by using Equation (1) with the values $\alpha > 0.30298$ and $\omega < 2.375477$.

$0.30298 < k < 1$, our new bounds are significantly better than what can be obtained solely from the bound $\alpha > 0.30298$ and $\omega < 2.375477$ through Equation (1), as illustrated in Figure 1. This suggests that non-negligible improvements can be obtained for all applications of rectangular matrix multiplications that rely on this simple linear interpolation — we will elaborate on this subject in Subsection 1.1.

Let us compare more precisely our results with those reported in [16]. For $k = 2$, we obtain $\omega(1, 1, 2) < 3.256689$ while Ke et al. [16] obtained $\omega(1, 1, 2) < 3.2699$ by using the basic construction F_q . Our improvements are of the same order for the other two values ($k = 0.8$ and $k = 0.5356$) analyzed in [16], as can be seen from Table 1. Note that the order of magnitude of the improvements here is similar to what was obtained in [10] by changing the basic construction from F_q to $F_q \otimes F_q$ for square matrix multiplication, which led to a improvement from $\omega < 2.38719$ to $\omega < 2.375477$.

A noteworthy point is that our algorithm directly leads to improved lower bounds on α while, as already mentioned, to obtain a nontrivial lower bound on α using the basic construction F_q (as done in [9]) a specific methodology was needed. Our approach can then be considered as a general framework

to study rectangular matrix multiplication, which leads to a unique optimization problem that gives upper bounds on $\omega(1, 1, k)$ for any value of k .

1.1 Applications

As mentioned in the beginning of the introduction, improvements on the time complexity of rectangular matrix multiplication give faster algorithms for a multitude of computational problems. In this subsection we describe quantitatively the improvements that our new upper bounds imply for some of these problems: sparse square matrix multiplication, the all-pairs shortest paths problem and computing dynamically the transitive closure of a graph.

Sparse square matrix multiplication. Yuster and Zwick [32] have shown how fast algorithms for rectangular matrix multiplication can be used to construct fast algorithms for computing the product of two sparse square matrices (this result has been generalized to the product of sparse rectangular matrices in [15], and the case where the output matrix is also sparse has been studied in [3]). More precisely, let M and M' be two $n \times n$ matrices such that each matrix has at most m non-zero entries, where $0 \leq m \leq n^2$. Yuster and Zwick [32] showed that the product of M and M' can be computed in time

$$O\left(\min(nm, n^{\omega(1,1,\lambda_m)+o(1)}, n^{\omega+o(1)})\right),$$

where λ_m is the solution of the equation $\lambda_m + \omega(1, 1, \lambda_m) = 2 \log_n(m)$. Using the upper bounds on $\omega(1, 1, k)$ of Equation (1) with the values $\alpha < 0.294$ and $\omega < 2.376$, this gives the complexity depicted in Figure 2.

These upper bounds can be of course directly improved by using the new upper bound on ω by Vassilevska Williams [27] and the new lower bound on α given in the present work, but the improvement is small. A more significant improvement can be obtained by using directly the upper bounds on $\omega(1, 1, k)$ presented in Figure 1, which gives the new upper bounds on the complexity of sparse matrix multiplication depicted in Figure 2. For example, for $m = n^{4/3}$, we obtain complexity $O(n^{2.087})$, which is better than the original upper bound $O(n^{2.1293\dots})$ obtained from Equation (1) with $\alpha > 0.294$ and $\omega < 2.376$. Note that replacing $\omega < 2.376$ with the the best known bound $\omega < 2.3727$ only decreases the latter bound to $O(n^{2.1287\dots})$. Thus, even if the algorithms presented in the present paper do not give any improvement on ω (i.e., for the product of dense square matrices), we do obtain improvements for computing the product of two sparse square matrices.

Graph algorithms. Zwick [34] has shown how to use rectangular matrix multiplication to compute, with high probability, the all-pairs shortest paths in weighted direct graphs where the weights are bounded integers. The time complexity obtained is $O(n^{2+\mu+\epsilon})$, for any constant $\epsilon > 0$, where μ is the solution of the equation $\omega(1, 1, \mu) = 1 + 2\mu$. Using the upper bounds on $\omega(1, 1, k)$ of Equation (1) with $\alpha > 0.294$ and $\omega < 2.376$, this gives $\mu < 0.575$ and thus complexity $O(n^{2.575+\epsilon})$. This reduction to rectangular matrix multiplication is the asymptotically fastest known approach for weighted directed graphs with small integer weights.

Our results (see Table 1) show that $\omega(1, 1, 0.5302) < 2.0604$, which gives the upper bound $\mu < 0.5302$. We thus obtain the following result.

Theorem 1.2. *There exists a randomized algorithm that computes the shortest paths between all pairs of vertices in a weighted directed graph with bounded integer weights in time $O(n^{2.5302})$, where n is the number of vertices in the graph.*

Note that, even if $\omega = 2$, the complexity of Zwick's algorithm is $O(n^{2.5+\epsilon})$. In this perspective, our improvements on the complexity of rectangular matrix multiplication offer a non-negligible speed-up for the all-pairs shortest paths problem in this setting.




Figure 2: Upper bounds on the exponent for the multiplication two $n \times n$ matrices with at most m non-zero entries. The horizontal axis represents $\log_n(m)$. The dashed line represents the results by Yuster and Zwick [32] and shows that the term $n^{\omega(1,1,\lambda_m)}$ dominates the complexity when $1 \leq \log_n(m) \leq (1 + \omega)/2$. The plain line represents the improvements we obtain.

The same approach can be used to improve several other existing graph algorithms. Let us describe another example: algorithms for computing dynamically the transitive closure of a graph. Demetrescu and Italiano [12] presented a randomized algorithm for the dynamic transitive closure of directly acyclic graph with n vertices that answers queries in $O(n^\mu)$ time, and performs updates in $O(n^{1+\mu+\epsilon})$ time, for any $\epsilon > 0$. Here μ is again the solution of the equation $\omega(1, 1, \mu) = 1 + 2\mu$. This was the first algorithm for this problem with subquadratic time complexity. This result have been generalized later to general graphs, with the same bounds, by Sankowski [21]. Our new upper bounds thus show the existence of an algorithm for the dynamic transitive closure that answers queries in $O(n^{0.5302})$ time and performs updates in $O(n^{1.5302})$ time.

1.2 Overview of our techniques and organization of the paper

Before presenting an overview of the techniques used in this paper, we will give an informal description of algebraic complexity theory (the contents of which will be superseded by the formal presentation of these notions in Section 2). In this paper we will use, for any positive integer n , the notation $[n]$ to represent the set $\{1, \dots, n\}$.

Trilinear forms and bilinear algorithms. The matrix multiplication of an $m \times n$ matrix by an $n \times p$ matrix can be represented by the following trilinear form, denoted as $\langle m, n, p \rangle$:

$$\langle m, n, p \rangle = \sum_{r=1}^m \sum_{s=1}^n \sum_{t=1}^p x_{rs} y_{st} z_{rt},$$

where x_{rs} , y_{st} and z_{rt} are formal variables. This form can be interpreted as follows: the (r, t) -th entry of the product of an $m \times n$ matrix M by an $n \times p$ matrix M' can be obtained by setting $x_{ij} = M_{ij}$ for all $(i, j) \in [m] \times [n]$ and $y_{ij} = M'_{ij}$ for all $(i, j) \in [n] \times [p]$, setting $z_{rt} = 1$ and setting all the other z -variables to zero. One can then think of the z -variables as formal variables used to record the entries of the matrix product.

More generally, a trilinear form t is represented as

$$t = \sum_{i \in A} \sum_{j \in B} \sum_{k \in C} t_{ijk} x_i y_j z_k.$$

where A, B and C are three sets, x_i, y_j and z_k are formal variables and the t_{ijk} 's are coefficients in a field \mathbb{F} . Note that the set of indexes for the trilinear form $\langle m, n, p \rangle$ are $A = [m] \times [n]$, $B = [n] \times [p]$ and $C = [m] \times [p]$.

An exact (bilinear) algorithm computing t corresponds to an equality of the form

$$t = \sum_{\ell=1}^r \left(\sum_{i \in A} \alpha_{\ell i} x_i \right) \left(\sum_{j \in B} \beta_{\ell j} y_j \right) \left(\sum_{k \in C} \gamma_{\ell k} z_k \right)$$

with coefficients $\alpha_{\ell i}, \beta_{\ell j}, \gamma_{\ell k}$ in \mathbb{F} . The minimum number r such that such a decomposition exists is called the rank of the trilinear form t , and denoted $R(t)$. The rank of a trilinear form is an upper bound on the complexity of a (bilinear) algorithm computing the form: it precisely expresses the number of multiplications needed for the computation, and it can be shown that the number of additions or scalar multiplications affect the cost only in a negligible way. For any $k > 0$, the quantity $\omega(1, 1, k)$ can then be equivalently defined as follows:

$$\omega(1, 1, k) = \inf \{ \tau \in \mathbb{R} \mid R(\langle n, n, \lfloor n^k \rfloor \rangle) = O(n^\tau) \}.$$

Approximate bilinear algorithms have been introduced to take advantage of the fact that the complexity of trilinear forms (and especially of matrix multiplication) may be reduced significantly by allowing small errors in the computation. Let λ be an indeterminate over \mathbb{F} , and let $\mathbb{F}[\lambda]$ denote the set of all polynomials over \mathbb{F} in λ . Let s be any nonnegative integer. A λ -approximate algorithm for t is an equality of the form

$$\lambda^s t + \lambda^{s+1} \left(\sum_{i \in A} \sum_{j \in B} \sum_{k \in C} d_{ijk} x_i y_j z_k \right) = \sum_{\ell=1}^r \left(\sum_{i \in A} \alpha_{\ell i} x_i \right) \left(\sum_{j \in B} \beta_{\ell j} y_j \right) \left(\sum_{k \in C} \gamma_{\ell k} z_k \right)$$

for coefficients $\alpha_{\ell i}, \beta_{\ell j}, \gamma_{\ell k}, d_{ijk}$ in $\mathbb{F}[\lambda]$ and some nonnegative integer s . Informally, this means that the form t can be computed by determining the coefficient of λ^s in the right hand side. The minimum number r such that such a decomposition exists is called the border rank of t and denoted $\underline{R}(t)$. It is known that the border rank is an upper bound on the complexity of an algorithm that approximates the trilinear form, and that any such approximation algorithm can be converted into an exact algorithm with essentially the same complexity.

A sum $\sum_i t_i$ of trilinear forms is a direct sum if the t_i 's do not share variables. Informally, Schönhage's asymptotic sum inequality [23] for rectangular matrix multiplication states that, if the form t can be converted (in the λ -approximation sense) into a direct sum of c trilinear forms, each form being isomorphic to $\langle m, m, m^k \rangle$, then

$$c \cdot m^{\omega(1,1,k)} \leq \underline{R}(t).$$

This suggests that good bounds on $\omega(1, 1, k)$ can be obtained if the form t can be used to derive many independent (i.e., not sharing any variables) matrix multiplications. This approach has been applied to derive almost all new bounds on matrix multiplication since its discovery in 1981 by Schönhage.

Overview of our techniques. Our algorithm uses, as its basic construction, the trilinear form $F_q \otimes F_q$ from [10], which can be written as a sum of fifteen terms T_{ijk} , for all fifteen nonnegative integers i, j, k such that $i + j + k = 4$:

$$F_q \otimes F_q = \sum_{\substack{0 \leq i, j, k \leq 4 \\ i + j + k = 4}} T_{ijk}.$$

If the sum were direct, then, from Schönhage's asymptotic sum inequality and since an upper bound on the rank of $F_q \otimes F_q$ is easy to obtain, this would reduce the problem to the analysis of each part T_{ijk} . This is unfortunately not the case: the T_{ijk} 's share variables. To solve this problem, the basic construction is manipulated in order to obtain a direct sum, similarly to [10]. The first step is to take the N -th tensor product of the basis construction, where N is a large integer. This gives:

$$(F_q \otimes F_q)^{\otimes N} = \sum_{IJK} T_{IJK}$$

where the sum is over all triples of sequences IJK with $I, J, K \in \{0, 1, 2, 3, 4\}^N$ such that $I_\ell + J_\ell + K_\ell = 4$ for all $\ell \in \{1, \dots, N\}$. Here each form T_{IJK} is the tensor product of N forms T_{ijk} . The sum is nevertheless not yet direct. The key idea of the next step is to zero variables in order to remove some forms and obtain a sum where any two non-zero forms T_{IJK} and $T_{I'J'K'}$ are such that $I \neq I'$, $J \neq J'$ and $K \neq K'$, which will imply that the sum is direct. Moreover, in order to be able to apply Schönhage's asymptotic sum inequality, we would like all the remaining T_{IJK} to be isomorphic to the same rectangular matrix product (i.e., there should exist values m and k such that each non-zero form T_{IJK} is isomorphic to the matrix multiplication $\langle m, m, m^k \rangle$). From here our approach differs from Coppersmith and Winograd [10], since our concern is upper bounds for rectangular matrix multiplication.

We will take fifteen integers a_{ijk} and show how the form $(F_q \otimes F_q)^{\otimes N}$ can be transformed, by zeroing variables, into a *direct sum* of a large number of forms T_{IJK} in which each T_{IJK} is such that

$$T_{IJK} \cong \bigotimes_{\substack{0 \leq i, j, k \leq 4 \\ i+j+k=4}} T_{ijk}^{\otimes a_{ijk} N}. \quad (2)$$

The main difference here is that, in [10], the symmetry of square multiplication implied that the a_{ijk} 's could be taken invariant under permutation of indices, with means that only four parameters (a_{004} , a_{013} , a_{022} and a_{112}) needed to be considered. In our case, we still impose the condition $a_{ijk} = a_{ikj}$, but not more. This reduces the number of parameters to nine: a_{004} , a_{400} , a_{013} , a_{103} , a_{301} , a_{022} , a_{202} , a_{112} and a_{211} . Many nontrivial technical problems arise from this larger number of parameters. In particular, the equations that occur during the analysis do not have a unique solution and an optimization step is necessary. This is similar to the difficulties that appeared in the analysis of the basis construction $F_q^{\otimes 4}$ (for square matrix multiplication) done in [24, 27]. We will show that this further optimization step essentially imposes the additional (nonlinear) constraint $a_{013}a_{202}a_{112} = a_{103}a_{022}a_{211}$.

Showing that $(F_q \otimes F_q)^{\otimes N}$ can be transformed into a direct sum of many isomorphic forms, as claimed, was also used in previous works based on the approach by Coppersmith and Winograd. Our setting is nevertheless more general than in [10], for the following two reasons. First, our approach is asymmetric since our parameters a_{ijk} are not invariant under permutations of the indices. Second, the technical problems due to the presence of many parameters require more precise arguments. The former complication was addressed implicitly in [9], and explicitly in [13, 16]. The latter complication was addressed in [24, 27]. Here we need to deal with these two complications simultaneously, which involves a careful analysis. Instead of approaching this task directly in the language of trilinear forms, we give a graph-theoretic interpretation of it, which will make the exposition more intuitive, and also simplify the analysis. Each form T_{IJK} will correspond to one vertex in a graph, and an edge in the graph will represent the fact that two forms share one index. We will interpret the task of converting the sum $(F_q \otimes F_q)^{\otimes N}$ into a direct sum (i.e., a sum where the non-zero forms T_{IJK} do not share any index) as the task of converting the graph, by using only simple graph operations, into an edgeless subgraph. We will present algorithms solving this latter graph-theoretic task and show that a large edgeless subgraph can be obtained, which means that many forms T_{IJK} not sharing any index can be constructed from $(F_q \otimes F_q)^{\otimes N}$.

Now that we have a direct sum of many forms, each form being isomorphic to (2), the only thing to do before applying Schönhage's asymptotic sum inequality is to show that the form (2) is isomorphic to

a direct sum of matrix products $\langle m, m, m^k \rangle$. Some of the T_{ijk} 's (more precisely, all the T_{ijk} 's except T_{112} , T_{121} and T_{211}) can be analyzed in a straightforward way, since they correspond to matrix products, as originally observed in [10]. The forms T_{112} , T_{121} and T_{211} are delicate to analyze since they do not correspond to matrix products. In [10] (see also [24, 27]), they were analyzed individually through the concept of “value”, a quantity that evaluates the number of square matrix products (and their size) that can be created from the form under consideration. This is nevertheless useless for estimating $\omega(1, 1, k)$, except when $k = 1$, because the value is intrinsically symmetric (in particular, the values of T_{112} , T_{121} and T_{211} are identical), while here we are precisely interested in breaking the symmetry in order to obtain bounds for rectangular matrix multiplication. Instead, we will analyze the term

$$T_{112}^{\otimes a_{112}N} \otimes T_{121}^{\otimes a_{121}N} \otimes T_{211}^{\otimes a_{211}N} \quad (3)$$

globally. This is the key new idea leading to our new bounds on α and more generally on $\omega(1, 1, k)$ for any $k \neq 1$. Note that this difficulty was not present in previous works on rectangular matrix multiplication [9, 13, 16]: for simpler basic constructions such as F_q , all the smaller parts correspond to matrix products. We will show that T_{112} , T_{121} and T_{211} can be converted into a large number of objects called “ \mathcal{C} -tensors” in Strassen’s terminology [26]. This will be done by relying on the graph-interpretation we introduced and showing how this conversion can be interpreted as finding large cliques in a graph (this is the main reason why we developed this graph-theoretic interpretation). While the fact that the form T_{112} corresponds to a sum of \mathcal{C} -tensors was briefly mentioned in [10], and a proof sketched, we will need a complete analysis here. We will in particular rely on the fact that the \mathcal{C} -tensors obtained from T_{112} , T_{121} and T_{211} are not identical. The success of our approach comes from the discovery that these \mathcal{C} -tensors are actually “complementary”: while the \mathcal{C} -tensors obtained individually from T_{112} , T_{121} and T_{211} do not give any improvement for the exponent of rectangular matrix multiplication, their combination (i.e., the \mathcal{C} -tensors corresponding to the whole term (3)) does lead to improvements when analyzed globally by the “laser method” developed by Strassen [26]. This will show that the form (3), and thus the form (2) as well, is isomorphic to a direct sum of matrix products $\langle m, m, m^k \rangle$.

Finally, Schönhage’s asymptotic sum inequality will give an inequality, depending on the parameters a_{ijk} , that involves $\omega(1, 1, k)$. Our new upper bounds on $\omega(1, 1, k)$ are obtained by optimizing these parameters. While this is done essentially through numerical calculations, the new lower bound on α requires a more careful analysis where the optimal values of all but a few of the parameters are found analytically.

Higher powers. A natural question is whether our bounds can be improved by taking higher tensor powers of F_q as the basic algorithm, i.e., taking $F_q^{\otimes r}$ for $r > 2$. As can be expected from the analysis for $r = 2$ we have outlined above, the analysis is much more difficult than in the square case, for two main reasons. The first reason is that the construction is not symmetric and thus more parameters have to be considered. This problem can be nevertheless addressed through a systematic framework similar to the one described in [27] — this is actually quite accessible without using a computer for $r = 4$. The second, and more fundamental, reason is that the analysis of the smaller parts is different from the square case since it does not use the concept of value. We solved this problem for $r = 2$ by applying Strassen’s laser method globally on the combination of T_{112} , T_{121} and T_{211} , which is the key technical contribution of this paper. For larger values of r the same approach can in principle be used, but other techniques seem to be necessary to convert these ideas into a systematic framework.

Organization of the paper. Section 2 describes formally the notions of trilinear forms and Strassen’s laser method. Section 3 presents in details the basic construction $F_q \otimes F_q$ from [10] and some of its properties. Section 4 describes the graph-theoretic problems that will arise in the analysis of our trilinear forms. Section 5 describes our algorithm, and Section 6 shows how to use this algorithm to derive the optimization problem giving our new upper bounds on $\omega(1, 1, k)$. Finally, the optimization is done in Section 7.

2 Preliminaries

In this section we present known results about trilinear forms, Strassen's laser method and Salem-Spencer sets that we will use in this paper. We refer to [5] for an extensive treatment of the first two topics.

2.1 Trilinear forms, degeneration of tensors and Schönhage's asymptotic sum inequality

It will be convenient for us to use an abstract approach and represent trilinear forms as tensors. Our presentation is independent from what was informally defined and stated in Subsection 1.2, but describes essentially the same contents. We thus encourage the reader who encounter these notions for the first time to refer at Subsection 1.2 for concrete illustrations.

We assume that \mathbb{F} is an arbitrary field.

Let $U = \mathbb{F}^u$, $V = \mathbb{F}^v$ and $W = \mathbb{F}^w$ be three vector spaces over \mathbb{F} , where u, v and w are three positive integers. A tensor t of format (u, v, w) is an element of $U \otimes V \otimes W = \mathbb{F}^{u \times v \times w}$, where \otimes denotes the tensor product. If we fix bases $\{x_i\}$, $\{y_j\}$ and $\{z_k\}$ of U , V and W , respectively, then we can express t as

$$t = \sum_{ijk} t_{ijk} x_i \otimes y_j \otimes z_k$$

for coefficients t_{ijk} in \mathbb{F} . The tensor t can then be represented by the 3-dimensional array $[t_{ijk}]$. We will often write $x_i \otimes y_j \otimes z_k$ simply as $x_i y_j z_k$.

The tensor corresponding to the matrix multiplication of an $m \times n$ matrix by an $n \times p$ matrix is the tensor of format $(m \times n, n \times p, m \times p)$ with coefficients

$$t_{ijk} = \begin{cases} 1 & \text{if } i = (r, s), j = (s, t) \text{ and } k = (r, t) \text{ for some integers } (r, s, t) \in [m] \times [n] \times [p] \\ 0 & \text{otherwise.} \end{cases}$$

This tensor will be denoted by $\langle m, n, p \rangle$.

Another important example is the tensor $\sum_{\ell=1}^n x_\ell y_\ell z_\ell$ of format (n, n, n) . This tensor is denoted $\langle n \rangle$ and corresponds to n independent scalar products.

Let λ be an indeterminate over \mathbb{F} and consider the extension $\mathbb{F}[\lambda]$ of \mathbb{F} , i.e., the set of all polynomials over \mathbb{F} in λ . A triple of matrices $\alpha \in \mathbb{F}[\lambda]^{u' \times u}$, $\beta \in \mathbb{F}[\lambda]^{v' \times v}$ and $\gamma \in \mathbb{F}[\lambda]^{w' \times w}$ transforms $t \in \mathbb{F}^{u \times v \times w}$ into the tensor $(\alpha \otimes \beta \otimes \gamma)t \in \mathbb{F}[\lambda]^{u' \times v' \times w'}$ defined as

$$(\alpha \otimes \beta \otimes \gamma)t = \sum_{ijk} t_{ijk} \alpha(x_i) \otimes \beta(y_j) \otimes \gamma(z_j).$$

This new tensor is called a restriction of t . Intuitively, the fact that a tensor t' is a restriction of t means that an algorithm computing t can be converted into an algorithm computing t' that uses the same amount of multiplications (i.e., an algorithm with essentially the same complexity). We now give the definition of degeneration of tensors.

Definition 2.1. Let $t \in \mathbb{F}^{u \times v \times w}$ and $t' \in \mathbb{F}^{u' \times v' \times w'}$ be two tensors. We say that t' is a degeneration of t , denoted $t' \trianglelefteq t$, if there exists matrices $\alpha \in \mathbb{F}[\lambda]^{u' \times u}$, $\beta \in \mathbb{F}[\lambda]^{v' \times v}$ and $\gamma \in \mathbb{F}[\lambda]^{w' \times w}$ such that

$$\lambda^s t' + \lambda^{s+1} t'' = (\alpha \otimes \beta \otimes \gamma)t$$

for some tensor $t'' \in \mathbb{F}[\lambda]^{u' \times v' \times w'}$ and some nonnegative integer s .

Intuitively, the fact that a tensor t' is a degeneration of a tensor t means that an algorithm computing t can be converted into an "approximate algorithm" computing t' with essentially the same complexity. The notion of degeneration can be used to define the notion of border rank.

Definition 2.2. Let t be a tensor. Then $\underline{R}(t) = \min\{r \in \mathbb{N} \mid t \trianglelefteq \langle r \rangle\}$.

Let $t \in U \otimes V \otimes W$ and $t' \in U' \otimes V' \otimes W'$ be two tensors. We can naturally define the direct sum $t \oplus t'$, which is a tensor in $(U \oplus U') \otimes (V \oplus V') \otimes (W \oplus W')$, and the tensor product $t \otimes t'$, which is a tensor in $(U \otimes U') \otimes (V \otimes V') \otimes (W \otimes W')$. For any integer $c \geq 1$, we will denote the tensor $t \oplus \cdots \oplus t$ (with c occurrences of t) by $c \cdot t$ and the tensor $t \otimes \cdots \otimes t$ (with c occurrences of t) by $t^{\otimes c}$. The degeneration of tensors has the following properties.

Proposition 2.1 (Proposition 15.25 in [5]). *Let t_1, t'_1, t_2 and t'_2 be four tensors. Suppose that $t'_1 \preceq t_1$ and $t'_2 \preceq t_2$. Then $t'_1 \oplus t'_2 \preceq t_1 \oplus t_2$ and $t'_1 \otimes t'_2 \preceq t_1 \otimes t_2$.*

Schönhage's asymptotic sum inequality [23] will be one of the main tools used to prove our bounds. Its original statement is for estimating the exponent of square matrix multiplication, but it can be easily generalized to estimate the exponent of rectangular matrix multiplication as well. We will use the following form, which has been also used implicitly in [13, 16]. A proof can be found in [17].

Theorem 2.1 (Schönhage's asymptotic sum inequality). *Let k, m and c be three positive integers. Let t be a tensor such that $c \cdot \langle m, m, m^k \rangle \preceq t$. Then*

$$c \cdot m^{\omega(1,1,k)} \leq \underline{R}(t).$$

Theorem 2.1 states that, if the form t can be degenerated (i.e., approximately converted, in the sense of Definition 2.1) into a direct sum of c forms, each being isomorphic to $\langle m, m, m^k \rangle$, then the inequality $c \cdot m^{\omega(1,1,k)} \leq \underline{R}(t)$ holds. Note that this is a powerful technique, since the concepts of degeneration and border rank refer to “approximate algorithms”, while $\omega(1, 1, k)$ refers to the complexity of exact algorithms for rectangular matrix multiplication.

2.2 Strassen's laser method and \mathcal{C} -tensors

Strassen [25] introduced in 1986 a new approach, often referred as the laser method, to derive upper bounds on the exponent of matrix multiplication. To the best of our knowledge, all the applications of this method have focused so far on square matrix multiplication, in which case several simplifications can be done due to the symmetry of the problem. In this paper we will nevertheless need the full power of the laser method, and in particular the notion of \mathcal{C} -tensor introduced in [25, 26], to derive our new bounds on the exponent of rectangular matrix multiplication. The exposition below will mainly follow [5].

Let $t \in U \otimes V \otimes W$ be a tensor. Suppose that U, V and W decompose as direct sums of subspaces as follows:

$$U = \bigoplus_{i \in S_U} U_i, \quad V = \bigoplus_{j \in S_V} V_j, \quad W = \bigoplus_{k \in S_W} W_k.$$

Denote by D this decomposition. We say that t is a \mathcal{C} -tensor with respect to D if t can be written as

$$t = \sum_{(i,j,k) \in S_U \times S_V \times S_W} t_{ijk}$$

where each t_{ijk} is a tensor in $U_i \otimes V_j \otimes W_k$. The support of t is defined as

$$\text{supp}_D(t) = \{(i, j, k) \in S_U \times S_V \times S_W \mid t_{ijk} \neq 0\},$$

and the nonzero t_{ijk} 's are called the components of t . We will usually omit the reference to D when there is no ambiguity or when the decomposition does not matter.

As a simple example, consider the complete decompositions of the spaces $U = \mathbb{F}^{m \times n}$, $V = \mathbb{F}^{n \times p}$ and $W = \mathbb{F}^{m \times p}$ (i.e., their decomposition as direct sums of one-dimensional subspaces, each subspace being spanned by one element of their basis). With respect to this decomposition, the tensor of matrix multiplication $\langle m, n, p \rangle$ is a \mathcal{C} -tensor with support

$$\text{supp}_c(\langle m, n, p \rangle) = \{(r, s), (s, t), (r, t) \mid (r, s, t) \in [m] \times [n] \times [p]\}$$

where each component is trivial (i.e., isomorphic to $\langle 1, 1, 1 \rangle$). In this paper the notation $\text{supp}_c(\langle m, n, p \rangle)$ will always refer to the support of $\langle m, n, p \rangle$ with respect to this complete decomposition.

We now introduce the concept of combinatorial degeneration. A subset Δ of $S_U \times S_V \times S_W$ is called diagonal if the three projections $\Delta \rightarrow S_U$, $\Delta \rightarrow S_V$ and $\Delta \rightarrow S_W$ are injective. Let Φ be a subset of $S_U \times S_V \times S_W$. A set $\Psi \subseteq \Phi$ is a combinatorial degeneration of Φ if there exists tree functions $a: S_U \rightarrow \mathbb{Z}$, $b: S_V \rightarrow \mathbb{Z}$ and $c: S_W \rightarrow \mathbb{Z}$ such that

- for all $(i, j, k) \in \Psi$, $a(i) + b(j) + c(k) = 0$;
- for all $(i, j, k) \in \Phi \setminus \Psi$, $a(i) + b(j) + c(k) > 0$.

The most useful application of combinatorial degeneration will be the following result, which states that a sum, over indices in a diagonal combinatorial degeneration of $\text{supp}_D(t)$, of the components t_{ijk} is direct.

Proposition 2.2 (Proposition 15.30 in [5]). *Let t be \mathcal{C} -tensor with support $\text{supp}_D(t)$ and components t_{ijk} . Let $\Delta \subseteq \text{supp}_D(t)$ be a combinatorial degeneration of $\text{supp}_D(t)$ and assume that Δ is diagonal. Then*

$$\bigoplus_{(i,j,k) \in \Delta} t_{ijk} \trianglelefteq t.$$

In this work we will construct \mathcal{C} -tensors where all the components are isomorphic to $\langle m, m, m^k \rangle$ for some values m and k . Proposition 2.2 then suggests that a good bound on the exponent of rectangular matrix multiplication can be derived, via Theorem 2.1, if the support of the \mathcal{C} -tensor contains a large diagonal combinatorial degeneration. When this support is isomorphic to $\text{supp}_c(\langle e, h, \ell \rangle)$ for some positive integers e, h and ℓ , a powerful tool to construct large diagonal combinatorial degenerations is given by the following result by Strassen (Theorem 6.6 in [26]), restated in our terminology.

Proposition 2.3 ([26]). *Let e_1, e_2 and e_3 be three positive integers such that $e_1 \leq e_2 \leq e_3$. For any permutation σ of $\{1, 2, 3\}$, there exists a diagonal set $\Delta \subseteq \text{supp}_c(\langle e_{\sigma(1)}, e_{\sigma(2)}, e_{\sigma(3)} \rangle)$ with*

$$|\Delta| = \begin{cases} e_1 e_2 - \lfloor \frac{(e_1 + e_2 - e_3)^2}{4} \rfloor & \text{if } e_1 + e_2 \geq e_3 \\ e_1 e_2 & \text{otherwise} \end{cases}$$

that is a combinatorial degeneration of $\text{supp}_c(\langle e_{\sigma(1)}, e_{\sigma(2)}, e_{\sigma(3)} \rangle)$. In particular, $|\Delta| \geq \lceil 3e_1 e_2 / 4 \rceil$.

Finally, we mention that the concept of \mathcal{C} -tensor is preserved by the tensor product. We will just state this property for the restricted class of \mathcal{C} -tensors that we will encounter in this paper (for which the precise decompositions do not matter), and refer to [5] or to Section 7 in [26] for a complete treatment.

Proposition 2.4. *Let t be a \mathcal{C} -tensor with support isomorphic to $\text{supp}_c(\langle e, h, \ell \rangle)$ in which each component is isomorphic to $\langle m, n, p \rangle$. Let t' be a \mathcal{C} -tensor with support isomorphic to $\text{supp}_c(\langle e', h', \ell' \rangle)$ in which each component is isomorphic to $\langle m', n', p' \rangle$. Then $t \otimes t'$ is a \mathcal{C} -tensor with support isomorphic to $\text{supp}_c(\langle ee', hh', \ell \ell' \rangle)$ in which each component is isomorphic to $\langle mm', nn', pp' \rangle$.*

2.3 Salem-Spencer sets

Let M be a positive integer and consider $\mathbb{Z}_M = \{0, 1, \dots, M-1\}$. We say that a set $B \subseteq \mathbb{Z}_M$ has no length-3 arithmetic progression if, for any three elements b_1, b_2 and b_3 in B ,

$$b_1 + b_2 = 2b_3 \pmod{M} \iff b_1 = b_2 = b_3.$$

Salem and Spenser have shown the existence of very dense sets with no length-3 arithmetic progression.

Theorem 2.2 ([20]). *For any $\epsilon > 0$, there exists an integer M_ϵ such that, for any integer $M > M_\epsilon$, there is a set $B \subseteq \mathbb{Z}_M$ of size $|B| > M^{1-\epsilon}$ with no length-3 arithmetic progression.*

We refer to these sets as Salem-Spencer sets. They can be constructed in time polynomial in M . Note that this construction has been improved by Behrend [4], but the above statement will be enough for our purpose.

3 Coppersmith-Winograd's construction

In this section we describe the construction by Coppersmith and Winograd [10], which we will use as the basis of our algorithm, and several of its properties.

We start with the simpler construction presented in Section 7 of [10]. For any positive integer q , let us define the following trilinear form F_q , where λ is an indeterminate over \mathbb{F} .

$$\begin{aligned} F_q &= \sum_{i=1}^q \lambda^{-2}(x_0 + \lambda x_i)(y_0 + \lambda y_i)(z_0 + \lambda z_i) - \\ &\lambda^{-3}(x_0 + \lambda^2 \sum_{i=1}^q x_i)(y_0 + \lambda^2 \sum_{i=1}^q y_i)(z_0 + \lambda^2 \sum_{i=1}^q z_i) + \\ &(\lambda^{-3} - q\lambda^{-2})(x_0 + \lambda^3 x_{q+1})(y_0 + \lambda^3 y_{q+1})(z_0 + \lambda^3 z_{q+1}) \end{aligned}$$

In this trilinear form the x -variables are x_0, x_1, \dots, x_{q+1} . Similarly, the number of y -variables is $(q+2)$ and the number of z -variables is $(q+2)$ as well. Define the form

$$F'_q = \sum_{i=1}^q (x_0 y_i z_i + x_i y_0 z_i + x_i y_i z_0) + x_0 y_0 z_{q+1} + x_0 y_{q+1} z_0 + x_{q+1} y_0 z_0.$$

It is easy to check that the form F_q can be written as $F_q = F'_q + \lambda \cdot F''_q$, where F''_q is a polynomial in λ and in the x -variables, y -variables and z -variables. In the language of Section 2, this means that $F'_q \trianglelefteq F_q$ and, informally, this means that an algorithm computing F_q can be converted into an algorithm computing F'_q with the same complexity. Note that $\underline{R}(F_q) \leq q+2$ since, by definition, F_q is the sum of $q+2$ products.

A more complex construction is proposed in Section 8 of [10]. It is obtained by taking the tensor product of F_q by itself. By Proposition 2.1 we know that $F'_q \otimes F'_q \trianglelefteq F_q \otimes F_q$. Consider the tensor product of F'_q by itself:

$$\begin{aligned} F'_q \otimes F'_q &= T_{004} + T_{040} + T_{400} + T_{013} + T_{031} + T_{103} + T_{130} + T_{301} + T_{310} + \\ &T_{022} + T_{202} + T_{220} + T_{112} + T_{121} + T_{211} \end{aligned}$$

where

$$\begin{aligned} T_{004} &= x_{0,0}^0 y_{0,0}^0 z_{q+1,q+1}^4 \\ T_{013} &= \sum_{i=1}^q x_{0,0}^0 y_{i,0}^1 z_{i,q+1}^3 + \sum_{k=1}^q x_{0,0}^0 y_{0,k}^1 z_{q+1,k}^3 \\ T_{022} &= x_{0,0}^0 y_{q+1,0}^2 z_{0,q+1}^2 + x_{0,0}^0 y_{0,q+1}^2 z_{q+1,0}^2 + \sum_{i,k=1}^q x_{0,0}^0 y_{i,k}^2 z_{i,k}^2 \\ T_{112} &= \sum_{i=1}^q x_{i,0}^1 y_{i,0}^1 z_{0,q+1}^2 + \sum_{k=1}^q x_{0,k}^1 y_{0,k}^1 z_{q+1,0}^2 + \sum_{i,k=1}^q x_{i,0}^1 y_{0,k}^1 z_{i,k}^2 + \sum_{i,k=1}^q x_{0,k}^1 y_{i,0}^1 z_{i,k}^2 \end{aligned}$$

and the other eleven terms are obtained by permuting the indexes of the x -variables, the y -variables and z -variables in the above expressions (e.g., $T_{040} = x_{0,0}^0 y_{q+1,q+1}^4 z_{0,0}^0$ and $T_{400} = x_{q+1,q+1}^4 y_{0,0}^0 z_{0,0}^0$).

Let us describe in more details the notations used here. The number of x -variables is $(q + 2)^2$. They are indexed as $x_{i,k}$, for $i, k \in \{0, 1, \dots, q + 1\}$. The superscript is assigned in the following way: the variable $x_{0,0}$ has superscript 0, the variables in $\{x_{i,0}, x_{0,k}\}_{1 \leq i, j \leq q}$ have superscript 1, the variables in $\{x_{q+1,0}, x_{i,k}, x_{0,q+1}\}_{1 \leq i, j \leq q}$ have superscript 2, the variables in $\{x_{q+1,k}, x_{i,q+1}\}_{1 \leq i, j \leq q}$ have superscript 3 and the variable $x_{q+1,q+1}$ has superscript 4. Note that the superscript is completely determined by the subscript. Similarly, the number of y -variables is $(q + 2)^2$, and the number of z -variables is $(q + 2)^2$ as well. The y -variables and the z -variables are assigned subscripts and superscripts exactly as for the x -variables. Observe that any term xyz that appears in T_{ijk} is such that x has superscript i , y has superscript j and z has superscript k .

We thus obtain

$$\sum_{\substack{0 \leq i, j, k \leq 4 \\ i+j+k=4}} T_{ijk} \leq F_q \otimes F_q.$$

Moreover we know that $\underline{R}(F_q \otimes F_q) \leq (q + 2)^2$, since $F_q \otimes F_q$ can be written using $(q + 2)^2$ multiplications.

We will later need to analyze all the forms T_{ijk} . It happens, as observed in [10], that most of these forms (all the forms except T_{112} , T_{121} and T_{211}) can be analyzed in a straightforward way, since they are isomorphic to the following matrix products:

$$\begin{aligned} T_{004} \cong T_{040} \cong T_{400} &\cong \langle 1, 1, 1 \rangle \\ T_{013} \cong T_{031} &\cong \langle 1, 1, 2q \rangle \\ T_{103} \cong T_{301} &\cong \langle 2q, 1, 1 \rangle \\ T_{130} \cong T_{310} &\cong \langle 1, 2q, 1 \rangle \\ T_{022} &\cong \langle 1, 1, q^2 + 2 \rangle \\ T_{202} &\cong \langle q^2 + 2, 1, 1 \rangle \\ T_{220} &\cong \langle 1, q^2 + 2, 1 \rangle. \end{aligned}$$

This can be seen from the definition of the trilinear form (or the tensor) corresponding to matrix multiplication described in Section 2. For example, the form T_{013} is isomorphic to the tensor $\sum_{\ell=1}^{2q} x_0 y_\ell z_\ell = \langle 1, 1, 2q \rangle$, which represents the product of a 1×1 matrix (a scalar) by a $1 \times 2q$ matrix (a row).

4 Graph-Theoretic Problems

In this section we describe and solve several graph-theoretic problems that will arise in the analysis of our trilinear forms. While the presentation given here is independent from the remaining of the paper, the reader may prefer to read Section 5 before going through this section.

4.1 Problem setting

Let τ be a fixed positive integer. Let N be a large integer and define the set

$$\Lambda = \{(I, J, K) \in [\tau]^N \times [\tau]^N \times [\tau]^N \mid I_\ell + J_\ell + K_\ell = \tau \text{ for all } \ell \in \{1, \dots, N\}\}.$$

Define the three coordinate functions $f_1, f_2, f_3: [\tau]^N \times [\tau]^N \times [\tau]^N \rightarrow [\tau]^N$ as follows.

$$\begin{aligned} f_1((I, J, K)) &= I \\ f_2((I, J, K)) &= J \\ f_3((I, J, K)) &= K \end{aligned}$$

From the definition of Λ , two distinct elements in Λ cannot agree on more than one coordinate. Since this simple observation will be crucial in our analysis, we state it explicitly as follows.

Fact 1. Let u and v be two elements in Λ . If $f_i(u) = f_i(v)$ for more than one index $i \in \{1, 2, 3\}$, then $u = v$.

Let U be a subset of Λ such that there exist integers $\mathcal{N}_1, \mathcal{N}_2$ and \mathcal{N}_3 for which the following property holds: for any $I \in [\tau]^N$,

$$\begin{aligned} |\{u \in U \mid f_1(u) = I\}| &\in \{0, \mathcal{N}_1\} \\ |\{u \in U \mid f_2(u) = I\}| &\in \{0, \mathcal{N}_2\} \\ |\{u \in U \mid f_3(u) = I\}| &\in \{0, \mathcal{N}_3\}. \end{aligned}$$

This means that, for any $i \in \{1, 2, 3\}$ and any $I \in [\tau]^N$, the size of the set $f_i(I)^{-1}$ is either 0 or \mathcal{N}_i . Let us write $|f_1(U)| = T_1$, $|f_2(U)| = T_2$ and $|f_3(U)| = T_3$. It is easy to see that

$$|U| = T_1 \mathcal{N}_1 = T_2 \mathcal{N}_2 = T_3 \mathcal{N}_3.$$

We are interested in stating asymptotic results holding when N goes to infinity. Through this section the \mathcal{N}_i 's and the T_i 's will be strictly increasing functions of N .

Let G be the (simple and undirected) graph with vertex set U in which two distinct vertices u and v are connected if and only there exists one index $i \in \{1, 2, 3\}$ such that $f_i(u) = f_i(v)$. The goal will be to modify the graph G to obtain a subgraph satisfying some specific properties. The only modification we allow is to remove all the vertices with a given sequence at a given position: given a sequence $I \in [\tau]^N$ and a position $s \in \{1, 2, 3\}$, remove all the vertices u (if any) such that $f_s(u) = I$. We call such an operation a removal operation. The reason why only such removal operations are allowed is that they will correspond, when considering trilinear forms, to setting to zero some variables, which is one of the only operations that can be performed on trilinear forms.

While not stated explicitly in graph-theoretic terms, the key technical result by Coppersmith and Winograd [10] is a method to convert, when $\mathcal{N}_1 = \mathcal{N}_2 = \mathcal{N}_3$, the graph G into an edgeless graph that still contains a non-negligible fraction of the vertices. We state this result in the following theorem.

Theorem 4.1 ([10]). *Suppose that $\mathcal{N}_1 = \mathcal{N}_2 = \mathcal{N}_3$. Then, for any constant $\epsilon > 0$, the graph G can be converted, with only removal operations, into an edgeless graph with $\Omega\left(\frac{T_1}{\mathcal{N}_1^\epsilon}\right)$ vertices.*

In this section we will give several generalizations of this result. In Subsection 4.2 we state our generalizations. Then, in Subsections 4.3–4.6, we describe the algorithms and prove the results. We stress that, in this section as in [10], the number of removal operations (i.e., the time complexity of the algorithms) is irrelevant. This is because, in the applications of these results to matrix multiplication, the parameter N will be treated as a large constant independent of the size of the matrices considered.

4.2 Statement of our results

The generalization we consider assume the existence of a known set $U^* \subseteq U$ such that

- $|f_i(U^*)| = T_i$ for each $i \in \{1, 2, 3\}$;
- there exist integers $\mathcal{N}_1^*, \mathcal{N}_2^*$ and \mathcal{N}_3^* such that

$$|\{u \in U \mid f_i(u) = I\}| = \mathcal{N}_i \Leftrightarrow |\{u \in U^* \mid f_i(u) = I\}| = \mathcal{N}_i^*$$

for each $I \in [\tau]^N$ and each $i \in \{1, 2, 3\}$.

Note that we have necessarily $T_1 \mathcal{N}_1^* = T_1 \mathcal{N}_2^* = T_1 \mathcal{N}_3^*$.

The first problem considered is again to convert the graph G into an edgeless subgraph that contains a non-negligible fraction of the vertices using only removal operations, but we additionally require that all the remaining vertices are in U^* . Our first result is the following theorem.

Theorem 4.2. *For any constant $\epsilon > 0$, the graph G can be converted, with only removal operations, into an edgeless graph with*

$$\Omega\left(\frac{T_1 \mathcal{N}_1^*}{(\mathcal{N}_1 + \mathcal{N}_2 + \mathcal{N}_3)^{1+\epsilon}}\right)$$

vertices, all of them being in U^ .*

Theorem 4.1 is a special case of Theorem 4.2, for $U^* = U$. The case $\mathcal{N}_1 = \mathcal{N}_2 = \mathcal{N}_3$ has been proved implicitly by Stothers [24] and Vassilevska Williams [27].

Our second problem deals with another kind of conversion. Remember that, from the definition of the graph G and Fact 1, for any edge connecting two vertices u and v in G there exists exactly one index $i \in \{1, 2, 3\}$ such that $f_i(u) = f_i(v)$. Let us define the concept of 1-clique as follows.

Definition 4.1. *A 1-clique in the graph G is a set $U' \subseteq U$ for which there exists a sequence $I \in [\tau^N]$ such that $f_1(u) = I$ for all $u \in U'$. The size of the 1-clique is $|U'|$.*

Our second result shows how to convert, using only removal operations, the graph G into a graph with vertices in U^* that is the disjoint union of many large 1-cliques. The formal statement follows.

Theorem 4.3. *Suppose that $\mathcal{N}_1 \geq \mathcal{N}_2 \geq \mathcal{N}_3$. Assume that $\frac{\mathcal{N}_2 T_1}{\mathcal{N}_1^*} + \frac{\mathcal{N}_2}{T_1} < \frac{1}{1024}$. Then, for any constant $\epsilon > 0$, the graph G can be converted, with only removal operations, into a graph satisfying the following conditions:*

- *all the vertices of the graph are in U^* ;*
- *each connected component is a 1-clique (i.e., the graph is a disjoint union of 1-cliques);*
- *among these 1-cliques, there are $\Omega\left(\frac{T_1}{\mathcal{N}_2^\epsilon}\right)$ 1-cliques that have size $\Omega\left(\frac{\mathcal{N}_1^*}{\mathcal{N}_2}\right)$.*

Note that, when only removal operations are allowed, the graph obtained is necessary a subgraph of G induced by a subset of its vertices. Theorem 4.3 thus states that there exist 1-cliques $U_r \subseteq U^*$ such that the graph obtained after the removal operations is the subgraph of G induced by $\cup_r U_r$, at least $\Omega(T_1/\mathcal{N}_2^\epsilon)$ of these U_r 's have size $\Omega(\mathcal{N}_1^*/\mathcal{N}_2)$, and for any $r \neq r'$ there is no edge with one extremity in U_r and the other extremity in $U_{r'}$. We mention that the constant $1/1024$ in the assumption of Theorem 4.3 is chosen only for concreteness. The same theorem actually holds even for weaker conditions on \mathcal{N}_1^*, T_1 and \mathcal{N}_2 , but this simpler version will be sufficient for our purpose.

4.3 Choice of the weight functions

Let M be a large prime number that will be chosen later. We take a Salem-Spencer set Γ with $|\Gamma| \geq M^{1-\epsilon}$, which existence is guaranteed by Theorem 2.2. Similarly to [10], we take $N + 2$ integers $\omega_0, \omega_1, \dots, \omega_{N+1}$ uniformly at random in $\mathbb{Z}_M = \{0, 1, \dots, M - 1\}$ and define three hash functions $b_1, b_2, b_3: [\tau]^N \rightarrow \mathbb{Z}_M$ as follows.

$$\begin{aligned} b_1(I) &= \omega_0 + \sum_{j=1}^N I_j \omega_j \pmod{M} \\ b_2(I) &= \omega_{N+1} + \sum_{j=1}^N I_j \omega_j \pmod{M} \\ b_3(I) &= \frac{1}{2} \times \left(\omega_0 + \omega_{N+1} + \sum_{j=1}^N (\tau - I_j) \omega_j \right) \pmod{M} \end{aligned}$$

A property of these functions is that, for a fixed sequence $I \in [\tau]^N$, the value $b_i(I)$ is uniformly distributed in \mathbb{Z}_M , for each $i \in \{1, 2, 3\}$. Note that the term ω_0 is not used in [10], but we introduce it to obtain a uniform distribution even if I is the all-zero sequence.

Let us introduce the notion of a compatible vertex.

Definition 4.2. A vertex $u \in U$ is compatible if $b_i(f_i(u)) \in \Gamma$ for each $i \in \{1, 2, 3\}$. Let V be the set of vertices in U that are compatible, and denote $V^* = V \cap U^*$.

We stress that the definition of compatibility (and thus the definitions of V and V^* too) depend of the choice of Γ and of the weights ω_i . Using the fact that Γ is a Salem-Spencer set we can give the following simple but very useful characterization of compatible vertices.

Lemma 4.1. Let $i \in \{1, 2, 3\}$ and $i' \in \{1, 2, 3\}$ be any two distinct indexes. Then a vertex $u \in U$ is compatible if and only if $b_i(f_i(u)) = b_{i'}(f_{i'}(u)) = b$ for some $b \in \Gamma$.

Proof. Let us take a vertex $u \in U$, and write $f_1(u) = I$, $f_2(u) = J$ and $f_3(u) = K$.

Note that, since $I_\ell + J_\ell + K_\ell = \tau$ for any index $\ell \in \{1, \dots, N\}$, the equality $b_1(I) + b_2(J) - 2b_3(K) = 0 \pmod{M}$ always holds (i.e., holds for all choices of the weights ω_j).

Suppose that $b_i(f_i(u)) = b_{i'}(f_{i'}(u)) = b$ for some $b \in \Gamma$, where i and i' are distinct. For instance, suppose that $i = 1$ and $i' = 2$. Then, since M is a prime, the above property implies that $b_3(K) = b$, which means that u is compatible. The same conclusion is true for the other choices of i and i' .

Now suppose that u is compatible. From the definition of a Spencer-Salem set, we can conclude that there exists an element $b \in \Gamma$ such that $b_1(I) = b_2(J) = b_3(K) = b$. \square

4.4 The first pruning

Similarly to [10], the first pruning simply eliminates all the nodes $u \in U$ that are not compatible. Note that this can be done using removal operations: for each $i \in \{1, 2, 3\}$, we remove all the vertices w such that $f_i(w) \notin b_i^{-1}(\Gamma)$. The vertices remaining are precisely those in V . Among those remaining vertices, the vertices in U^* are precisely those in $V^* = V \cap U^*$.

We now evaluate the expectation of $|V^*|$. The proof is very similar to what was shown in [10] for the case $U = U^*$, and to what was shown in [24, 27] for $\mathcal{N}_1^* = \mathcal{N}_2^* = \mathcal{N}_3^*$.

Lemma 4.2. $\mathbb{E}[|V^*|] = \frac{T_1 \mathcal{N}_1^* |\Gamma|}{M^2}$.

Proof. We use Lemma 4.1 with $i = 1$ and $i' = 2$. Remember that $|U^*| = T_1 \mathcal{N}_1^*$. For each vertex $u \in U^*$ and each value $b \in \mathbb{Z}_M$, the probability that $b_1(f_1(u)) = b_2(f_2(u)) = b$ is $1/M^2$. Note that the two events $b_1(f_1(u)) = b$ and $b_2(f_2(u)) = b$ are independent even when $f_1(u) = f_2(u)$ due to the terms ω_0 and ω_{N+1} in the hash functions. \square

Let E be the edge set of the subgraph of G induced by V : it consists of all edges connecting two distinct vertices u and v in V such that $f_i(u) = f_i(v)$ for some $i \in \{1, 2, 3\}$. Let $E' \subseteq E$ be the subset of edges in E with (at least) one extremity in V^* . Let $E'' \subseteq E'$ be the subset of edges in E' connecting two vertices u and v such that $f_1(u) \neq f_1(v)$ (which means that either $f_2(u) = f_2(v)$ or $f_3(u) = f_3(v)$). The following lemma gives upper bounds on the expectations of $|E'|$ and $|E''|$. The proof can be considered as a generalization of similar statements in [10, 24, 27].

Lemma 4.3. $\mathbb{E}[|E'|] \leq \frac{T_1 \mathcal{N}_1^* (\mathcal{N}_1 + \mathcal{N}_2 + \mathcal{N}_3 - 3) |\Gamma|}{M^3}$ and $\mathbb{E}[|E''|] \leq \frac{T_1 \mathcal{N}_1^* (\mathcal{N}_2 + \mathcal{N}_3 - 2) |\Gamma|}{M^3}$.

Proof. We show that, for any index $i \in \{1, 2, 3\}$, the expected number of ordered pairs (u, v) with $u \in V^*$ and $v \in V \setminus \{u\}$ such that $f_i(u) = f_i(v)$ is exactly

$$\frac{T_1 \mathcal{N}_1^* (\mathcal{N}_i - 1) |\Gamma|}{M^3}.$$

The expectation of the number of edges (i.e., unordered pairs) is necessarily smaller.

The factor $T_1 \mathcal{N}_1^*$ counts the number of vertices in U^* . For any vertex $u \in U^*$, there are exactly $\mathcal{N}_i - 1$ vertices $v \in U \setminus \{u\}$ such that $f_i(v) = f_i(u)$.

Let u be a vertex in U^* and v be a vertex in $U \setminus \{u\}$ such that $f_i(v) = f_i(u)$. Take another index $i' \in \{1, 2, 3\} \setminus \{i\}$ arbitrarily. From Lemma 4.1, both u and v are in V if and only if

$$b_i(f_i(u)) = b_{i'}(f_{i'}(u)) = b_{i'}(f_{i'}(v)) = b$$

for some element $b \in \Gamma$. This happens with probability $|\Gamma|/M^3$ since the random variables $b_i(f_i(u))$, $b_{i'}(f_{i'}(u))$, and $b_{i'}(f_{i'}(v))$ are mutually independent. From the linearity of the expectation, the expected number of ordered pairs is $T_1 \mathcal{N}_1^* (\mathcal{N}_i - 1)$ times this probability. \square

Lemmas 4.2 and 4.3 focused on expected values and were proven, essentially, by the linearity of the expectation. This was sufficient for the applications to square matrix multiplication presented in [10, 24, 27], and this will be sufficient for proving Theorem 4.2 as well. Since, in order to prove Theorem 4.3, we will need a more precise analysis of the behavior of the random variables considered, we now prove the following lemma, which gives a lower bound on the probability that the subgraph induced by V^* has many large 1-cliques.

Lemma 4.4. *With probability (on the choice of the weights ω_i) at least $1 - \frac{4MT_1}{\mathcal{N}_1^*} - \frac{4M}{T_1|\Gamma|}$, there exists a set $R \subseteq [\tau]^N$ satisfying the following two conditions:*

- $|R| \geq \frac{T_1|\Gamma|}{2M}$;
- for any $I \in R$, there exist at least $\frac{\mathcal{N}_1^*}{2M}$ vertices $u \in V^*$ such that $f_1(u) = I$.

Proof. Let us consider the set $f_1(U^*) = \{f_1(u) \mid u \in U^*\}$. For each element $I \in f_1(U^*)$, define the set

$$S_I = \{u \in U^* \mid f_1(u) = I\}.$$

Note that $|f_1(U^*)| = T_1$, and $|S_I| = \mathcal{N}_1^*$ for each $I \in f_1(U^*)$.

Fix an element $I \in f_1(U^*)$ and define

$$X_I = |\{u \in S_I \mid b_2(f_2(u)) = b_1(I)\}|.$$

This random variable represents the number of vertices from S_I that are mapped by $b_2 \circ f_2$ into $b_1(I)$. For any vertex $u = (I, J, K) \in S_I$, we have the equivalence

$$b_2(f_2(u)) = b_1(I) \iff \sum_{i=1}^N (J_i - I_i) \omega_i = \omega_0 - \omega_{N+1}.$$

Thus the probability of the event $b_2(f_2(u)) = b_1(I)$ is $1/M$. Thus $\mathbb{E}[X_I] = \frac{\mathcal{N}_1^*}{M}$. Note that, for any two distinct elements u and v in S_I , the two events $b_2(f_2(u)) = b_1(I)$ and $b_2(f_2(v)) = b_1(I)$ are independent. From this pairwise independence, $\text{var}[X_I] = \frac{\mathcal{N}_1^*}{M} (1 - \frac{1}{M})$. This gives

$$\Pr \left[\left| X_I - \frac{\mathcal{N}_1^*}{M} \right| \geq \frac{\mathcal{N}_1^*}{2M} \right] \leq \Pr \left[\left| X_I - \frac{\mathcal{N}_1^*}{M} \right| \geq \sqrt{\frac{\mathcal{N}_1^*}{4M}} \cdot \sqrt{\text{var}(X_I)} \right] \leq \frac{4M}{\mathcal{N}_1^*},$$

where the second inequality is obtained by Chebyshev's inequality.

By the union bound we can conclude that

$$\Pr \left[|X_I| \geq \frac{\mathcal{N}_1^*}{2M} \text{ for all } I \in f_1(U^*) \right] \geq 1 - \frac{4MT_1}{\mathcal{N}_1^*}.$$




Figure 3: The second pruning (first version)

Define $R = \{I \in f_1(U^*) \mid b_1(I) \in \Gamma\}$. For each element $I \in f_1(U^*)$, the variable $b_1(I)$ is distributed uniformly at random in \mathbb{Z}_M . Moreover, the variables $b_1(I)$'s are pairwise independent. Thus, by Chebyshev's inequality, we obtain

$$\Pr \left[|R| \geq \frac{T_1 |\Gamma|}{2M} \right] \geq 1 - \frac{4M}{T_1 |\Gamma|}.$$

From the union bound we can conclude that, with probability at least $1 - \frac{4MT_1}{\mathcal{N}_1^*} - \frac{4M}{T_1 |\Gamma|}$, the inequality $|R| \geq T_1 |\Gamma| / (2M)$ holds and simultaneously, for each element $I \in R$, there exist at least $\mathcal{N}_1^* / (2M)$ vertices $u \in U^*$ such that $f_1(u) = I$ and $b_2(f_2(u)) = b_1(I)$. These vertices are in V^* by Lemma 4.1. \square

4.5 The second pruning: first version and proof of Theorem 4.2

In this subsection M is an arbitrary prime number such that $2(\mathcal{N}_1 + \mathcal{N}_2 + \mathcal{N}_3) < M < 4(\mathcal{N}_1 + \mathcal{N}_2 + \mathcal{N}_3)$.

The first pruning has transformed the graph G into the subgraph induced by V . The second pruning, similarly to [10], will further modify this subgraph by removing vertices in order to obtain a subgraph consisting of isolated vertices from V^* (i.e., an edgeless graph). This is done by constructing greedily a set $L \subseteq V^*$ of isolated vertices. Initially $L = \emptyset$ and, at each iteration, either one remaining vertex in $V^* \setminus L$ will be added to L or several vertices in V will be removed. This will be repeated until there is no remaining vertex in $V^* \setminus L$. Finally, all the remaining vertices not in L will be removed. The detailed procedure is described in Figure 3, where \bar{V} represents the set of remaining vertices (initially $\bar{V} = V$). Note that the procedure slightly differs from what was done in [10, 24, 27] since we need to take in consideration the asymmetry of the problem.

Let \bar{V}_f denote the contents of \bar{V} at the end of the procedure. The following proposition, shown using the same ideas as in [10], shows that what we obtain is a large set of isolated vertices from V^* .

Proposition 4.1. *The subgraph of G induced by \bar{V}_f is an edgeless graph. Moreover, $\bar{V}_f \subseteq V^*$ and the expectation of $|\bar{V}_f|$ (over the choices of ω_j) is*

$$\Omega \left(\frac{T_1 \mathcal{N}_1^*}{(\mathcal{N}_1 + \mathcal{N}_2 + \mathcal{N}_3)^{1+\epsilon}} \right).$$

Proof. Let L_f denote the contents of L at the end of the procedure. First observe that $\overline{V}_f \subseteq L_f$, due to Step 3. Note that any vertex added to L cannot be later removed from \overline{V} , since it has no neighbor. Thus $L_f \subseteq \overline{V}_f$, and we conclude that $\overline{V}_f = L_f$. This shows in particular that $\overline{V}_f \subseteq V^*$. Moreover, since each vertex in L has no neighbor, the subgraph induced by \overline{V}_f is edgeless.

To prove the second part, we will show an upper bound on the number of vertices in V^* removed from \overline{V} during the loop of Step 2. The bound will be obtained by considering the number of edges from E' remaining in the subgraph induced by \overline{V} .

Let us consider what happens during Step 2.2. Let $u \in (\overline{V} \cap V^*) \setminus L$ be the vertex currently examined. Suppose that another vertex in \overline{V} sharing one index with u is found. For example, suppose that we find another vertex $v \in \overline{V}$ with $f_1(v) = f_1(u)$. Let $S = \{w \in \overline{V} \cap V^* \mid f_2(w) = f_2(u)\}$ be the set of vertices in V^* eliminated by the consequent removal operation. Observe that this removal operation will eliminate at least $\binom{|S|}{2} + 1$ new edges from E' : the edges between two vertices in S , and the edge connecting u and v . Since $\binom{|S|}{2} + 1 \geq |S|$, the number of vertices in V^* removed during one execution of Step 2.2 is at most the number of edges eliminated from E' .

The total number of vertices from V^* that are removed by the procedure during the loop of Step 2 is thus at most $|E'|$, which means that $|\overline{V}_f \cap V^*| \geq |V^*| - |E'|$. Since $\overline{V} \subseteq V^*$, Lemmas 4.2 and 4.3 imply that the expected number of vertices in \overline{V}_f is at least

$$\frac{T_1 \mathcal{N}_1^* |\Gamma|}{M^2} \left(1 - \frac{(\mathcal{N}_1 + \mathcal{N}_2 + \mathcal{N}_3 - 3)}{M} \right) \geq \frac{T_1 \mathcal{N}_1^* |\Gamma|}{2M^2},$$

where the inequality follows from the choice of M . Since $M = O(\mathcal{N}_1 + \mathcal{N}_2 + \mathcal{N}_3)$ and $|\Gamma| \geq M^{1-\epsilon}$, we conclude that $\mathbb{E}[|\overline{V}_f|] = \Omega\left(\frac{T_1 \mathcal{N}_1^*}{(\mathcal{N}_1 + \mathcal{N}_2 + \mathcal{N}_3)^{1+\epsilon}}\right)$. \square

Theorem 4.2 follows from Proposition 4.1 by fixing a choice of the weights $\omega_0, \omega_1, \dots, \omega_{N+1}$ for which $|\overline{V}_f| \geq \mathbb{E}[|\overline{V}_f|]$ (such a choice necessarily exists from the definition of the expectation).

4.6 The second pruning: second version and proof of Theorem 4.3

In this subsection M is an arbitrary prime such that $64\mathcal{N}_2 < M < 128\mathcal{N}_2$.

The first version of the pruning described in Subsection 4.5 was designed to obtain an edgeless subgraph of G . In this subsection we describe how to modify it to obtain a union of many large 1-cliques instead. The detailed procedure of the new pruning algorithm is described in Figure 4. The only difference is that, at Step 2.2, a vertex is not added in L only if it is connected to another vertex with the same second or third index.

Let \overline{V}_f denote the contents of \overline{V} at the end of the procedure. By slightly modifying the arguments of the previous subsection, it is easy to see that the resulting graph has only vertices from V^* and is a disjoint union of 1-cliques (i.e., each connected component is a 1-clique).

Proposition 4.2. *The subgraph of G induced by \overline{V}_f is a disjoint union of 1-cliques. Moreover, $\overline{V}_f \subseteq V^*$ and $|\overline{V}_f| \geq |V^*| - |E''|$.*

Proof. Let L_f denote the contents of L at the end of the procedure. Due to Step 3, we know that $\overline{V}_f \subseteq L_f$. Moreover, any vertex added to L cannot be later removed from \overline{V} , since it has no neighbor with the same second or third index. Thus $L_f \subseteq \overline{V}_f$, and we conclude that $\overline{V}_f = L_f$, which shows in particular that $\overline{V}_f \subseteq V^*$. Furthermore, since each vertex in L has no neighbor with the same second or third index, the subgraph induced by \overline{V}_f is a disjoint union of 1-cliques.

The inequality $|\overline{V}_f| \geq |V^*| - |E''|$ is obtained as in the proof of Proposition 4.1, but replacing the edge set E' by E'' . \square

We now give the proof of Theorem 4.3, by using Lemmas 4.3 and 4.4 to evaluate the size and the numbers of 1-cliques in the resulting graph.




Figure 4: The second pruning (second version)

Proof of Theorem 4.3. Using Lemma 4.3 with the value $M > 64\mathcal{N}_2$ and Markov's bound, we conclude that

$$\Pr \left[|E''| \leq \frac{T_1 \mathcal{N}_1^* |\Gamma|}{16M^2} \right] \geq \frac{1}{2}.$$

Note that the conditions $M < 128\mathcal{N}_2$ and $\frac{\mathcal{N}_2 T_1}{\mathcal{N}_1^*} + \frac{\mathcal{N}_2}{T_1} < \frac{1}{1024}$ imply that

$$1 - \frac{4MT_1}{\mathcal{N}_1^*} - \frac{4M}{T_1 |\Gamma|} > 1 - 512 \left(\frac{\mathcal{N}_2 T_1}{\mathcal{N}_1^*} + \frac{\mathcal{N}_2}{T_1 |\Gamma|} \right) > 1 - 512 \left(\frac{\mathcal{N}_2 T_1}{\mathcal{N}_1^*} + \frac{\mathcal{N}_2}{T_1} \right) > \frac{1}{2}.$$

Thus the probability that a set R as in Lemma 4.4 exists and the inequality $|E''| \leq \frac{T_1 \mathcal{N}_1^* |\Gamma|}{16M^2}$ simultaneously holds is positive. There then exists a choice of the weights $\omega_0, \omega_1, \dots, \omega_{N+1}$ such that this happens. Let us take such a choice.

From Proposition 4.2 we know that at most $|E''|$ vertices in V^* are removed by the second pruning. Then there exists a set $R' \subseteq R$ of size $|R'| \geq T_1 |\Gamma| / (4M)$ such that, for any $r' \in R'$, there are at least $\mathcal{N}_1^* / (4M)$ vertices u with $f_1(u) = r'$ remaining after the second pruning. This is because, otherwise, from the properties of the set R stated in Lemma 4.4 it would be necessary to remove more than $T_1 \mathcal{N}_1^* |\Gamma| / (16M^2)$ vertices during the second pruning. \square

5 Algorithm for Rectangular Matrix Multiplication

In this section we present our algorithm, which essentially consists in the two algorithmic steps described in Subsections 5.2 and 5.4. We first start by explaining the construction we will use.

5.1 Our construction

Let $a_{004}, a_{400}, a_{013}, a_{103}, a_{301}, a_{022}, a_{202}, a_{112}, a_{211}$ be nine arbitrary positive¹ rational numbers such that

$$2a_{004} + a_{400} + 2a_{013} + 2a_{103} + 2a_{301} + a_{022} + 2a_{202} + 2a_{112} + a_{211} = 1$$

¹The hypothesis that each a_{ijk} is not zero is made only for convenience (all the bounds presented in this paper are obtained using positive values for these parameters). More specifically, this hypothesis is used only when approximating quantities like $(a_{ijk}N)!$ using Stirling's inequality. Without the hypothesis it would be necessary to treat the (trivial) case $a_{ijk} = 0$ separately.

and

$$a_{013}a_{202}a_{112} = a_{103}a_{022}a_{211}.$$

Let us define rational numbers $A_0, A_1, A_2, A_3, A_4, B_0, B_1, B_2, B_3, B_4$ as follows:

$$\begin{aligned} A_0 &= 2a_{004} + 2a_{013} + a_{022} \\ A_1 &= 2a_{103} + 2a_{112} \\ A_2 &= 2a_{202} + a_{211} \\ A_3 &= 2a_{301} \\ A_4 &= a_{400} \\ B_0 &= a_{004} + a_{400} + a_{103} + a_{301} + a_{202} \\ B_1 &= a_{013} + a_{301} + a_{112} + a_{211} \\ B_2 &= a_{022} + a_{202} + a_{112} \\ B_3 &= a_{013} + a_{103} \\ B_4 &= a_{004}. \end{aligned}$$

Note that $\sum_{i=0}^4 A_i = \sum_{i=0}^4 B_i = 1$.

It will be convenient to define six additional numbers $a_{040}, a_{031}, a_{130}, a_{310}, a_{220}$ and a_{121} as $a_{040} = a_{004}, a_{031} = a_{013}, a_{130} = a_{103}, a_{310} = a_{301}, a_{220} = a_{202}$ and $a_{121} = a_{112}$. We can then rewrite concisely the A_i 's and the B_j 's as follows.

$$\begin{aligned} A_i &= \sum_{\substack{0 \leq j, k \leq 4 \\ i+j+k=4}} a_{ijk} \quad \text{for } i = 0, 1, 2, 3, 4 \\ B_j &= \sum_{\substack{0 \leq i, k \leq 4 \\ i+j+k=4}} a_{ijk} \quad \text{for } j = 0, 1, 2, 3, 4 \end{aligned}$$

Let N be a large enough positive integer such each Na_{ijk} is an integer. We rise the construction $F_q \otimes F_q$ described in Section 3 to the N -th power. Observe that $(F'_q \otimes F'_q)^{\otimes N} \subseteq (F_q \otimes F_q)^{\otimes N}$ and

$$(F'_q \otimes F'_q)^{\otimes N} = \sum_{IJK} T_{IJK},$$

where the sum is over all triples of sequences IJK with $I, J, K \in \{0, 1, 2, 3, 4\}^N$ such that $I_\ell + J_\ell + K_\ell = 4$ for all $\ell \in \{1, \dots, N\}$. Here we use the notation $T_{IJK} = T_{I_1 J_1 K_1} \otimes \dots \otimes T_{I_N J_N K_N}$. Note that there are 15^N terms T_{IJK} in the above sum. In the tensor product the number of x -variables is $(q+2)^{2N}$. The number of y -variables and z -variables is also $(q+2)^{2N}$. Remember that in the original construction, each x -variable was indexed by a superscript in $\{0, 1, 2, 3, 4\}$. Each x -variable in the tensor product is thus indexed by a sequence of N such superscripts, i.e., by an element $I \in \{0, 1, 2, 3, 4\}^N$. The same is true for the y -variables and the z -variables. Note that the x -variables appearing in T_{IJK} have superscript I , the y -variables appearing in T_{IJK} have superscript J , and the z -variables appearing in T_{IJK} have superscript K .

Let us introduce the following definition.

Definition 5.1. Let $\bar{a}_{004}, \bar{a}_{040}, \bar{a}_{400}, \bar{a}_{013}, \bar{a}_{031}, \bar{a}_{103}, \bar{a}_{130}, \bar{a}_{301}, \bar{a}_{310}, \bar{a}_{022}, \bar{a}_{202}, \bar{a}_{220}, \bar{a}_{112}, \bar{a}_{121}, \bar{a}_{211}$ be fifteen nonnegative rational numbers. We say that a triple IJK is of type $[\bar{a}_{ijk}]$ if

$$|\{\ell \in \{1, \dots, N\} \mid I_\ell = i, J_\ell = j \text{ and } K_\ell = k\}| = \bar{a}_{ijk}N$$

for all 15 combinations of positive i, j, k with $i + j + k = 4$.

With a slight abuse of notation, we will say that a form T_{IJK} is of type $[\bar{a}_{ijk}]$ if the triple IJK is of type $[\bar{a}_{ijk}]$.

5.2 The first step

We set to zero all x -variables except those satisfying the following condition: their superscript I has exactly A_0N coordinates with value 0, A_1N coordinates with value 1, A_2N coordinates with value 2, A_3N coordinates with value 3 and A_4N coordinates with value 4. We will say that such a sequence I is of type A . There are

$$T_X = \binom{N}{A_0N, \dots, A_4N} = \Theta \left(\frac{1}{N^2} \left(\frac{1}{A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4}} \right)^N \right)$$

sequences I of type A (the approximation is done using Stirling's formula). After the zeroing operation, all forms T_{IJK} such that I is not of type A disappear (i.e., become zero).

We process the y -variables and the z -variables slightly differently. We set to zero all y -variables except those satisfying the following condition: their superscript J has exactly B_0N coordinates with value 0, B_1N coordinates with value 1, B_2N coordinates with value 2, B_3N coordinates with value 3 and B_4N coordinates with value 4. We will say that such a sequence is of type B . There are

$$T_Y = \binom{N}{B_0N, \dots, B_4N} = \Theta \left(\frac{1}{N^2} \left(\frac{1}{B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4}} \right)^N \right)$$

sequences J of type B . Similarly, we set to zero all z -variables except those such that their superscript K is of type B (there are T_Y such sequences).

After these three zeroing operations, the forms T_{IJK} remaining are precisely those such that I is of type A , J is of type B , and K is of type B . Equivalently, the forms remaining are precisely the forms T_{IJK} that are of type $[\bar{a}_{ijk}]$ with fifteen numbers \bar{a}_{ijk} (for all fifteen combinations of positive i, j, k such that $i + j + k = 4$) satisfying the following four conditions:

$$\bar{a}_{ijk}N \in \{0, 1, \dots, N\} \quad \text{for all } i, j, k; \quad (4)$$

$$A_i = \sum_{j,k: i+j+k=4} \bar{a}_{ijk} \quad \text{for } i = 0, 1, 2, 3, 4; \quad (5)$$

$$B_j = \sum_{i,k: i+j+k=4} \bar{a}_{ijk} \quad \text{for } j = 0, 1, 2, 3, 4; \quad (6)$$

$$B_k = \sum_{i,j: i+j+k=4} \bar{a}_{ijk} \quad \text{for } k = 0, 1, 2, 3, 4. \quad (7)$$

Let I be a fixed sequence of type A . The number of non-zeros forms T_{IJK} with this sequence I as its first index is thus precisely

$$\mathcal{N}_X = \sum_{[\bar{a}_{ijk}]} \binom{A_0N}{\bar{a}_{004}N, \bar{a}_{040}N, \bar{a}_{013}N, \bar{a}_{031}N, \bar{a}_{022}N} \binom{A_1N}{\bar{a}_{103}N, \bar{a}_{130}N, \bar{a}_{112}N, \bar{a}_{121}N} \times \\ \binom{A_2N}{\bar{a}_{202}N, \bar{a}_{220}N, \bar{a}_{211}N} \binom{A_3N}{\bar{a}_{301}N, \bar{a}_{310}N} \binom{A_4N}{\bar{a}_{400}N},$$

where the sum is over all the choices of fifteen parameters \bar{a}_{ijk} 's satisfying conditions (4)–(7).

For a fixed sequence J of type B , the number of non-zeros forms T_{IJK} with this sequence J as its second index is

$$\mathcal{N}_Y = \sum_{[\bar{a}_{ijk}]} \binom{B_0N}{\bar{a}_{004}N, \bar{a}_{400}N, \bar{a}_{103}N, \bar{a}_{301}N, \bar{a}_{202}N} \binom{B_1N}{\bar{a}_{013}, \bar{a}_{310}, \bar{a}_{112}, \bar{a}_{211}N} \times \\ \binom{B_2N}{\bar{a}_{022}N, \bar{a}_{220}N, \bar{a}_{121}N} \binom{B_3N}{\bar{a}_{031}N, \bar{a}_{130}N} \binom{B_4N}{\bar{a}_{040}N},$$

where the sum is again over all the choices of fifteen parameters \bar{a}_{ijk} 's satisfying conditions (5)–(7). Similarly, for a fixed sequence K of type B , the number of non-zeros forms T_{IJK} with this sequence K as its third index is

$$\mathcal{N}_Z = \sum_{[\bar{a}_{ijk}]} \binom{B_0 N}{\bar{a}_{040} N, \bar{a}_{400} N, \bar{a}_{130} N, \bar{a}_{310} N, \bar{a}_{220} N} \binom{B_1 N}{\bar{a}_{031}, \bar{a}_{301}, \bar{a}_{121}, \bar{a}_{211} N} \times \\ \binom{B_2 N}{\bar{a}_{022} N, \bar{a}_{202} N, \bar{a}_{112} N} \binom{B_3 N}{\bar{a}_{013} N, \bar{a}_{103} N} \binom{B_4 N}{\bar{a}_{004} N}.$$

The total number of remaining triples is

$$T_X \mathcal{N}_X = T_Y \mathcal{N}_Y = T_Z \mathcal{N}_Z. \quad (8)$$

Note that this implies that $\mathcal{N}_Y = \mathcal{N}_Z$.

We will also be interested in the number of remaining forms T_{IJK} of type $[a_{ijk}]$. For a fixed sequence I of type A , the number of non-zeros forms T_{IJK} of type $[a_{ijk}]$ with this sequence I as its first index is

$$\mathcal{N}_X^* = \binom{A_0 N}{a_{004} N, a_{004} N, a_{013} N, a_{013} N, a_{022} N} \binom{A_1 N}{a_{103} N, a_{103} N, a_{112} N, a_{112} N} \times \\ \binom{A_2 N}{a_{202} N, a_{202} N, a_{211} N} \binom{A_3 N}{a_{301} N, a_{301} N} \binom{A_4 N}{a_{400} N}.$$

For a fixed sequence J of type B , the number of non-zeros forms T_{IJK} of type $[a_{ijk}]$ with this sequence J as its second index is

$$\mathcal{N}_Y^* = \binom{B_0 N}{a_{004} N, a_{400} N, a_{103} N, a_{301} N, a_{202} N} \binom{B_1 N}{a_{013}, a_{301}, a_{112}, a_{211} N} \times \\ \binom{B_2 N}{a_{022} N, a_{202} N, a_{112} N} \binom{B_3 N}{a_{013} N, a_{103} N} \binom{B_4 N}{a_{004} N}.$$

We have

$$T_X \mathcal{N}_X^* = T_Y \mathcal{N}_Y^*. \quad (9)$$

5.3 Approximation

In this subsection we will use the notation $[\bar{a}_{ijk}]$ to represent an arbitrary set of fifteen parameters \bar{a}_{ijk} such that $0 \leq \bar{a}_{ijk} \leq 1$ for each i, j, k . Let $c([\bar{a}_{ijk}])$ denote the number of nonzero elements among these fifteen parameters. Consider the following expression:

$$g([\bar{a}_{ijk}]) = \left(\frac{\bar{a}_{004} \bar{a}_{040} \bar{a}_{400} \bar{a}_{013} \bar{a}_{031} \bar{a}_{103} \bar{a}_{130} \bar{a}_{301} \bar{a}_{310} \bar{a}_{022} \bar{a}_{202} \bar{a}_{220} \bar{a}_{112} \bar{a}_{121} \bar{a}_{211}}{a_{004} a_{040} a_{400} a_{013} a_{031} a_{103} a_{130} a_{301} a_{310} a_{022} a_{202} a_{220} a_{112} a_{121} a_{211}} \right)^{-1}.$$

Using Stirling's formula, we can give the following approximations.

$$\mathcal{N}_X = \Theta \left(\sum_{[\bar{a}_{ijk}]} \frac{[A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4} \times g([\bar{a}_{ijk}])]^N}{N^{(c([\bar{a}_{ijk}]) - 5)/2}} \right) \\ \mathcal{N}_Y = \Theta \left(\sum_{[\bar{a}_{ijk}]} \frac{[B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4} \times g([\bar{a}_{ijk}])]^N}{N^{(c([\bar{a}_{ijk}]) - 5)/2}} \right) \\ \mathcal{N}_X^* = \Theta \left(\frac{1}{N^5} [A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4} \times g([a_{ijk}])]^N \right) \\ \mathcal{N}_Y^* = \Theta \left(\frac{1}{N^5} [B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4} \times g([a_{ijk}])]^N \right)$$

The first two sums are again over all the choices of fifteen parameters \bar{a}_{ijk} 's satisfying conditions (4)–(7). Note that, for any $[\bar{a}_{ijk}]$ satisfying these four conditions, $c([\bar{a}_{ijk}]) \geq 5$, since the A_i 's are non-zero.

We know that $\mathcal{N}_X^* \leq \mathcal{N}_X$ and $\mathcal{N}_Y^* \leq \mathcal{N}_Y$, by definition. The following proposition shows that \mathcal{N}_X and \mathcal{N}_Y can actually be approximated by \mathcal{N}_X^* and \mathcal{N}_Y^* , respectively.

Proposition 5.1. $\mathcal{N}_X = O(N^8 \mathcal{N}_X^*)$ and $\mathcal{N}_Y = O(N^8 \mathcal{N}_Y^*)$.

Proof. Any set of values $[\bar{a}_{ijk}]$ that satisfies conditions (4)–(7) is such that $\bar{a}_{004} = a_{004}$, $\bar{a}_{040} = a_{040}$ and $\bar{a}_{400} = a_{400}$. Moreover, the other values a_{ijk} depend only on \bar{a}_{103} , \bar{a}_{031} and \bar{a}_{301} :

$$\begin{aligned}\bar{a}_{013} &= B_3 N - \bar{a}_{103} \\ \bar{a}_{130} &= B_3 N - \bar{a}_{031} \\ \bar{a}_{310} &= A_3 N - \bar{a}_{301} \\ \bar{a}_{022} &= (A_0 - 2B_4 - B_3)N + \bar{a}_{103} - \bar{a}_{031} \\ \bar{a}_{202} &= (-A_4 + B_0 - B_4)N - \bar{a}_{103} - \bar{a}_{301} \\ \bar{a}_{220} &= (-A_3 - A_4 + B_0 - B_3 - B_4)N + \bar{a}_{031} + \bar{a}_{301} \\ \bar{a}_{112} &= (-A_0 + A_4 - B_0 + B_2 + B_3 + 3B_4)N + \bar{a}_{031} + \bar{a}_{301} \\ \bar{a}_{121} &= (-A_0 + A_3 + A_4 - B_0 + B_2 + 2B_3 + 3B_4)N - \bar{a}_{103} - \bar{a}_{301} \\ \bar{a}_{211} &= (A_2 + A_3 + 2A_4 - 2B_0 + B_3 + 2B_4)N - \bar{a}_{031} + \bar{a}_{103}.\end{aligned}$$

Note that there are at most $(N + 1)$ choices for each \bar{a}_{103} , \bar{a}_{031} and \bar{a}_{301} , from condition (4). There are thus at most $(N + 1)^3$ choices for the values $[\bar{a}_{ijk}]$ satisfying conditions (4)–(7).

We now show that the expression $g([\bar{a}_{ijk}])$ is maximized for the values $[\bar{a}_{ijk}] = [a_{ijk}]$. Let us take the logarithm of the expression g . Since this is a concave function on a convex domain, a local optimum of $\log g$ is a global maximum of g . The partial derivatives of $\log g$ are as follows.

$$\begin{aligned}\frac{\partial \log g}{\partial \bar{a}_{103}} &= \log(\bar{a}_{013}) - \log(\bar{a}_{103}) - \log(\bar{a}_{022}) + \log(\bar{a}_{202}) + \log(\bar{a}_{121}) - \log(\bar{a}_{211}) \\ \frac{\partial \log g}{\partial \bar{a}_{031}} &= -\log(\bar{a}_{031}) + \log(\bar{a}_{130}) + \log(\bar{a}_{022}) - \log(\bar{a}_{220}) - \log(\bar{a}_{112}) + \log(\bar{a}_{211}) \\ \frac{\partial \log g}{\partial \bar{a}_{301}} &= -\log(\bar{a}_{301}) + \log(\bar{a}_{310}) + \log(\bar{a}_{202}) - \log(\bar{a}_{220}) - \log(\bar{a}_{112}) + \log(\bar{a}_{121})\end{aligned}$$

The values $[\bar{a}_{ijk}] = [a_{ijk}]$ satisfy $\frac{\partial \log g}{\partial \bar{a}_{103}} = \frac{\partial \log g}{\partial \bar{a}_{031}} = \frac{\partial \log g}{\partial \bar{a}_{301}} = 0$ since $a_{013}a_{202}a_{112} = a_{103}a_{022}a_{211}$.

We conclude that

$$\mathcal{N}_X = O\left((N + 1)^3 \left[A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4} \times g([a_{ijk}])\right]^N\right) = O(N^8 \mathcal{N}_X^*),$$

and similarly $\mathcal{N}_Y = O(N^8 \mathcal{N}_Y^*)$. □

5.4 The second step

We will now apply the results of Section 4, by associating to U the set of all forms T_{IJK} that are of type $[\bar{a}_{ijk}]$ for values \bar{a}_{ijk} satisfying conditions (4)–(7), and to U^* the set of all forms T_{IJK} of type $[a_{ijk}]$. Note that all the conditions of Subsection 4.1 are satisfied: we have $\tau = 4$, and values $T_1 = T_X$, $T_2 = T_3 = T_Y$, $\mathcal{N}_1 = \mathcal{N}_X$, $\mathcal{N}_2 = \mathcal{N}_3 = \mathcal{N}_Y$, $\mathcal{N}_1^* = \mathcal{N}_X^*$ and $\mathcal{N}_2^* = \mathcal{N}_3^* = \mathcal{N}_Y^*$.

With this association we can use the graph-theoretic framework developed in Section 4. The initial graph G , as defined in Subsection 4.1, corresponds to the current sum of trilinear forms (each vertex corresponds to a form T_{IJK} of type $[\bar{a}_{ijk}]$ for values \bar{a}_{ijk} satisfying conditions (4)–(7)). A removal operation on G corresponds to zeroing variables with a given superscript. For instance, removing all

vertices $u \in G$ such that $f_1(u) = I$ corresponds to zeroing all the x -variables with superscript I . Our goal is to zero variables in order to obtain a sum of several forms T_{IJK} satisfying the following two conditions. First, each form in the sum is of type $[a_{ijk}]$. Then, the forms in the sum do not share any index (i.e., if T_{IJK} and $T_{I'J'K'}$ are in the sum, then $I \neq I'$, $J \neq J'$ and $K \neq K'$), which implies that the same variable does not appear in more than one form, and thus that the sum is direct. This corresponds to constructing an edgeless subgraph of G in which all the vertices are in U^* , so we are in a situation where Theorem 4.2 can be applied.

Suppose that the inequality

$$A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4} \geq B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4}$$

holds. Then $T_X = O(T_Y)$, and Equalities (8) and (9) imply that $\mathcal{N}_Y = O(\mathcal{N}_X)$ and $\mathcal{N}_Y^* = O(\mathcal{N}_X^*)$. From Proposition 5.1 we then obtain the relation $\mathcal{N}_1 + \mathcal{N}_2 + \mathcal{N}_3 = O(N^8 \mathcal{N}_X^*)$. By the above discussion and Theorem 4.2, we can obtain a direct sum of

$$\Omega \left(\frac{T_X \mathcal{N}_X^*}{(N^8 \mathcal{N}_X^*)^{1+\epsilon}} \right)$$

forms, all of type $[a_{ijk}]$. By using the trivial upper bound $\mathcal{N}_X^* \leq 15^N$, we obtain the following theorem.

Theorem 5.1. *Let q be any positive integer and $a_{004}, a_{400}, a_{013}, a_{103}, a_{301}, a_{022}, a_{202}, a_{112}$ and a_{211} be any nine positive rational numbers satisfying the following three conditions:*

- $2a_{004} + a_{400} + 2a_{013} + 2a_{103} + 2a_{301} + a_{022} + 2a_{202} + 2a_{112} + a_{211} = 1$;
- $a_{013}a_{202}a_{112} = a_{103}a_{022}a_{211}$;
- $A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4} \geq B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4}$.

Then, for any constant $\epsilon > 0$, the trilinear form $(F_q \otimes F_q)^{\otimes N}$ can be converted (i.e., degenerated in the sense of Definition 2.1) into a direct sum of

$$\Omega \left(\frac{1}{N^{10+8\epsilon} 15^{N\epsilon}} \left[\frac{1}{A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4}} \right]^N \right)$$

forms, each form being isomorphic to

$$\bigotimes_{\substack{0 \leq i,j,k \leq 4 \\ i+j+k=4}} T_{ijk}^{\otimes a_{ijk}N}.$$

6 Upper Bounds on the Exponent of Rectangular Matrix Multiplication

Theorem 5.1 showed how the form $(F_q \otimes F_q)^{\otimes N}$ can be used to obtain a direct sum of many forms T_{IJK} such that

$$T_{IJK} \cong \bigotimes_{i,j,k:i+j+k=4} T_{ijk}^{\otimes a_{ijk}N}.$$

In order to apply Schönage's asymptotic sum inequality (Theorem 2.1), we need to analyze the smaller forms T_{ijk} . All the forms except T_{112} , T_{121} and T_{211} correspond to matrix multiplications, as described in Section 3. In Subsection 6.1 we analyze the forms T_{112} , T_{121} and T_{211} . Then, in Subsection 6.2, we put all our results together and prove our main result.

6.1 The forms T_{112} , T_{121} and T_{211}

Let us recall the definition of these three forms.

$$\begin{aligned}
T_{112} &= \sum_{i=1}^q x_{i,0}^1 y_{i,0}^1 z_{0,q+1}^2 + \sum_{k=1}^q x_{0,k}^1 y_{0,k}^1 z_{q+1,0}^2 + \sum_{i,k=1}^q x_{i,0}^1 y_{0,k}^1 z_{i,k}^2 + \sum_{i,k=1}^q x_{0,k}^1 y_{i,0}^1 z_{i,k}^2 \\
T_{121} &= \sum_{i=1}^q x_{i,0}^1 y_{0,q+1}^2 z_{i,0}^1 + \sum_{k=1}^q x_{0,k}^1 y_{q+1,0}^2 z_{0,k}^1 + \sum_{i,k=1}^q x_{i,0}^1 y_{i,k}^2 z_{0,k}^1 + \sum_{i,k=1}^q x_{0,k}^1 y_{i,k}^2 z_{i,0}^1 \\
T_{211} &= \sum_{i=1}^q x_{0,q+1}^2 y_{i,0}^1 z_{i,0}^1 + \sum_{k=1}^q x_{q+1,0}^2 y_{0,k}^1 z_{0,k}^1 + \sum_{i,k=1}^q x_{i,k}^2 y_{i,0}^1 z_{0,k}^1 + \sum_{i,k=1}^q x_{i,k}^2 y_{0,k}^1 z_{i,0}^1
\end{aligned}$$

We first focus on the form T_{211} . It will be convenient to write $T_{211} = t_{011} + t_{101} + t_{110} + t_{200}$, where

$$\begin{aligned}
t_{011} &= \sum_{i=1}^q x_{0,q+1}^2 y_{i,0}^1 z_{i,0}^1, \\
t_{101} &= \sum_{i,k=1}^q x_{i,k}^1 y_{0,k}^0 z_{i,0}^1, \\
t_{110} &= \sum_{i,k=1}^q x_{i,k}^1 y_{i,0}^1 z_{0,k}^0, \\
t_{200} &= \sum_{k=1}^q x_{q+1,0}^2 y_{0,k}^0 z_{0,k}^0.
\end{aligned}$$

Note that the superscripts in these forms differ from the original superscripts. They are nevertheless uniquely determined by the subscripts of the variables. Observe that $t_{011} \cong t_{200} \cong \langle 1, 1, q \rangle$, which corresponds to the product of a scalar by a row vector, and $t_{101} \cong t_{110} \cong \langle q, q, 1 \rangle$, which corresponds to the product of a $q \times q$ matrix by a column vector.

The following proposition states that tensor powers of T_{211} can be used to construct a direct sum of several trilinear forms, each one being a \mathcal{C} -tensor in which the support and all the components are isomorphic to a rectangular matrix product.

Proposition 6.1. *Let b be any constant such that $0.916027 < b \leq 1$. Then there exists a constant $c \geq 1$ depending only on b such that, for any $\epsilon > 0$ and any large enough integer m , the form $T_{211}^{\otimes 2m}$ can be converted into a direct sum of*

$$\Omega \left(\frac{1}{mc^{2\epsilon m}} \cdot \left[\frac{2}{(2b)^b (1-b)^{1-b}} \right]^{2m} \right)$$

trilinear forms, each form being a \mathcal{C} -tensor in which:

- each component is isomorphic to $\langle q^{2bm}, q^{2bm}, q^{2(1-b)m} \rangle$;
- the support is isomorphic to $\text{supp}_c(\langle 1, 1, H \rangle)$, where $H = \Omega \left(\frac{1}{\sqrt{m}} \cdot [(2b)^b (1-b)^{(1-b)}]^{2m} \right)$.

Remark. Proposition 6.1 uses the convention $0^0 = 1$. For the case $b = 1$, the proposition states that the form $T_{112}^{\otimes 2m}$ can be used to construct at least one tensor with support isomorphic to $\text{supp}_c(\langle 1, 1, H \rangle)$ for $H = \Omega(4^m / \sqrt{m})$, each component being isomorphic to $\langle q^{2m}, q^{2m}, 1 \rangle$.

Proof of Proposition 6.1. For simplicity we suppose that bm is an integer (otherwise, we can work with $\lfloor bm \rfloor$, which gives the same asymptotic complexity).

Let S be the set of all triples IJK with $I \in \{0, 1, 2\}^{2m}$ and $J, K \in \{0, 1\}^{2m}$ such that $I_\ell + J_\ell + K_\ell = 2$ for all $\ell \in \{1, \dots, 2m\}$. We raise the form T_{211} to the $2m$ -th power. This gives the form

$$\sum_{IJK \in S} t_{IJK}$$

where $t_{IJK} = t_{I_1 J_1 K_1} \otimes \cdots \otimes t_{I_{2m} J_{2m} K_{2m}}$. Each x -variable in the tensor product has for superscript a sequence in $\{0, 1, 2\}^{2m}$, and each y -variable or z -variable has for superscript a sequence in $\{0, 1\}^{2m}$. Let us decompose the space of x -variables as $\bigoplus_{I \in \{0, 1, 2\}^{2m}} X_I$, where X_I denotes the subspace of x -variables with superscript I . Similarly, decompose the space of y -variables as $\bigoplus_{J \in \{0, 1\}^{2m}} Y_J$ and the space of z -variables as $\bigoplus_{K \in \{0, 1\}^{2m}} Z_K$. The form $\sum t_{IJK}$ above is then a \mathcal{C} -tensor with respect to this decomposition, with support S . We will now modify this form (by zeroing variables, which will modify its support) in order to obtain a simple expression for each of its components.

We zero all the x -variables except those for which the superscript I has $(1-b)m$ coordinates with value 0, $(1-b)m$ coordinates with value 2 and $2bm$ coordinates with value 1. We zero all the y -variables and z -variables except those for which the superscript has m coordinates with value 0 and m coordinates with value 1. After these zeroing operations, the only forms remaining in the sum are those corresponding to indexes IJK satisfying the following four conditions.

$$\begin{aligned} |\{\ell \in \{1, \dots, 2m\} \mid I_\ell = 0, J_\ell = 1 \text{ and } K_\ell = 1\}| &= (1-b)m \\ |\{\ell \in \{1, \dots, 2m\} \mid I_\ell = 1, J_\ell = 0 \text{ and } K_\ell = 1\}| &= bm \\ |\{\ell \in \{1, \dots, 2m\} \mid I_\ell = 1, J_\ell = 1 \text{ and } K_\ell = 0\}| &= bm \\ |\{\ell \in \{1, \dots, 2m\} \mid I_\ell = 2, J_\ell = 0 \text{ and } K_\ell = 0\}| &= (1-b)m \end{aligned}$$

This means that each form t_{IJK} in this new sum (i.e., each component in the corresponding \mathcal{C} -tensor) is isomorphic to

$$t_{011}^{\otimes(1-b)m} \otimes t_{101}^{\otimes bm} \otimes t_{110}^{\otimes bm} \otimes t_{200}^{\otimes(1-b)m} \cong \langle q^{2bm}, q^{2bm}, q^{2(1-b)m} \rangle.$$

We now analyze the support of the new sum (the decomposition considered is unchanged).

The case $b = 1$ can be analyzed directly: there are $\binom{2m}{m} = \Theta(4^m / \sqrt{m})$ forms in the sum, all of them sharing the same first index (the all-one sequence $\mathbf{1} = 1 \cdots 1$). This sum is then $\sum_J t_{1JK}$, where for each t_{1JK} the sequence K is uniquely determined by J . The support of the sum is thus isomorphic to $\text{supp}_c(\langle 1, 1, \binom{2m}{m} \rangle)$.

To analyze the case $b < 1$, we will interpret the sum in the framework developed in Section 4, by letting U be the set of triples IJK satisfying the above four conditions. Indeed, all the requirements for U are satisfied: we have $\tau = 2$ and

$$\begin{aligned} T_1 &= \binom{2m}{(1-b)m, (1-b)m, 2mb} = \Theta\left(\frac{1}{m} \cdot \left[\frac{2}{(2b)^b(1-b)^{1-b}}\right]^{2m}\right) \\ \mathcal{N}_1 &= \binom{2mb}{mb} = \Theta\left(\frac{1}{\sqrt{m}} \cdot [2^b]^{2m}\right) \\ T_2 = T_3 &= \binom{2m}{m} = \Theta\left(\frac{1}{\sqrt{m}} \cdot [2]^{2m}\right) \\ \mathcal{N}_2 = \mathcal{N}_3 &= \binom{m}{m(1-b)} \binom{m}{m(1-b)} = \Theta\left(\frac{1}{m} \cdot \left[\frac{1}{b^b(1-b)^{1-b}}\right]^{2m}\right). \end{aligned}$$

Note in particular that $N_1 > N_2$, since $(2b)^b(1-b)^{1-b} > 1$ for any $b \geq 0.773$. Choose $U^* = U$ (which means that $\mathcal{N}_i = \mathcal{N}_i^*$ for each $i \in \{1, 2, 3\}$). The correspondence with the graph-theoretic interpretation of Section 4 is as follows. Each vertex in the graph G defined in Subsection 4.1 corresponds to one form t_{IJK} in the sum, which is isomorphic to $\langle q^{2bm}, q^{2bm}, q^{2(1-b)m} \rangle$ from the discussion above. One

removal operation corresponds to zeroing variables. A 1-clique of length n corresponds to a sum of n forms sharing their first index, which is precisely a \mathcal{C} -tensor with support isomorphic to $\text{supp}_c(\langle 1, 1, n \rangle)$.

For any value $b > 0.916027$ we have

$$\frac{2}{(2b)^{2b}(1-b)^{2(1-b)}} < 1$$

and thus $T_1\mathcal{N}_2/\mathcal{N}_1 = o(1)$. By Theorem 4.3, for any $\epsilon > 0$ we can then convert the sum into a direct sum of

$$\Omega\left(\frac{T_1}{\mathcal{N}_2^\epsilon}\right) = \Omega\left(\frac{1}{m^{1-\epsilon}} \cdot \left[\frac{2}{2^b (b^b(1-b)^{1-b})^{1-\epsilon}}\right]^{2m}\right)$$

\mathcal{C} -tensors, each tensor having support isomorphic to $\text{supp}_c(\langle 1, 1, n \rangle)$ for

$$n = \Omega\left(\frac{\mathcal{N}_1}{\mathcal{N}_2}\right) = \Omega\left(\sqrt{m} \cdot \left[(2b)^b(1-b)^{(1-b)}\right]^{2m}\right)$$

and components isomorphic to $\langle q^{2bm}, q^{2bm}, q^{2(1-b)m} \rangle$. Finally, note that

$$\frac{1}{m^{1-\epsilon}} \cdot \left[\frac{2}{2^b (b^b(1-b)^{1-b})^{1-\epsilon}}\right]^{2m} \geq \frac{1}{mc^{2\epsilon m}} \left[\frac{2}{(2b)^b(1-b)^{1-b}}\right]^{2m}$$

for some constant $c \geq 1$ depending only on b , since $b^b(1-b)^{1-b} < 1$. \square

The forms T_{112} and T_{121} can be analyzed in the same way as T_{211} by permuting the roles of the x -variables, the y -variables and the z -variables. Similarly to the statement of Proposition 6.1, the form $T_{112}^{\otimes 2m}$ gives a direct sum of \mathcal{C} -tensors with support isomorphic to $\langle 1, H, 1 \rangle$, each component in the tensors being isomorphic to $\langle q^{2bm}, q^{2(1-b)m}, q^{2bm} \rangle$. The form $T_{121}^{\otimes 2m}$ gives a direct sum of \mathcal{C} -tensors with support isomorphic to $\text{supp}_c(\langle H, 1, 1 \rangle)$, each component being isomorphic to $\langle q^{2(1-b)m}, q^{2bm}, q^{2bm} \rangle$.

Suppose that different constants are used to treat each of the three forms: the forms T_{112} and T_{121} are processed with some constant b , while T_{211} is processed with another constant \tilde{b} . For any fixed values a_{112}, a_{211} and any $\epsilon > 0$, the form $T_{112}^{\otimes a_{112}N} \otimes T_{112}^{\otimes a_{112}N} \otimes T_{211}^{\otimes a_{211}N}$ can then be used to construct a direct sum of

$$\Omega\left(\frac{1}{N^3 c'^{N\epsilon}} \cdot \left[\frac{2}{(2b)^b(1-b)^{1-b}}\right]^{2a_{112}N} \times \left[\frac{2}{(2\tilde{b})^{\tilde{b}}(1-\tilde{b})^{1-\tilde{b}}}\right]^{a_{211}N}\right)$$

\mathcal{C} -tensors, for some value $c' \geq 1$ depending only on b and \tilde{b} . Each of these \mathcal{C} -tensors has a support isomorphic to $\text{supp}_c(\langle H_{112}, H_{112}, H_{211} \rangle)$, where

$$\begin{aligned} H_{112} &= \Omega\left(\frac{1}{\sqrt{N}} \cdot \left[(2b)^b(1-b)^{(1-b)}\right]^{a_{112}N}\right) \\ H_{211} &= \Omega\left(\frac{1}{\sqrt{N}} \cdot \left[(2\tilde{b})^{\tilde{b}}(1-\tilde{b})^{(1-\tilde{b})}\right]^{a_{211}N}\right). \end{aligned}$$

In all these \mathcal{C} -tensors, each component is isomorphic to the rectangular matrix multiplication

$$\langle q^{(a_{112}+a_{211}\tilde{b})N}, q^{(a_{112}+a_{211}\tilde{b})N}, q^{(2a_{112}b+a_{211}(1-\tilde{b}))N} \rangle.$$

We can then use Propositions 2.2 and 2.3 to convert each \mathcal{C} -tensor into a direct sum of at least $\frac{3}{4}H_{112} \times \min(H_{112}, H_{211})$ trilinear forms, each isomorphic to $\langle q^{(a_{112}+a_{211}\tilde{b})N}, q^{(a_{112}+a_{211}\tilde{b})N}, q^{(2a_{112}b+a_{211}(1-\tilde{b}))N} \rangle$. We thus obtain the following result.

Proposition 6.2. *Let a_{112} and a_{211} be any two positive constants. Let b and \tilde{b} be any two constants such that $0.916027 < b, \tilde{b} \leq 1$. Then there exists a constant $c' \geq 1$ such that, for any $\epsilon > 0$, the form*

$$T_{112}^{\otimes a_{112}N} \otimes T_{121}^{\otimes a_{112}N} \otimes T_{211}^{\otimes a_{211}N}$$

can be converted into a direct sum of

$$\Omega \left(\frac{1}{N^4 c'^{N\epsilon}} \cdot \left[\frac{2^{2a_{112}+a_{211}}}{\max \left([(2b)^b(1-b)^{1-b}]^{a_{112}}, [(2\tilde{b})^{\tilde{b}}(1-\tilde{b})^{1-\tilde{b}}]^{a_{211}} \right)} \right]^N \right)$$

forms, each form being isomorphic to $\langle q^{(a_{112}+a_{211}\tilde{b})N}, q^{(a_{112}+a_{211}\tilde{b})N}, q^{(2a_{112}b+a_{211}(1-\tilde{b}))N} \rangle$.

6.2 Main theorem

Let us define the following three quantities.

$$\begin{aligned} Q &= (2q)^{a_{103}+a_{301}} \times (q^2+2)^{a_{202}} \times q^{a_{112}+a_{211}\tilde{b}} \\ R &= (2q)^{2a_{013}} \times (q^2+2)^{a_{022}} \times q^{2a_{112}b+(1-\tilde{b})a_{211}} \\ \mathcal{M} &= \frac{2^{2a_{112}+a_{211}}}{A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4}} \times \frac{1}{\max \left([(2b)^b(1-b)^{1-b}]^{a_{112}}, [(2\tilde{b})^{\tilde{b}}(1-\tilde{b})^{1-\tilde{b}}]^{a_{211}} \right)}. \end{aligned}$$

Our main theorem gives an upper bound on $\omega(1, 1, k)$ that depends on these quantities.

Theorem 6.1. *Let q be any positive integer and b, \tilde{b} be such that $0.916027 < b, \tilde{b} \leq 1$. Let $a_{004}, a_{400}, a_{013}, a_{103}, a_{301}, a_{022}, a_{202}, a_{112}$ and a_{211} be any nine positive rational numbers satisfying the following three conditions:*

- $2a_{004} + a_{400} + 2a_{013} + 2a_{103} + 2a_{301} + a_{022} + 2a_{202} + 2a_{112} + a_{211} = 1$;
- $a_{013}a_{202}a_{112} = a_{103}a_{022}a_{211}$;
- $A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4} \geq B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4}$.

Then

$$\mathcal{M}Q^{w(1,1,\frac{\log R}{\log Q})} \leq (q+2)^2.$$

Proof. Let $\epsilon > 0$ be an arbitrary positive value. Let N be a large integer and consider the trilinear form $(F_q \otimes F_q)^{\otimes N}$. Theorem 5.1 shows that this form can be used to obtain a direct sum of

$$r_1 = \Omega \left(\frac{1}{N^{10+8\epsilon} 15^{N\epsilon}} \left[\frac{1}{A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4}} \right]^N \right)$$

forms, each isomorphic to

$$\bigotimes_{i,j,k: i+j+k=4} T_{ijk}^{\otimes a_{ijk}N}.$$

All the terms T_{ijk} in this form, except T_{112}, T_{121} and T_{211} , correspond to matrix multiplications and have been analyzed in Section 3. By Proposition 6.2 the part $T_{112}^{\otimes a_{112}N} \otimes T_{121}^{\otimes a_{112}N} \otimes T_{211}^{\otimes a_{211}N}$ can be used to obtain a direct sum of

$$r_2 = \Omega \left(\frac{1}{N^4 c'^{N\epsilon}} \cdot \left[\frac{2^{2a_{112}+a_{211}}}{\max \left([(2b)^b(1-b)^{1-b}]^{a_{112}}, [(2\tilde{b})^{\tilde{b}}(1-\tilde{b})^{1-\tilde{b}}]^{a_{211}} \right)} \right]^N \right)$$

matrix multiplications $\langle q^{(a_{112}+a_{211}\tilde{b})N}, q^{(a_{112}+a_{211}\tilde{b})N}, q^{(2a_{112}b+(1-\tilde{b})a_{211})N} \rangle$.

This means that the trilinear form $(F_q \otimes F_q)^{\otimes N}$ can be converted into a direct sum of $r_1 r_2$ matrix multiplications $\langle Q^N, Q^N, R^N \rangle$. In other words:

$$r_1 r_2 \cdot \langle Q^N, Q^N, R^N \rangle \leq (F_q \otimes F_q)^{\otimes N}.$$

Since $\underline{R}(F_q \otimes F_q) \leq (q+2)^2$, as mentioned in Section 3, we know that $\underline{R}((F_q \otimes F_q)^{\otimes N}) \leq (q+2)^{2N}$. By Schönhage's asymptotic sum inequality (Theorem 2.1) we then conclude that

$$r_1 r_2 \times Q^{N\omega(1,1,\frac{\log R}{\log Q})} \leq (q+2)^{2N}.$$

Taking the N -th root, we obtain:

$$\frac{1}{(15c')^{\epsilon N^{(14+8\epsilon)/N}}} \times \mathcal{M}Q^{\omega(1,1,\frac{\log R}{\log Q})} \leq (q+2)^2.$$

For any $\epsilon > 0$ the above inequality holds for large enough integers N . By letting N grow to infinity, and then letting ϵ decrease to zero, we conclude that $\mathcal{M}Q^{\omega(1,1,\frac{\log R}{\log Q})} \leq (q+2)^2$. \square

7 Optimization

In this section we use Theorem 6.1 to derive numerical upper bounds on the exponent of rectangular matrix multiplication, and prove Theorem 1.1.

7.1 Square matrix multiplication

In this subsection we briefly show that our results give, for the exponent of square matrix multiplication, the same upper bound as the bound obtained by Coppersmith and Winograd [10].

Due to the symmetry of square matrix multiplication, we take $b = \tilde{b}$, $a_{400} = a_{004}$, $a_{103} = a_{013} = a_{301}$, $a_{022} = a_{202}$ and $a_{112} = a_{211}$. Then only six parameters remain, and the conditions $a_{013}a_{202}a_{112} = a_{103}a_{022}a_{211}$ and $A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4} = B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4}$ are immediately satisfied.

Theorem 6.1 shows that

$$\frac{8^{a_{112}} \times [(2q)^{2a_{013}} \times (q^2 + 2)^{a_{202}} \times q^{(1+b)a_{112}}]^{\omega(1,1,1)}}{A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4} \times [(2b)^b (1-b)^{(1-b)}]^{a_{112}}} \leq (q+2)^2.$$

By choosing $q = 6$, $b = 0.9724317$, $a_{103} = 0.012506$, $a_{202} = 0.102546$, $a_{112} = 0.205542$ and $a_{004} = 0.0007/3$, we obtain the upper bound $\omega(1,1,1) < 2.375477$.

This upper bound on the exponent of square matrix multiplication is exactly the same value as in [10]. This is not a coincidence. Indeed, by setting $b = \frac{q^{\omega(1,1,1)}}{q^{\omega(1,1,1)}+2}$, which is larger than 0.916027 for $q \geq 5$, we obtain

$$\frac{q^{b\omega(1,1,1)}}{(2b)^b (1-b)^{(1-b)}} = \frac{q^{\omega(1,1,1)} + 2}{2}$$

and our inequality becomes

$$\frac{(2q)^{2a_{013}\omega(1,1,1)} \times (q^2 + 2)^{a_{202}\omega(1,1,1)} \times (4q^{\omega(1,1,1)}(q^{\omega(1,1,1)} + 2))^{a_{112}}}{A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4}} \leq (q+2)^2,$$

which is exactly the same optimization problem as in Section 8 of [10].

q	5	5	6
b	0.984599222	0.968978515	0.94866036
\tilde{b}	0.919886704	0.938616630	0.99996514
a_{400}	0.004942000	0.001498500	0.00000090
a_{103}	0.010965995	0.014456894	0.01553556
a_{301}	0.055710210	0.031215255	0.00079349
a_{022}	0.037622078	0.065869083	0.22704392
a_{202}	0.138698196	0.118190058	0.05836108
a_{112}	0.145715589	0.178553843	0.20388121
a_{211}	0.245013049	0.226835534	0.13394891
a_{004}	0.00011...	0.000246...	0.001224...
a_{013}	0.00500...	0.010235...	0.039707...
$A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4}$	0.326588...	0.3265988...	0.339123647...
$B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4}$	0.326587...	0.3265987...	0.339123642...
$\log R / \log Q$	0.530200005...	0.750000001...	2.00000004...
$(2 \log(q+2) - \log \mathcal{M}) / \log Q$	2.060395...	2.190086...	3.256688...

Table 2: Three solutions for our optimization problem. The first ten rows give (exact) values of the ten parameters. The numeral values of the next four rows show that the three conditions $0 < a_{004}, a_{013} \leq 1$ and $A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4} \geq B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4}$ are satisfied. The numerical values of the last two rows show that $\omega(1, 1, 0.5302) < 2.060396$, $\omega(1, 1, 0.75) < 2.190087$ and $\omega(1, 1, 2) < 3.256689$.

7.2 Rectangular matrix multiplication

In this subsection we explain how to use Theorem 6.1 to derive an upper bound on $\omega(1, 1, k)$ for an arbitrary value k , and show how to obtain the results stated in Table 1 and Figure 1.

We use the following strategy. We take a positive integer q , seven positive rational numbers a_{400} , a_{103} , a_{301} , a_{022} , a_{202} , a_{112} and a_{211} , and two values b, \tilde{b} such that $0.916027 < b, \tilde{b} \leq 1$. We then fix

$$a_{013} = \frac{a_{103} a_{022} a_{211}}{a_{202} a_{112}}$$

and

$$a_{004} = \frac{1 - (a_{400} + 2a_{013} + 2a_{103} + 2a_{301} + a_{022} + 2a_{202} + 2a_{112} + a_{211})}{2}.$$

The conditions that have to be satisfied are:

- $0 < a_{004}, a_{013} \leq 1$;
- $A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4} \geq B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4}$.

If these conditions are satisfied, by Theorem 6.1 this gives the upper bound

$$\omega\left(1, 1, \frac{\log R}{\log Q}\right) \leq \frac{2 \log(q+2) - \log \mathcal{M}}{\log Q}.$$

The above discussion reduces the problem of finding an upper bound on $\omega(1, 1, k)$ to solving a nonlinear optimization problem. The upper bounds presented in Table 1 are obtained precisely by solving this optimization problem using Maple. We show exact values of the parameters proving that $\omega(1, 1, 0.5302) < 2.060396$, $\omega(1, 1, 0.75) < 2.190087$ and $\omega(1, 1, 2) < 3.256689$ in Table 2.

7.3 The value α

In this subsection we describe how to use Theorem 6.1 to obtain a lower bound on the value α , the largest value such that the product of an $n \times n^\alpha$ matrix by an $n^\alpha \times n$ matrix can be computed with $O(n^{2+\epsilon})$ arithmetic operations for any $\epsilon > 0$. The analysis is more delicate than in the previous subsection, since we will need to exhibit parameters such that $\mathcal{M}Q^2 = (q+2)^2$, with an equality rather than an inequality, and is done by finding analytically the optimal values of all but a few parameters.

Let q be an integer such that $q \geq 5$. For convenience, we will write $\kappa = 1/(q+2)^2$. Let a_{112} and a_{211} be any rational numbers such that $0 < a_{112} < q\kappa$ and $0 < a_{211} < (q^2+2)\kappa$. We set the parameters $b, \tilde{b}, a_{004}, a_{103}, a_{202}$ and a_{301} as follows:

$$\begin{aligned} b &= 1 \\ \tilde{b} &= q^2/(q^2+2) \\ a_{400} &= \kappa \\ a_{103} &= q\kappa - a_{112} \\ a_{202} &= ((q^2+2)\kappa - a_{211})/2 \\ a_{301} &= q\kappa. \end{aligned}$$

Putting these values in the formula for Q , we obtain:

$$\begin{aligned} Q &= (2q)^{q\kappa+a_{103}} \times (q^2+2)^{\frac{(q^2+2)\kappa-a_{211}}{2}} \times q^{a_{112}+q^2a_{211}/(q^2+2)} \\ &= (2q)^{2q\kappa} \times (q^2+2)^{\frac{(q^2+2)\kappa}{2}} \times 2^{-a_{112}} \times \left(\frac{q^{q^2/(q^2+2)}}{\sqrt{q^2+2}} \right)^{a_{211}}. \end{aligned}$$

Observe that $A_1 = A_3 = 2q\kappa$, $A_2 = (q^2+2)\kappa$, $A_4 = \kappa$ and $A_0 = 1 - (A_1 + A_2 + A_3 + A_4) = \kappa$. Then we obtain the following equality.

$$\frac{1}{A_0^{A_0} A_1^{A_1} A_2^{A_2} A_3^{A_3} A_4^{A_4}} = \frac{(q+2)^2}{(2q)^{4q\kappa} (q^2+2)^{(q^2+2)\kappa}}$$

The following lemma shows that, when a_{112} is small enough, the condition $\mathcal{M}Q^2 = (q+2)^2$ is satisfied.

Lemma 7.1. *Suppose that*

$$a_{112} \leq \left(1 + \frac{2q^2}{q^2+2} \log_2(q) - \log_2(q^2+2) \right) a_{211}. \quad (10)$$

Then $\mathcal{M}Q^2 = (q+2)^2$.

Proof. Our choice for b and \tilde{b} gives

$$\begin{aligned} [(2b)^b(1-b)^{1-b}]^{a_{112}} &= 2^{a_{112}} \\ [(2\tilde{b})^{\tilde{b}}(1-\tilde{b})^{1-\tilde{b}}]^{a_{211}} &= \left[\frac{2}{q^2+2} \cdot q^{\frac{2q^2}{q^2+2}} \right]^{a_{211}}. \end{aligned}$$

Inequality (10) then implies that $[(2b)^b(1-b)^{1-b}]^{a_{112}} \leq [(2\tilde{b})^{\tilde{b}}(1-\tilde{b})^{1-\tilde{b}}]^{a_{211}}$. In consequence,

$$\mathcal{M} = \frac{(q+2)^2}{(2q)^{4q\kappa} (q^2+2)^{(q^2+2)\kappa}} \times 4^{a_{112}} \times \left[\frac{q^2+2}{q^{2q^2/(q^2+2)}} \right]^{a_{211}},$$

which gives $\mathcal{M}Q^2 = (q+2)^2$. □

We now explain how to determine the three remaining parameters a_{004} , a_{013} and a_{022} . Remember that the parameters should satisfy the equalities

$$a_{013} = \frac{a_{103}a_{211}}{a_{202}a_{112}}a_{022}$$

and

$$2a_{004} + a_{400} + 2a_{013} + 2a_{103} + 2a_{301} + a_{022} + 2a_{202} + 2a_{112} + a_{211} = 1.$$

From our choice of parameters, the second equality can be rewritten as $2a_{004} + 2a_{013} + a_{022} = \kappa$. Since the parameter a_{004} should be positive, we obtain the condition

$$\left(\frac{4(q\kappa - a_{112})a_{211}}{((q^2 + 2)\kappa - a_{211})a_{112}} + 1 \right) a_{022} < \kappa. \quad (11)$$

If a_{022} , a_{112} and a_{211} satisfy this inequality, then the parameter a_{004} is fixed:

$$a_{004} = \left(\kappa - \left(\frac{4(q\kappa - a_{112})a_{211}}{((q^2 + 2)\kappa - a_{211})a_{112}} + 1 \right) a_{022} \right) / 2.$$

Note that Inequality (11) forces the value a_{013} to be at most 1.

All the values are thus determined by the choice of q , a_{022} , a_{112} and a_{211} . In particular, we obtain

$$R = (2q)^{\frac{4(q\kappa - a_{112})a_{211}}{((q^2 + 2)\kappa - a_{211})a_{112}}} \times (q^2 + 2)^{a_{022}} \times q^{2a_{112} + \frac{2}{q^2 + 2}a_{211}}.$$

We can similarly express the values of B_0 , B_1 , B_2 , B_3 and B_4 in function of these four parameters. We then want to solve the following optimization problem.

Maximize $\frac{\log R}{\log Q}$ subject to

- $0 \leq a_{022} \leq 1$, $0 < a_{112} \leq 5\kappa$ and $0 \leq a_{211} \leq (q^2 + 2)\kappa$;
- q is an integer such that $q \geq 5$;
- Inequalities (10) and (11) hold;
- $\frac{(2q)^{4q\kappa} (q^2 + 2)^{(q^2 + 2)\kappa}}{(q + 2)^2} \geq B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4}$.

By taking the values $q = 5$, $a_{022} = 0.0174853$, $a_{112} = 0.0945442$ and $a_{211} = 0.1773724$, we obtain the value $\alpha \geq \frac{\log R}{\log Q} > 0.30298$. These parameters satisfy all the constraints. We obtain in particular the following numerical values.

$$\begin{aligned} \frac{(2q)^{4q\kappa} (q^2 + 2)^{(q^2 + 2)\kappa}}{(q + 2)^2} &= 0.3211277\dots \\ B_0^{B_0} B_1^{B_1} B_2^{B_2} B_3^{B_3} B_4^{B_4} &= 0.3211276\dots \\ R &= 1.475744\dots \\ Q &= 3.612672\dots \end{aligned}$$

A more precise lower bound on α can be found using optimization software and high precision arithmetic. Using Maple and truncating the result of the optimization after the 25th digit, we find that for $q = 5$ the values

$$\begin{aligned} a_{022} &= 0.0174853267797595451457284 \\ a_{112} &= 0.0945442542111395375830367 \\ a_{211} &= 0.1773724081899825630904504 \end{aligned}$$

give the lower bound

$$\alpha > 0.3029805825293869820274449.$$

Acknowledgments

The author is grateful to Virginia Vassilevska Williams and Ryan Williams for helpful correspondence about Andrew Stothers' work, and to Virginia Vassilevska Williams for suggesting that the next step is to use higher tensor powers of the basic construction to improve rectangular matrix multiplication. He also acknowledges support from the JSPS and the MEXT, under the grant-in-aids Nos. 22800006 and 24700005.

References

- [1] ALON, N., SHPILKA, A., AND UMANS, C. On sunflowers and matrix multiplication. In *Proceedings of the 27th Conference on Computational Complexity* (to appear, 2012).
- [2] ALON, N., AND YUSTER, R. Fast algorithms for maximum subset matching and all-pairs shortest paths in graphs with a (not so) small vertex cover. In *Proceedings of the 15th Annual European Symposium on Algorithms* (2007), pp. 175–186.
- [3] AMOSSEN, R. R., AND PAGH, R. Faster join-projects and sparse matrix multiplications. In *Proceedings of the 12th International Conference on Database Theory* (2009), pp. 121–126.
- [4] BEHREND, F. A. On sets of integers which contain no three terms in arithmetical progression. *Proceedings of the National Academy of Sciences of the United States of America* 32, 331–332 (1946).
- [5] BÜRGISSER, P., CLAUSEN, M., AND SHOKROLLAHI, M. A. *Algebraic complexity theory*. Springer, 1997.
- [6] COHN, H., KLEINBERG, R. D., SZEGEDY, B., AND UMANS, C. Group-theoretic algorithms for matrix multiplication. In *Proceedings of the 46th Annual IEEE Symposium on Foundations of Computer Science* (2005), pp. 379–388.
- [7] COHN, H., AND UMANS, C. A group-theoretic approach to fast matrix multiplication. In *Proceedings of the 44th Symposium on Foundations of Computer Science* (2003), pp. 438–449.
- [8] COPPERSMITH, D. Rapid multiplication of rectangular matrices. *SIAM Journal on Computing* 11, 3 (1982), 467–471.
- [9] COPPERSMITH, D. Rectangular matrix multiplication revisited. *Journal of Complexity* 13, 1 (1997), 42–49.
- [10] COPPERSMITH, D., AND WINOGRAD, S. Matrix multiplication via arithmetic progressions. *Journal of Symbolic Computation* 9, 3 (1990), 251–280.
- [11] CZUMAJ, A., KOWALUK, M., AND LINGAS, A. Faster algorithms for finding lowest common ancestors in directed acyclic graphs. *Theoretical Computer Science* 380, 1-2 (2007), 37–46.
- [12] DEMETRESCU, C., AND ITALIANO, G. F. Fully dynamic transitive closure: Breaking through the $o(n^2)$ barrier. In *Proceedings of the 41st Annual Symposium on Foundations of Computer Science* (2000), pp. 381–389.
- [13] HUANG, X., AND PAN, V. Y. Fast rectangular matrix multiplication and applications. *Journal of Complexity* 14, 2 (1998), 257–299.
- [14] KAPLAN, H., RUBIN, N., SHARIR, M., AND VERBIN, E. Counting colors in boxes. In *Proceedings of the 18th Annual ACM-SIAM Symposium on Discrete Algorithms* (2007), pp. 785–794.

- [15] KAPLAN, H., SHARIR, M., AND VERBIN, E. Colored intersection searching via sparse rectangular matrix multiplication. In *Proceedings of the 22nd ACM Symposium on Computational Geometry* (2006), pp. 52–60.
- [16] KE, S., ZENG, B., HAN, W., AND PAN, V. Y. Fast rectangular matrix multiplication and some applications. *Science in China Series A: Mathematics* 51, 3 (2008), 389–406.
- [17] LOTTI, G., AND ROMANI, F. On the asymptotic complexity of rectangular matrix multiplication. *Theoretical Computer Science* 23 (1983), 171–185.
- [18] PATRASCU, M., AND WILLIAMS, R. On the possibility of faster SAT algorithms. In *Proceedings of the 21st Annual ACM-SIAM Symposium on Discrete Algorithms* (2010), pp. 1065–1075.
- [19] RODITTY, L., AND SHAPIRA, A. All-pairs shortest paths with a sublinear additive error. *ACM Transactions on Algorithms* 7, 4 (2011), 45.
- [20] SALEM, R., AND SPENCER, D. C. On sets of integers which contain no three terms in arithmetical progression. *Proceedings of the National Academy of Sciences of the United States of America* 28 (1942), 561–563.
- [21] SANKOWSKI, P. Dynamic transitive closure via dynamic matrix inverse (extended abstract). In *Proceedings of the 45th Symposium on Foundations of Computer Science* (2004), pp. 509–517.
- [22] SANKOWSKI, P., AND MUCHA, M. Fast dynamic transitive closure with lookahead. *Algorithmica* 56, 2 (2010), 180–197.
- [23] SCHÖNHAGE, A. Partial and total matrix multiplication. *SIAM Journal on Computing* 10, 3 (1981), 434–455.
- [24] STOTHERS, A. *On the Complexity of Matrix Multiplication*. PhD thesis, University of Edinburgh, 2010.
- [25] STRASSEN, V. The asymptotic spectrum of tensors and the exponent of matrix multiplication. In *Proceedings of the 27th Annual Symposium on Foundations of Computer Science* (1986), pp. 49–54.
- [26] STRASSEN, V. Relative bilinear complexity and matrix multiplication. *Journal für die reine und angewandte Mathematik* 375-376 (1987), 406–443.
- [27] VASSILEVSKA WILLIAMS, V. Multiplying matrices faster than Coppersmith-Winograd. In *Proceedings of the 44th ACM Symposium on Theory of Computing* (to appear, 2012).
- [28] WILLIAMS, R. Non-uniform ACC circuit lower bounds. In *Proceedings of the 26th Annual IEEE Conference on Computational Complexity* (2011), pp. 115–125.
- [29] YUSTER, R. Efficient algorithms on sets of permutations, dominance, and real-weighted APSP. In *Proceedings of the 20th Annual ACM-SIAM Symposium on Discrete Algorithms* (2009), pp. 950–957.
- [30] YUSTER, R. Computing the diameter polynomially faster than APSP. *CoRR abs/1011.6181* (2010).
- [31] YUSTER, R., AND ZWICK, U. Detecting short directed cycles using rectangular matrix multiplication and dynamic programming. In *Proceedings of the 15th Annual ACM-SIAM Symposium on Discrete Algorithms* (2004), pp. 254–260.
- [32] YUSTER, R., AND ZWICK, U. Fast sparse matrix multiplication. *ACM Transactions on Algorithms* 1, 1 (2005), 2–13.

- [33] ZWICK, U. All pairs lightest shortest paths. In *Proceedings of the 31st Annual ACM Symposium on Theory of Computing* (1999), pp. 61–69.
- [34] ZWICK, U. All pairs shortest paths using bridging sets and rectangular matrix multiplication. *Journal of the ACM* 49, 3 (2002), 289–317.