Advanced Algorithms 2020 10/02, 2020

Exercise 3

Lecturer: Mohsen Ghaffari *Teaching Assistang: Julian Portmann*

1 Warm-up [Recommended]

It will be useful to recall the Chernoff bound that says that for independent Bernoulli distributed random variables $X_1, \ldots, X_n$ with $\mathbf{E}[X_i] = p$ we have

$$\Pr\left[|\sum_{1}^{n} X_i - pn| \geq \varepsilon pn\right] \leq 2\mathrm{e}^{-\frac{\varepsilon^2 pn}{3}}$$

for $0 < \varepsilon < 1$.

Target shooting: Suppose we have a ground set S and we wish to estimate the size of a subset $T \subseteq S$. We can do it by repeatedly sampling from S: we sample m times a uniform element from S and let X_i to be an indicator for the event of the i-th sample being an element of T. One can use the Chernoff bound to prove that the variable $(X_1 + X_2 + \ldots X_m)/m$ is within $(1+\varepsilon)$ multiplicative error of $|T|/|S|$ with probability at least $1-\delta$ if we set

$$m = \Theta\left(\frac{|S|}{|T|}\varepsilon^{-2}\log(1/\delta)\right).$$

1. Argue that the lemma really follows from the Chernoff bound.
2. Argue that if we sample only $O(\frac{|S|}{|T|})$ elements, with constant probability we never sample an element from T throughout the whole procedure and, hence, we cannot get a reasonable estimate.
3. Suppose that we have a FPRAS that returns value that is in the correct range $[(1-\varepsilon)OPT, (1+\varepsilon)OPT]$ with probability 2/3. Recall the median trick from the lecture and show how to amplify the probability to $1-\delta$ with $O(\log 1/\delta)$ calls of the original algorithm.

2 Counting satisfying assignments

In the lecture we have seen a FPRAS that estimates the number of satisfying assignments of a given DNF formula. Suppose that we wish to solve this problem and we know that one of the clauses contains only 10 variables. Can you propose a faster algorithm than the one you know from the lecture?

3 Counting independent sets [Recommended]

By an independent set of a graph G we mean a subset $S \subseteq V(G)$ of its vertices such that there is no edge with both endpoints in S. In the lecture we have seen an algorithm that estimates the number of colorings of a given graph if there is a way to sample them. In this exercise we wish to do the same for the problem of counting the number of independent sets. This means following: You are given a graph G and want to design a FPRAS for the number of independent sets in G. You are given a procedure that takes some graph as an input and outputs a uniformly random independent set from this graph as an answer.