



# Glyphosate pathways to modern diseases VI: Prions, amyloidoses and autoimmune neurological diseases

Anthony Samsel<sup>1</sup> and Stephanie Seneff<sup>2,\*</sup>

<sup>1</sup> Samsel Environmental and Public Health Services, Deerfield, NH 03037, USA

<sup>2</sup> Computer Science and Artificial Intelligence Laboratory, MIT, Cambridge, MA 02139, USA

Usage of the herbicide glyphosate on core crops in the USA has increased exponentially over the past two decades, in step with the exponential increase in autoimmune diseases including autism, multiple sclerosis, inflammatory bowel disease, type 1 diabetes, coeliac disease, neuromyelitis optica and many others. In this paper we explain how glyphosate, acting as a non-coding amino acid analogue of glycine, could erroneously be integrated with or incorporated into protein synthesis in place of glycine, producing a defective product that resists proteolysis. Whether produced by a microbe or present in a food source, such a peptide could lead to autoimmune disease through molecular mimicry. We discuss similarities in other naturally produced disease-causing amino acid analogues, such as the herbicide glufosinate and the insecticide L-canavanine, and provide multiple examples of glycine-containing short peptides linked to autoimmune disease, particularly with respect to multiple sclerosis. Most disturbing is the presence of glyphosate in many popular vaccines including the measles, mumps and rubella (MMR) vaccine, which we have verified here for the first time. Contamination may come through bovine protein, bovine calf serum, bovine casein, egg protein and/or gelatin. Gelatin sourced from the skin and bones of pigs and cattle given glyphosate-contaminated feed contains the herbicide. Collagen, the principal component of gelatin, contains very high levels of glycine, as do the digestive enzymes: pepsin, trypsin and lipase. The live measles virus could produce glyphosate-containing haemagglutinin, which might induce an autoimmune attack on myelin basic protein, commonly observed in autism. Regulatory agencies urgently need to reconsider the risks associated with the indiscriminate use of glyphosate to control weeds.

**Keywords:** autism, autoimmune disease, collagen, glycine, glyphosate, multiple sclerosis, protein misfolding, vaccines

## 1. INTRODUCTION

At first glance, multiple sclerosis (MS) and autism appear to have little in common, aside from the fact that both are neurological diseases. Autism is a condition with prenatal or early childhood onset, characterized by repetitive behaviours, impaired social interaction and cognitive impairment. The male:female ratio for autism is 4:1, while multiple sclerosis is twice as common in women as in men; its first symptoms usually begin in early adulthood to involve impaired lower limb mobility, although in later stages it affects both mental and physical capabilities. Both conditions are, however, associated with inflammatory autoimmune features [1, 2], and both diseases are viewed as having an environmental and a genetic component [3–6].

A study comparing a population of 658 MS patients with the general population found an association between MS and increased rates of asthma, inflammatory bowel disease (IBD), type 1 diabetes mellitus, pernicious anaemia and autoimmune thyroid disease [7], all of which

have also been linked to autism [8–11]. These conditions are all considered to be *autoimmune diseases*, which can be triggered through molecular mimicry, where an antibody responding to a foreign protein that resembles a native protein becomes sensitized to the native protein as well [12]. A paper by Shoenfeld and Aron-Maor in 2000 developed the argument that both autism and MS may be examples of an autoimmune reaction via mimicry following exposure to an antigenic stimulus, possibly from an infection or through vaccination [13]. They further propose specifically that myelin basic protein (MBP) and other proteins constituting the myelin sheath are attacked by the immune system in both autism and MS. This has been recognized by many others in autism [14, 15] and MS [16–20]. In 1982, Weizman et al. reported a cell-mediated autoimmune response to human MBP in 76% of the autistic children studied [16]. Immune sensitization to the myelin sheath proteins could arise either through mimicry as a consequence of exposure of the immune system to a foreign antigen with a similar peptide sequence that is

\* Corresponding author. E-mail: seneff@csail.mit.edu

resistant to clearance, or because the proteins themselves have been altered in some way that renders them defective, exposed and/or resistant to proteolysis.

Unlike DNA synthesis, protein synthesis is highly prone to error [21, 22]. It appears that biological systems have adopted a strategy of allowing coding errors to survive during active synthesis, but use protein misfolding as a criterion to mark a defective peptide for degradation and recycling through ubiquitination. It is estimated that 15% of average-length proteins will have at least one misincorporated amino acid. Typically, 10–15% of random substitutions disrupt protein function, mostly because of misfolding [22]. Such destabilization causes protein–protein aggregation, and can lead to multiple neurological diseases and amyloidoses. Drummond et al. propose that early-forming toxic oligomers of amyloidogenic proteins are enriched with missense errors [22].

Glyphosate is the active ingredient in the pervasive herbicide Roundup and in many other formulations of herbicides used to control weeds on agricultural, residential and public land worldwide. A recent study based in Germany involving 399 urine samples from adults not involved in agricultural work revealed glyphosate residues above the detection limit in the urine of 32% of the subjects, and residues of AMPA, a metabolite, in 40% [23]. In a paper published in 2014, Swanson et al. showed a remarkable correlation between the rising rate of glyphosate usage on corn (maize) and soy crops in the USA and an alarming rise in a number of different chronic diseases [24]. Additional strong correlations for other conditions and diseases are provided in two follow-on papers [25, 26]. While correlation does not necessarily mean causation, causation becomes much more likely if a plausible mechanism can be found. Swanson et al. found a remarkable 0.98 correlation coefficient between the rise in autism rates in the USA and the use of glyphosate on crops ( $P$ -value  $\leq 9.6 \times 10^{-6}$ ). The correlation for multiple sclerosis was not as high, but still highly significant at 0.83 ( $P$ -value  $\leq 1.1 \times 10^{-5}$ ). IBD had a correlation coefficient of 0.94 ( $P$ -value  $\leq 7.1 \times 10^{-8}$ ) (see Table 1 for other diseases).

Table 1. Correlations between time trends in several diseases and conditions recorded by the US Centers for Disease Control (CDC) with glyphosate usage on corn (maize) and soy crops reported by the USDA. Data reproduced from [23] and [25].

Disease	Correlation coefficient (R)	$P$ -value
Autism (prevalence)	0.98	$9.6 \times 10^{-6}$
MS (deaths)	0.83	$1.1 \times 10^{-5}$
IBD	0.94	$7.1 \times 10^{-8}$
Anaemia	0.90	$1.8 \times 10^{-4}$
Diabetes (prevalence)	0.97	$9.2 \times 10^{-9}$
Thyroid cancer (incidence)	0.99	$7.6 \times 10^{-9}$

IBD, especially among children, is an emerging global epidemic [27] that is linked to autism [28, 29]. Impairment of intestinal barrier function is a core feature of IBD [30]. Increased intestinal permeability promotes infiltration of unmetabolized peptides into the lymph system and general circulation. This provides an opportunity for an immune antigenic response, which by molecular mimicry can lead to an attack on crucial proteins in the brain and spinal column. Disturbances of collagen texture are a major factor leading to the onset of diverticular disease and IBD along with the disturbed wound-healing mechanisms seen in the pathogenesis of anastomatic leakage following large bowel surgery [31].

In a recent paper [32], we suggested that glyphosate, a non-coding amino acid analogue of glycine, could substitute for glycine in error during protein synthesis. Such misincorporation and disruption of proteostasis could explain the strong correlations observed between glyphosate usage and multiple modern diseases. *In this paper, we show that this could be one of the most important mechanisms by which glyphosate could induce multiple autoimmune diseases.*

A prime site for initiation of the disease process is the colon, where misfolded collagen, resistant to degradation, could lead to an autoimmune disease and, subsequently, a leaky gut. Autoantibodies against type VII collagen have been detected in up to 68% of IBD patients [33]. Glycine is the most common amino acid in collagen, making up one fourth of the residues in the protein. Proline is also a very common component of collagen and, as we discuss later in this paper, proline resists hydrolysis. Incomplete collagen degradation by matrix metalloproteinases in the gut could lead to the accumulation of short pro–gly–pro peptides that are resistant to proteolysis. These could then induce the infiltration of neutrophils or the activation of resident immune cells to induce an inflammatory response [34].

An unpublished study conducted by Monsanto and submitted to the US Environmental Protection Agency (EPA) traced the accumulation of radiolabeled glyphosate in various tissues of rats following low-dose oral administration (10 mg/kg body weight) [35]. By far the highest accumulation was found in the bones (Table 11 in [36]). Radioactive levels in the colon were 4–6 times as high as those in the stomach and small intestine.

The production of novel non-coding amino acids by plants and microbes wards off predators. The toxicity of these products may be due to the fact that they replace coding analogues during protein synthesis. Examples include: azetidine-2-carboxylic acid (Aze), a proline analogue [37, 38]; glufosinate, a glutamate analogue that is also a popular herbicide [39];  $\beta$ -N-methylamino-L-alanine

(BMAA), an analogue of serine [40]; and L-canavanine, a natural analogue of L-arginine that is exploited as an insecticide [41, 42].

A remarkable true-life story involving a 119-day Alaskan wilderness experiment conducted by Christopher McCandless was recounted in the book *Into the Wild* by Jon Krakauer (later made into a popular movie) [43]. McCandless was thought to have died in the wilderness from starvation; however, Krakauer always suspected a toxin in the seeds of the wild potato, *Hedysarum alpinum*, which formed a staple of his diet in his last month of life. Krakauer had originally suspected a poisonous alkaloid but, through later research, was able to identify a significant level of L-canavanine in the wild potato seeds and published a paper on this analysis with several other authors in 2016 [42].

A key factor in L-canavanine's toxicity is its ability to insinuate itself into peptides in place of L-arginine. L-canavanine can be assimilated into essentially any protein to create aberrant canavanyl proteins that can disrupt many fundamentally important biochemical reactions across a broad spectrum of organisms [41, 44]. L-canavanine is exploited in agriculture as a potent insecticide against the tobacco hornworm [45], although the tobacco budworm has developed tolerance with a unique enzyme, canavanine hydrolase, which can quickly metabolize it [46]. Larvae exposed to L-canavanine incorporate it into the protein lysozyme, resulting in a 48% loss in catalytic activity [41]. Furthermore, dipterocins B and C of *Protoformia terranova*, but not dipterocin A, are negatively impacted by L-canavanine. The distinction is that dipterocin A has histidine at position 38 instead of the L-arginine found in the other two dipterocins. Presciently, with respect to glyphosate, Rosenthal wrote: "These insect studies support the view that the biological effects of canavanine result from its incorporation into a protein, resulting in an alteration in protein conformation that leads ultimately to impairment of protein function" [41].

## 2. SHIKIMATE PATHWAY INHIBITION REVISITED

The shikimate pathway enzyme, 5-enolpyruvylshikimate-3-phosphate synthase (EPSPS) is believed to be the main target of glyphosate's toxicity to plants [47]. A 1991 paper by Padgett et al. describes studies to gain insight into the mechanism by which glyphosate disrupts EPSPS [47]. Surprisingly, it is not understood exactly how glyphosate binds to the active site.

The microbes *Klebsiella pneumoniae*, *Escherichia coli* [47, 48] and *Agrobacterium sp.* strain CP4 [48, 49] have all evolved to produce versions of EPSPS that are glyphosate-resistant. The CP4 variant has been widely exploited by importing it into genetically modified

glyphosate-resistant crops [48]. Insight can be gained by investigating the alterations to the peptide sequence that afforded resistance. All three mutations involved replacing a glycine residue at the active site with alanine [47, 48]. In the case of *E. coli*, the mutated enzyme is about 72 times *less* efficient than the wild-type enzyme, but 69 times *more* efficient in the presence of glyphosate. Changing the DNA code from glycine to alanine completely disables glyphosate's inhibiting effects on the enzyme [48].

Substitution of gly-96 at the active site in *E. coli* by serine leads to a version of the enzyme that is unable to bind PEP, most likely due to steric hindrance. The authors speculated that the hydroxymethyl group of serine displaces the phosphate of PEP and functions as a nucleophile. In fact, this mutated enzyme achieves a kind of reverse reaction, breaking EPSP down into shikimate-3-phosphate and pyruvate via hydrolysis.

We propose that substitution of gly-96 (gly-100 in the CP4 variant) by glyphosate during protein synthesis could explain its disruption of the enzyme's function. One can expect that the highly reactive and bulky glyphosate molecule, if substituted for gly-96, would behave more like serine than alanine. An additional disruptive factor is glyphosate's chelation of manganese, which would disrupt the catalytic action of EPSPS. A cell containing both wild-type and glyphosate-substituted forms of the enzyme would arguably circuitously convert PEP to pyruvate via EPSP without producing ATP from ADP; i.e., would waste the energy in the phosphate bond, as shown in Fig. 1, and end up with excess pyruvate and a deficiency in EPSP.




Figure 1. Diagram of the hypothetical pathway by which glyphosate substitution for glycine in EPSPS could result in the synthesis of pyruvate from PEP without generating ATP; i.e., wasting the energy in the phosphate group, as discussed in the text.

### 3. GLYPHOSATE AS A GLYCINE ANALOGUE

While glyphosate's main mechanism of toxicity to plants is considered to be disruption of the shikimate pathway, it is also likely that it disrupts other biological pathways where glycine is either a substrate or a ligand, due to the fact that it is a glycine analogue. It has been proposed that, through glycine mimicry, glyphosate's rôle as a ligand to NMDA receptors in the brain could explain its known ability to activate NMDA receptors and cause neuronal damage [49, 50]. In [51], acute exposure of rat hippocampal slices to Roundup (0.00005–0.1%) for 30 minutes caused oxidative stress and neuronal cell death, which was attributed to NMDA receptor activation. Glyphosate also interferes with the synthesis of porphyrin, a precursor to haem, by disrupting the first step in the pathway where glycine is substrate [52].

N-substituted glycine "peptoids" are an attractive class of synthetic molecules that can be constructed by linking component N-substituted glycines at sequential nitrogen-carbon bonds; they are directly analogous to the linking of amino acids into peptides [53]. Glyphosate is of course an N-substituted glycine, where the nitrogen side chain is a methyl phosphonyl group. Part of the attraction of peptoids is that they are highly resistant to proteolysis, just as is the amino acid proline, in which the carbon side chain circles back and binds to the peptide nitrogen. Impaired ability to break down proline-rich gliadin has been proposed as a contributing factor in coeliac disease and gluten intolerance [54]. This can explain why common cereals with high proline contents are especially problematic to gluten-sensitive individuals [55, 56].

Glyphosate is probably particularly problematic when it substitutes for N-terminal glycines in proteins where these glycines are highly conserved and play a significant rôle. Several proteins rely on an N-terminal glycine for anchoring to the plasma membrane (e.g., endothelial nitric oxide synthase (eNOS) [57]) or to the cytoskeleton (e.g., Kelch-like ECH-associated protein 1 (KEAP1) [58]). Protein N-myristoylation and prenylation depend on an amide bond to the N-terminal glycine residue [59]. For example, myristoylated G proteins involved in many signaling mechanisms depend on an N-terminal glycine residue [59]. This would be disrupted if the nitrogen atom has a side chain through glyphosate substitution for the terminal glycine.

N-nitrosoamino acids form a reasonable model for N-nitrosoglyphosate, a carcinogenic derivative of glyphosate that was of concern to the EPA during Monsanto's early studies. N-nitrosoproline is particularly relevant because proline, like glyphosate, has an extra carbon atom bound to the nitrogen atom. With respect to non-coding amino acids, and especially the incorporation

of N-nitrosoamino acids into peptides and proteins, R.C. Massey remarked: "In addition to their presence as free N-nitrosoamino acids, species such as N-nitrosoproline (NPRO) and N-nitroso-4-hydroxyproline (HONPRO) may exist in a peptide- or protein-bound form as a result of N-nitrosation of an N-terminal imino acid residue" [62]. Tricker et al. [63] and Kubacki et al. [64] devised high performance liquid chromatography-thermal energy analyser (HPLC-TEA) techniques for analysis of multiple dipeptides with a nitrosylated N-terminal, including N-nitrosopropylalanine (NPROALA), N-nitrosopropyl-4-hydroxyproline (NPROHOPRO) and N-nitrosopropylglycine (NPROGLY) [63, 64]. Tricker notes that the average recoveries for NPROALA, NPROHOPRO and NPROGLY, 200 µg of which was added to cured meat, were between 69 and 88%. Tricker also used the method to analyse the nitroso-tripeptide N-nitrosopropylglycylglycine [65].

Nitrosamines of glyphosate (N-phosphonomethylglycine), its salts and esters include: N-nitrosoglyphosate (NNG) (Monsanto CP 76976), N-nitrosoiminodiacetic acid (NNIDA), N-nitrosoglyphosate sodium salt (NNGNa), N-nitrosoglyphosate isopropylamine ester (NNGIPA), N-nitrosoglyphosate potassium salt (NNGK), the metabolite N-nitrosoAMPA (NNAMPA), the metabolites N-nitrosodimethyl amine (NDMA) and N-nitrosarcosine (NSAR), which occur in glyphosate products or may be generated *in vivo* or in soils and waterways. N-nitroso compounds derived from secondary amines are considered carcinogenic.

Monsanto glyphosate documents reveal analysis and quantification of five nitrosamines of concern [61]. Out of six lots of Roundup analysed for NNG, four lots contained NNG residues of 0.61 to 0.78 ppm and two lots had residues from 0.22 to 0.40 ppm NNG. Analysis of six lots of Monsanto Rodeo revealed NNG residues in the range 0.13–0.49 ppm.

Recently, a powerful metatranscriptome study on bacterial gene expression following glyphosate treatment was conducted on microbes growing within the rhizosphere of glyphosate-tolerant corn [66]. RNA transcript abundance was compared between control and glyphosate-treated samples in order to characterize which protein genes were upregulated or downregulated. While they found many changes in gene expression, most striking to us was the upregulation of genes involved in both protein synthesis and protein hydrolysis. The ribosomal proteins L16p (L10e) and Firmicutes ribosomal L7Ae family proteins involved in the synthesis of the ribosomal large subunit increased 1.4- and two-fold, respectively, and the small subunit ribosomal protein S11p (S14e) increased 1.5-fold. Upregulation of genes involved in protein degradation was even more dramatic. For

example, transcripts for a proteasome  $\beta$  2 subunit (EC 3.4.25.1) increased 4.3-fold and aminopeptidase YpdF increased threefold. An explanation could be an increase in the number of proteins that fail to fold properly due to glyphosate substitution for glycine in the protein. These authors also suggested a potential shift towards an increase in glyphosate-tolerant bacteria, a point that will become important later in this paper.

These results are corroborated by a study on pea plants grown in hydroponic culture, which revealed that glyphosate induced a significant increase in two major systems for proteolytic degradation: the ubiquitin-26 S proteasome system and papain-like cysteine proteases [67]. It also increased the total free amino acid content and decreased the soluble protein in the root system.

#### 4. GLYPHOSATE-CONTAMINATED COLLAGEN AND PROTEOLYSIS RESISTANCE

We mentioned in the Introduction the gly-pro-gly peptide sequence that is common in collagen and linked to autoimmune disease. There are several enzymes in multiple organisms that are devoted to the proteolysis of peptide sequences containing proline, particularly the gly-pro sequence. These include enzymes that detach a terminal proline, enzymes that detach a dipeptide sequence where the second residue is a proline molecule and the first one is often glycine, and enzymes that break apart the X-pro dipeptide to release two free amino acids, one of which is proline. Certain pathogens have special modified versions of these enzymes, and there are genetic diseases related to pathologies in these enzymes. Substitution of glyphosate for glycine in this sequence is likely to cause extra stress to the enzymes that break down these sequences, potentially leading to autoimmune disease.

Prolyl aminopeptidase is an enzyme that detaches a terminal proline residue from a peptide. The enzyme is expressed predominantly by pathogenic bacteria in the gut, in particular *Serratia marcescens*, a common pathogen in the gut as well as in the urinary tract; it is often multiply antibiotic-resistant and is a serious threat in hospital-acquired infection [34]. This enzyme is especially important to the pathogens for degrading collagen, providing amino acids as fuel. It is conceivable that the pathogens are able to degrade glyphosate-contaminated peptides terminating in proline whereas the human form of the enzyme is not. It is intriguing that the *S. marcescens* version of prolyl aminopeptidase is unusual in having extra space at the active site [34], which could potentially accommodate the larger glyphosate molecule adjacent to the terminal proline residue. This might also contribute to glyphosate's observed effect on the gut microbiome: excessive growth of pathogens.

Multiple strains of the toxic mould *Aspergillus* secrete an X-prolyl dipeptidyl aminopeptidase (X-PDAP) that is important for digesting collagen because it can separate out an X-pro pair to bypass the difficult step of breaking the X-pro bond. Research has shown that this enzyme is essential for hydrolysing proline-containing peptides [69, 70]. It is likely that it becomes even more essential when X is glyphosate, as the peptoid sequence glyphosate-proline is likely almost impossible to break. Since gly-pro is a very common sequence in collagen, glyphosate-pro is likely to impede the breakdown of collagen fragments, which may then encourage *Aspergillus* infection in both plants and animals. Glyphosate has been shown to increase the growth rate of *Aspergillus* [71].

The most disturbing question is, what happens in the absence of pathogens that can effectively clear collagen peptides contaminated with glyphosate? As we will see later in this paper, antibodies to collagen are linked to antibodies to vaccines. A genetic defect in the enzyme prolylase, which can break apart the very common gly-pro dipeptide to release the individual amino acids, leads to a severe disease with mental deficiencies and multiple skin lesions [72]. Intriguingly, a common plant pathogen, *Xanthomonas campestris*, which causes blight on multiple plant species has a unique variant of prolylase with two mutations, a substitution of tyrosine for gly-385 and valine for tyr-387, two highly conserved residues in the peptide sequence [73]. Is it possible that swapping out glycine affords protection from glyphosate substitution for this residue? We hypothesize that peptides derived from multiple proline and glyphosate-contaminated proteins, which are highly resistant to proteolysis, are causing an autoimmune epidemic that is an important contributor to autism and other autoimmune disorders.

#### 5. BMAA AND ALS IN GUAM

$\beta$ -N-methylamino-L-alanine (BMAA) is another noncoding amino acid and an analogue of serine [40]. BMAA is synthesized by cyanobacteria, the microbes responsible for the toxic algal blooms that occur in lakes experiencing an accumulation of nitrogen and phosphate nutrients following hot, rainy weather [74]. An *in vitro* study by Dunlop et al. in 2013 demonstrated that BMAA can be misincorporated into human proteins, causing protein misfolding that could lead to neurological diseases [40].

BMAA has, in fact, been linked to several neurodegenerative diseases, including Parkinson's, Alzheimer's and amyotrophic lateral sclerosis (ALS) [75]. A 2013 study linked an ALS cluster in Chesapeake Bay to consumption of BMAA-contaminated crabs [76]. A study in France investigated an ALS cluster near a lagoon that supplied oysters and mussels to the local

population. The authors demonstrated that the shellfish were contaminated with BMAA, but also remarked that there was intensive chemical-based agriculture in the region [77]. Interestingly, cyanobacteria have been found to be remarkably resistant to glyphosate [78, 79], and this could contribute to the recent record-setting algal blooms in the Great Lakes region, where glyphosate is extensively used on genetically modified (GM) Roundup-Ready crops [80].

One likely molecule that could be adversely affected by BMAA is the glutamate transporter, whose defective expression has been linked to ALS [81]. Glutamate excitotoxicity in motor neurons is associated with ALS, and this could be caused by an impaired glutamate transport system. Ordinarily, astrocytes quickly clear glutamate from the synapse, following its release by neurons, and the transporter is essential for this clearance. A conserved serine-rich motif in the glutamate transporter forms a reentrant loop, similar to a structure found in many ion channels [82]. This loop is crucial for the enzyme's proper function, and would be disrupted by substitution of BMAA for serine.

An interesting detective story has evolved around an epidemic of a complex neurological condition termed amyotrophic lateral sclerosis–Parkinsonism dementia complex (ALS–PDC), which reached epidemic proportions during a short interval after World War II among the native Chamorro people on the small island of Guam in the South Pacific. At the peak of the epidemic, the natives had a hundredfold increased risk to ALS and Parkinson's disease compared to the risk in the general human population.

A plausible explanation for this epidemic relates to a popular native food source: seeds from the cycad trees [83–85]. Cycad seeds contain BMAA, likely derived from associated cyanobacteria. However, what is especially interesting is that the BMAA becomes concentrated in the skin of fruit bats that feed on the cycad seeds. Fruit bats were a popular delicacy among the natives, who ate every part of them, including the skin. Increased access to firearms from the USA during the war may have made it easier to kill the bats, on which the natives then feasted, ultimately leading to the natives' near-extinction through the accumulation of BMAA in their brains [86]. Meanwhile the near-extermination of the bats through the hunting removed the presumed source of the epidemic [83].

However, the warfare also led to the accumulation of many toxic chemicals in the soil, which could have encouraged the proliferation of cyanobacteria, which are especially resilient in the face of stressors. The bats' demise was undoubtedly hastened by the accumulation of

excess BMAA in their tissues. A measurement of the amount of BMAA in three dried specimens of fruit bats from Guam taken from a museum in Berkeley found concentrations between 1200 and 7500  $\mu\text{g/g}$ , which indicates up to hundredfold bioamplification over the level in the seeds of the cycad tree [87].

There have been inconsistent results in measuring the levels of BMAA in different tissue samples, but this has been explained recently by the realization that any BMAA incorporated into proteins may be missed in analysis without sufficient proteolysis. Ince et al. wrote: "When the insoluble, protein-containing fraction following TCA (trichloroacetic acid) extraction is further hydrolysed to release BMAA from protein, there is a further pool of protein-bound BMAA that is present in a ratio of between 60:1 and 120:1 compared with the pool of free BMAA" [84, p. 348]. We believe that this point has great significance when it comes to glyphosate: we highly suspect that different methodologies used to measure glyphosate contamination in any situation where there is a significant protein-bound component may yield different results depending on the degree to which protein hydrolysis is carried out.

## 6. GLYPHOSATE CONTAMINATION IN COLLAGEN, ENZYMES, GELATIN AND VACCINES

Gelatin is commonly used as an excipient stabilizer in vaccines, particularly the live virus vaccines. Gelatin is derived from animal skin and bone, especially of pigs and cattle; they may be fed glyphosate-contaminated forages, including GM Roundup-Ready corn and soy feed, which are sometimes supplemented with GM Roundup-Ready beet pulp. Gelatin is mainly derived by partial hydrolysis from the collagen in skin and bone. 26% of the amino acids in collagen are glycine; proline and hydroxyproline together make up 18% [88]; and glutamate constitutes 6%. All three of these components are problematic. The proline could be substituted by Aze from the sugar beet, the glycine could be substituted by residual glyphosate in the feed, and glutamate is a neurotransmitter but known to be neurotoxic at high concentrations; it works together with glycine to excite NMDA receptors in the brain. The vaccine virus may incorporate some of the noncoding amino acids into its own proteins to produce versions of them that resist proteolysis and induce autoimmunity through molecular mimicry.

One of us (Samsel) analysed a number of animal protein products for glyphosate. These included the bones of pigs, cows, horses' hooves, bees and bee products, collagen and gelatin products, vitamins, protein powders, enzymes and vaccines. Results are shown in Tables 2 and 3. Both high performance liquid

chromatography with tandem mass spectrometry (HPLC–MSMS) and enzyme-linked immunosorbent assay (ELISA) methods were utilized. It has been shown that both HPLC and ELISA are comparable in terms of accuracy and precision for detection and quantification of glyphosate in water-based analysis and including Nanopure, tap and river waters. Water-based solvents for

glyphosate demonstrate a detection limit of 0.6 ng/mL and a linear functional range of 1–25 ng/mL [200]. However, HPLC was not able to achieve detection below 5 ppb;<sup>1</sup> hence, in cases including water-based vaccines, analysis using numerous sample runs was made including using two independent labs to test the same samples.

Table 2. Residues of glyphosate found in animal-based products that were reported to the US Food and Drug Administration (FDA) by Samsel Environmental & Public Health Services. The limit of detection for glyphosate using hot water extraction is 0.075 parts per billion (ppb).<sup>1</sup>

Protein substrate	Type	Test date	Glyphosate residue (ppb) <sup>1</sup>
GELATIN	JELL-O ORANGE #07 JAN 2018 DB02 02:36	29 July 2016	9.00
GELATIN	POWER-MAX PROTEIN POWDER ADVANCED NUTRITION	29 July 2016	14.94
GELATIN	DISNEY GUMMIES VITAMINS	9 August 2016	8.27
GELATIN	FLINTSTONES GUMMIES VITAMINS	9 August 2016	5.32
ORAGEL	CHILDREN'S ORAGEL 7.5% BENZOCAINE FORMULA	26 September 2016	2.81

HPLC–MSMS was also later used, where the method detection limit (MDL) permitted, for additional confirmation and quantification of glyphosate in digestive enzymes and collagens. Spiked sample recoveries were done for all samples tested. Freshly prepared glyphosate standard solutions were run as controls and results were calculated based on a standard curve.

In 1989, Monsanto researchers conducted an experiment on exposure of bluegill sunfish to <sup>14</sup>C-radiolabeled glyphosate [89]. One of us (Samsel) obtained the (unpublished) report from the EPA through the Freedom of Information Act. The researchers had found that, with EDTA extraction, the amount of radiolabel in tissue samples was much higher than the amount of detected glyphosate. They decided to apply a digestive enzyme, proteinase K, and discovered that this “caused a substantial improvement in extractability”. It brought the yield from 17–20% in the case of EDTA to 57–70% following digestion with proteinase K. They summed up as follows: “Proteinase K hydrolyses proteins to amino acids and small oligopeptides, suggesting that a significant portion of the <sup>14</sup>C activity residing in the bluegill sunfish tissue was tightly associated with *or incorporated into protein*” (present authors’ emphasis). In this context it is important to recall that a 60- to 120-fold higher detection level of BMAA was obtained following protein *hydrolysis* of contaminated proteins [84].

Since Monsanto found bioaccumulation of glyphosate in all animal tissues, with the highest levels in the bones and marrow [35, 36], one would expect that all tissues derived from animals fed a diet containing glyphosate residues and used for food by people around the globe would be contaminated. Knowing that the bioaccumulation of glyphosate would be evident in the vast majority of animals raised for market and fed a contaminated diet, as well as their products; and suspecting the possibility of contamination of even the digestive enzymes derived from these animals, one of us (Samsel) decided to analyse random samples.

Results from various gelatin-based products, along with the results for several different vaccines (discussed later) were reported to the FDA by Samsel Environmental & Public Health Services in August 2016. Table 2 shows results for glyphosate residues found in these gelatin-based products. The highest level found in a gelatin sample was almost 15 ppb.<sup>1</sup>

Having found glyphosate in animal gelatins, analysing the collagen at the source was a logical next step. Tissues from pork and cattle obtained from a local supermarket, commercially available collagen sourced from industrially-raised swine and oxen, as well as the purified digestive enzymes pepsin, lipase and trypsin, derived from pigs, were selected for evaluation. Three methods of laboratory analysis were used to determine if

<sup>1</sup> Parts per (US) billion. To put this into perspective, 1 ppb = 1 µg/kg, and 1 µg of glyphosate (N-phosphonomethylglycine) contains  $3.561 \times 10^{12}$  molecules of the substance, each one of which could integrate with a protein.

glyphosate was present in porcine pepsin and in the glycine-rich collagen from the tissues of pigs and cattle, protein sources that are regularly consumed by Americans. The results are given in Table 3.

Glyphosate integration with enzymes is a serious consideration, as glyphosate may serve as an enzyme inhibitor like other phosphonates [90–92]. Inhibition and immobilization of enzymes may occur via three basic categories: covalent linkage; adsorption on a carrier; or entrapment within macromolecules [93].

Inhibition of enzymes may be reversible or irreversible. Types of reversible enzyme inhibition include competitive, noncompetitive and uncompetitive. *Irreversible* inhibitors covalently bond to the functional groups of the active site, thus permanently inactivating catalytic activity. Irreversible inhibition includes two types: group-specific inhibition and “suicide” inhibition.

The importance of fully functional digestive enzymes cannot be understated. They are essential for metabolic function, as they convert food into nutrients and other molecules that are then available to cells for tissue and organ growth, maintenance and repair. The precursor trypsinogen, produced in the pancreas, is enzymatically transformed into the serine protease trypsin. Trypsin catalyses the hydrolysis of proteins into peptides and provides substrates for further enzymatic hydrolysis for protein absorption.

Pepsin, a primary protease of digestion, is also responsible for the metabolism of dietary protein.

Pepsin’s cleavage of peptide bonds is responsible for the availability of the aromatic amino acids phenylalanine, tyrosine and tryptophan. It is also responsible for the cleavage and release of several other amino acids, including valine, glycine, histamine, glutamine, alanine and leucine.

Lipase participates in cell signaling, inflammation and metabolism. Pancreatic lipase is the catalyst for the hydrolysis of dietary lipids, which include fats, oils, cholesterol esters and triglycerides [94]. Triglyceride triester is metabolized for utilization as glucose and three fatty acids. Glyphosate integration into and inhibition of lipase could induce excessive bioaccumulation of fatty material in the blood vessels, gut, liver, spleen and other organs, as well as mimic lysosomal acid lipase deficiency. It would also allow for an increase in triglycerides in the blood, leading to numerous disease cascades, including malabsorption, fatty liver disease, jaundice, failure to thrive in infants, calcification of the adrenal gland, anaemia, hypercholesterolaemia, biliary dysfunction, decreased HDL, increased LDL, blood clots, fat-enlarged hepatocytes and liver fibrosis and failure. Samsel found that radiolabeled glyphosate was not detectable by HPLC–MSMS in samples of lipase deliberately spiked for analysis, suggesting that glyphosate may irreversibly inhibit lipase. On the other hand, pepsin and trypsin had good spike recoveries, demonstrating reversibility as glyphosate was released from the protein.

Table 3. Integration of glyphosate residues in various proteins, assessed using three testing methods.<sup>a</sup>

Protein substrate (Method)	Type	Glyphosate residue (ppb)
Bone (ELISA)	Bovine leg	11.56
Bone marrow (ELISA)	Bovine leg marrow	4.22
Bone (ELISA)	Porcine foot	9.81
Skin (ELISA)	Porcine	0.325
Gelatin (ELISA)	Bovine, Sigma Aldrich, gel strength 225 Type B	2.04
Collagen (ELISA)	Bovine I & III	120.18
Collagen (GC-MS)	Bovine I & III	130 µg/kg
Collagen (HPLC-MSMS)	Bovine I & III	95 µg/kg
Pepsin (ELISA)	Purified porcine enzyme	< 40.00
Pepsin (GC-MS)	Purified porcine enzyme	430 µg/kg
Pepsin (HPLC-MSMS)	Purified porcine enzyme	290 µg/kg
Trypsin (ELISA)	Purified porcine enzyme	61.99
Lipase (ELISA)	Purified porcine enzyme	24.43
Bee bread (HPLC-MSMS)	Bee bread	2300 µg/kg
Bees (HPLC-MSMS)	<i>Apis mellifera</i>	< 10 µg/kg trace
Honey & comb (HPLC-MSMS)	Honey	< 10 µg/kg trace

<sup>a</sup> The trace amount found in the bee substrates appeared as a small peak, which directly corresponded to glyphosate, complete with retention time and molecular features confirming contamination using HPLC–MSMS.



Table 3 shows results for various bovine and porcine products, including enzymes, bone, bone marrow, skin, collagen and gelatin. Acid hydrolysis was used on the bovine and porcine skin, bones and marrow, which were shaken and digested with 0.15 M hydrochloric acid for 24 h. The analysis methods were ELISA, gas chromatography–mass spectrometry (GC–MS) and HPLC–MSMS. All of the tested products were contaminated, with the highest level detected being 430 µg/kg in porcine pepsin (via GC–MS).

Additional evidence of glyphosate accumulation was found by Samsel in 2015 in the bodies of dead bees, bee bread and honey from bee hives suspected of colony collapse disorder (CCD), and these are also shown in the table. Colony collapse disorder (CCD) is an ever-increasing problem threatening pollination of crops globally. It may share a similar aetiology to that of Alzheimer’s disease with regard to learning and memory within the bee’s brain. Integration of glyphosate with the structural proteins and enzymes of the bee may affect protein folding and function. Additionally, glyphosate may also affect the digestive enzymes and bacterial homeostasis within the digestive system, which in turn may affect the quality of the honey produced. Glyphosate in bees may become part of their chitin, which has a structural function, in their bodies, analogous to glyphosate becoming part of the collagens of humans and other animals.

The results in Table 3 show ubiquitous contamination of the bee and bee products. Honey is derived from nectar and is the source of carbohydrates in the bee diet, whereas pollen turned into bee bread supplies the fats and proteins. Royal jelly, made from the secretions of the glands found in the hypopharynx of the worker bees, is fed to the queen and developing larvae [96].

Results for nineteen different vaccines, from five manufacturers, are shown in Table 4. Some vaccines do not contain live viruses and do not involve gelatin in their preparation, but many involve the use of eggs, bovine calf serum, fetal bovine serum or bovine proteins [95]. Engerix Hepatitis B vaccine is manufactured through a novel procedure, which involves culturing genetically engineered *Saccharomyces cerevisiae* yeast cells that carry the surface antigen gene of the hepatitis B virus. The procedures result in a product that can contain up to 5% yeast proteins, which could be a source of glyphosate if the yeast is grown on broths or media that utilize glyphosate-contaminated nutrient sources such as animal or plant proteins.

Vaccines that tested negative for glyphosate included Merck’s Hep-B vaccine, most of the pneumococcal vaccines and the sterile diluent included as a control. Gelatin is not listed as an ingredient in any of these vaccines, nor is bovine serum. In contrast, all of the vaccines that listed gelatin as an excipient tested positive for glyphosate, and nearly all of them also included bovine serum (including Varicella, MMR-II, MMRV and Zoster).

It is significant that MMR-II consistently contained the highest levels of glyphosate, significantly more than any of the other vaccines. This vaccine uses up to 12% hydrolysed gelatin as an excipient–stabilizer; as well as foetal bovine serum albumin, human serum albumin and residual chick embryo; all of which are contaminated by glyphosate during animal production.

## 7. EVIDENCE FOR A ROLE FOR COLLAGEN IN VACCINE ADVERSE REACTIONS

Post-vaccination allergic reactions to MMR and varicella vaccines have been linked to the gelatin excipient, and confirmed through observation of induced gelatin-specific IgE antibodies [97–100]. 24 out of 26 children with allergic reactions to vaccines (e.g., anaphylactic shock) had anti-gelatin IgE ranging from 1.2 to 250 µg/mL. Seven were allergic to gelatin-containing foods. A pool of 26 control children all tested negative for anti-gelatin IgE [99]. A study from 2009 that looked at gelatin sensitivity in children who were sensitive to cows’ milk, beef and/or pork as determined by IgE antibody levels [101] found that 16% of beef-sensitized children and 38% of pork-sensitized children had IgE antibodies to beef- or pork-derived gelatins that were cross-reactive with each other.

In a published case study, a 2-month-old baby developed Kawasaki disease one day after receiving its first dose of Infanrix (DTaP-IPV-Hib) and Prevenar, a pneumococcal conjugate vaccine [102]. Kawasaki disease is an acute, multisystemic vasculitis whose occurrence very early in life is extremely rare. Extensive tests for the presence of infection with multiple bacteria and viruses were all negative. We suggest that glyphosate contamination in one or both of the vaccines may have contributed to the vasculitis through glyphosate uptake into common proteins such as collagen in the vasculature to induce the autoimmune reaction.

Kelso (1993) reported the case of a 17-year-old girl who experienced anaphylaxis within minutes of receiving an MMR vaccine [98]. The girl described the event as “kind of like what happens when I eat Jell-O<sup>2</sup>”. Further testing found gelatin to be the component of the vaccine

<sup>2</sup> Jell-O is a proprietary brand of gelatin-based desserts, popular in the USA, and manufactured by Kraft Foods, part of the Kraft Heinz Company, headquartered in Chicago.

Table 4. Glyphosate levels in vaccines determined by ELISA reported to the US CDC, NIH, FDA and UN WHO of the Americas in September 2016 by Samsel Environmental & Public Health Services.<sup>a</sup>

Vaccine undiluted	Manufacturer	Lot number Exp date	Test date Lab #	Glyphosate residue (ppb)	% Recovery in spiked sample
DTaP ADACEL	SANOFI PASTEUR	58160-820-43	7-15-2016	0.109	82%
DTaP	SANOFI PASTEUR	NDC 3-30-2018 C50418A	LAB #1 5-11-2016	< 0.075	81%
DTaP ADACEL	SANOFI PASTEUR	9-2-2018 NDC 58160-820-43 3-30-2018	LAB #1 7-12-2016 LAB #2	ND	-
HEPATITIS-B	MERCK	LO16427	5-11-2016	< 0.075	97%
HEPATITIS ENGERIX-B	GLAXOSMITH- KLINE	NDC 58160-820-43 6-1-2018	7-15-2016 LAB #1	0.337	73%
INFLUENZA FLUZONE QUAD	SANOFI PASTEUR	6762	7-15-2016	0.170	95%
INFLUENZA	NOVARTIS	6-30-2016 1573 3P	LAB #1 5-11-2016	0.227	106%
Pneumococcal PNEUMOVAX 23	MERCK	05/2016 700281601	LAB #1 9-19-2016	0.112	118%
MMR II	MERCK	5-18-2017 7002151400	LAB #1 7-15-2016	3.740	-
MMR II	MERCK	9-9-2017 009545	LAB #1 5-11-2016	2.963	-
MMR II	MERCK	3-19-2017 7002151400	LAB #1 9-19-2016	3.154	-
MMR II	MERCK	9-9-2017 7002151400	LAB #1 7-12-2016	2.90	-
MMRV PROQUAD	MERCK	9-9-2017 7002305700	LAB #2 9-19-2016	0.659	103%
MMRV PROQUAD	MERCK	9-12-2017 7002305700	LAB #1 7-15-2016	0.512	86%
MRV PROQUAD	MERCK	9-12-2017 7002305700	LAB #1 7-12-2016	0.43	-
Pneumococcal PNEUMOVAX 23	MERCK	9-12-2017 700281601	LAB #2 7-15-2016	< 0.075	77%
Pneumococcal PREVNAR 13	WYETH	5-18-2017 73332	LAB #1 5-11-2016	< 0.075	82%
Pneumococcal PNEUMOVAX 23	MERCK	07/2017 7002681601	LAB #1 7-12-2016	ND	-
STERILE DILUENT	MERCK, SHARP & DOHME	5-18-2017 LO 40058	LAB #2 7-15-2016	< 0.075	97%
VARICELLA VARIVAX	MERCK	5-11-2018 7002025000	LAB #1 7-15-2016	0.556	84%
MVARICELLA VARIVAX	MERCK	2-8-2018 7002025000	LAB #1 7-12-2016	0.41	-
ZOSTER ZOSTAVAX	MERCK	2-8-2018 7002502401	LAB #2 9-19-2016	0.620	95%
ZOSTER ZOSTAVAX	MERCK	6-1-2017 7002602401	LAB #1 7-15-2016	0.558	98%
ZOSTER ZOSTAVAX	MERCK	6-1-2017 7002602401	LAB #1 7-12-2016	0.42	-
ZOSTER ZOSTAVAX	MERCK	6-1-2017 7002602401	LAB #2		

<sup>a</sup> Limits of detection for glyphosate in vaccines in parts per billion (ppb):<sup>1</sup> 0.075 (LAB #1); 0.15 (LAB #2).

to which the girl was allergic. The connexion may be to misfolded proteins, which include the collagens and associated partially hydrolysed gelatins. Indeed, both Jell-O and vaccines have been contaminated by glyphosate, as we reported in the previous section.

Puppies immunized with the rabies vaccine and a multivalent canine vaccine were compared to unvaccinated

control puppies [103]. The vaccinated puppies, but not the unvaccinated ones, developed autoantibodies to their own collagen. A follow-up study where either just the rabies vaccine or just the multivalent vaccine was administered produced a similar result. The authors suggested that this could explain issues of joint pain that are currently common among dogs, particularly as they age.

## 8. MULTIPLE SCLEROSIS (MS)

### 8.1 Sugar beet and MS

The world obtains 30% of its sugar supply from beet sugar. While sugar cane is grown in tropical regions, sugar beet requires a temperate climate. The highest incidences of MS worldwide are in the USA, Canada and western Europe [5], where most of the beet sugar is produced. MS rates are higher in the northern states of the USA compared to the south, corresponding to the distribution of sugar beet cultivation. MS rates in Canada are highest in the Alberta prairie region, at the centre of the Canadian sugar beet industry [104]. Studies on migrants have shown that those who move from a low-risk to a high-risk area tend to adopt high-risk only if they migrated during childhood [105]. This implicates local environmental factors acting before adolescence. Tokachi province in Japan hosts only 0.3% of the population, but produces 45% of the sugar beet consumed in Japan [37]; this province has the highest rate of MS among all Asian populations [106].

A fascinating proposition how sugar beet could cause MS implicates a unique noncoding amino acid that is produced by sugar beet, namely Aze. Both proline and Aze have a unique structure for an amino acid: the side chain loops back round to connect up to the nitrogen atom. In the case of Aze, there are only 3 carbons in the ring instead of the 4 carbons in proline (Fig. 2). It has been shown experimentally that Aze can be inserted by mistake into proteins in place of proline [38].

Myelin basic protein (MBP) is an essential protein for maintaining the myelin sheath, and it interacts with actin, tubulin, calmodulin and SH3 domains [107]. It

assembles actin filaments and microtubules, binds actin filaments and SH3 domains to membrane surfaces, and participates in signal transduction in oligodendrocytes and myelin. A central proline-rich region in MBP is functionally significant [108–110] and, in particular, is a binding site for Fyn-SH3, a key regulatory protein [111]. Proline substitutions of the SH3 ligand decrease its affinity for the Fyn-SH3 domain [108]. Fyn is localized to the cytoplasmic leaflet of the oligodendrocyte plasma membrane, where it participates in numerous signaling pathways during development of the central nervous system [112, 113]. Phosphorylation at a polyproline structure in the Fyn-binding region of MBP affects its structure.

A study using recombinant murine MBP inserted into *E. coli* strains demonstrated conclusively that Aze makes its way into MBP, substituting for up to three of the eleven possible proline sites. Molecular modeling of a proline-rich region of the recombinant MBP illustrated that misincorporation of Aze at any site would cause a severe bend in the polypeptide chain, and that multiple Aze substitutions would completely disrupt the structure of MBP [114, 115].

A possible concern regarding Aze is that over 90% of the sugar beet grown in the USA and Canada is genetically engineered to resist glyphosate. Therefore, the crops are exposed to significant amounts of glyphosate. The electronic *Code of Federal Regulations e-CFR 180.364 Glyphosate; Tolerances for Residues*, allows up to 25 ppm residue of glyphosate in dried sugar beet pulp. In 1999, Monsanto realized that its GM sugar beet crop well exceeded the upper limit established by the EPA for glyphosate residues. They requested, and were granted, a 125-fold increase in the upper residue limit for dried beet pulp (from 0.2 to 25 ppm). At the same time, the upper limit for fresh beet was increased fiftyfold to 10 ppm.

Glyphosate has been shown to increase the risk of root rot in sugar beet, caused by fungi [116]. Aze has been demonstrated to have antifungal activity [117]. Plants tend to increase synthesis of toxins under stress conditions, and it is plausible that an increased potential for root rot would result in increased synthesis of Aze. This is especially likely given that plants increase proline synthesis under a variety of different stress conditions [118]. However, to our knowledge, whether glyphosate causes an increase in either proline or Aze synthesis in sugar beet has not been investigated.

Consumption of milk worldwide is strongly correlated with MS risk (Spearman's correlation test = 0.836;  $P < 0.001$ ) [119]. For the past several decades, cows' feed has been supplemented with either beet




Figure 2. Molecular structures of the coding amino acids proline, L-arginine, glycine and glutamic acid; and their respective noncoding analogues Aze, L-canavanine, glyphosate and glufosinate.

molasses or sugar beet pulp, left as a residue after the sugar has been extracted [120]. Aze has been experimentally found in three sugar beet by-products that are fed to farm animals: sugar beet molasses, and both shredded and pelleted sugar beet pulp [38]. Casein is relatively enriched in proline [121]. If cows are exposed to Aze from the sugar beet, it will likely get inserted by mistake into casein, causing it to resist proteolysis. MBP's critical proline-rich sequence is vulnerable to misincorporation of Aze. The characteristic plaques of MS show loss of MBP within lesions in axon sheaths [107]. It is unclear whether this autoimmune reaction would arise through molecular mimicry from antibodies to unmetabolized peptides from casein or as a direct result of improperly folded MBP due to Aze insertion.

Glyphosate, an analogue of glycine, can be expected to be found in all tissues, including the milk of all mammals consuming glyphosate residues in the diet. Radiolabeled glyphosate studies conducted with lactating goats found  $^{13}\text{C}$  and  $^{14}\text{C}$  residues of glyphosate (N-phosphonomethylglycine), N-acetylglyphosate and other radiolabeled metabolites in milk. Monsanto found daily average  $^{14}\text{C}$  residue levels from 19 to 86 ppb, with levels falling after five days of depuration to 6 ppb prior to sacrifice for organ examination. Results disseminated by Monsanto indicate that lactating animals (goats) fed a diet containing glyphosate and AMPA can be expected to have measured residue levels in edible tissues and milk [122]. In 2007 Dupont, in a similar study, examined the metabolism of N-acetylglyphosate in lactating goats. Detectable residues of N-acetylglyphosate, glyphosate and AMPA were detected in milk and other tissues. Milk, liver and kidney each contained 0.03% of the administered dose. Individual daily radiolabeled residues in the milk ranged from 0.030 to 0.036  $\mu\text{g/g}$  [123].

*Lactobacillus* plays an important rôle in metabolizing casein in the human gut. A detailed study of the prolyl aminopeptidase from *Lactobacillus* revealed that it is a member of the class of  $\alpha/\beta$  hydrolases. Multiple sequence alignment has revealed three distinct highly conserved regions in this family and all three contain at least two highly conserved glycines [124] that would be vulnerable to displacement by glyphosate. The motif gly-x-ser-x-gly-gly characterizes the domain surrounding the catalytic serine residue of prolyl oligopeptidases in general. The glycine residues in this motif contribute to the correct positioning of the catalytic serine with respect to its substrate. A second glycine-rich domain appears essential to activity, as it likely corresponds to the oxyanion hole. The function of the third highly conserved glycine-rich domain, with the motif asp-x-x-gly-x-gly-x-ser, remains unknown. *Lactobacillus*

*spp.* are also highly dependent on manganese to protect them from oxidative damage, hence glyphosate's preferential chelation of manganese likely harms *Lactobacillus* [125].

An examination of collagen in the jugular veins of MS patients undergoing surgical reconstruction revealed an abnormal collagen structure, characterized by thin, loosely packed type III fibres [126]. Collagen is rich in proline. If too many of the prolines in procollagen are displaced by Aze, the polypeptide does not fold into a stable triple-helical conformation, which is a prerequisite for normal secretion of procollagen [127]. This reduces the release of procollagen and the misfolded molecules are subjected to proteolysis for recycling, resulting in the useless expenditure of energy for building and degrading procollagen molecules. Those that are released can be expected to produce defective collagen matrices. Collagen is even more highly enriched in glycine than in proline, as its core structure consists of a triple peptide repeat, where glycine is always the third residue of the triplet, and proline and hydroxyproline often occupy the other two positions [128]. Glyphosate substitution for glycine in structural proteins; i.e., collagen, elastin, fibronectin and laminin; would contribute to disrupted folding as well as defective strength and elasticity.

Conserved prolines also play a crucial rôle in ion channel gating, the regulation of hypoxia-inducible factor (HIF) and embryogenesis; in fact, substituting Aze for proline is a technique used to test whether a particular proline residue is critical to the protein's proper functioning [37].

## 8.2 Rôle of *Acinetobacter* and *Pseudomonas aeruginosa* in MS

A series of papers by Ebringer et al. have suggested an important rôle for the Gram-negative bacteria *Acinetobacter* and *Pseudomonas aeruginosa* in MS [129–131] as well as a proposed link to prion diseases. Their most recent paper in *Medical Hypotheses* presents the evidence to support this idea from multiple dimensions [130]. First, MS patients were shown to have elevated levels of antibodies to these two microbes but not to the common gut microbe *E. coli* [132, 116]. They have autoantibodies to MBP and myelin oligodendrocyte glycoprotein (MOG) [131]. MS patients are also prone to sinusitis and *Acinetobacter* is one of the most common microbes found in nasal sinuses. Ebringer et al. also proposed that the increased prevalence of sinusitis in colder climates may explain the geographical distribution of MS in more northerly latitudes [130]. *P. aeruginosa* causes upper respiratory infections and it is among the microbes that have developed multiple antibiotic

resistance in recent years, presenting a huge problem in hospital infection [133]. *Acinetobacter* has also become resistant to multiple antibiotics [134].

The number of microbial species that can metabolize glyphosate is quite small. A 1996 study showed that *Acinetobacter* is able to fully metabolize both glyphosate and AMPA and utilize these molecules as a source of phosphorus [135]. A study of agricultural soil heavily polluted with glyphosate identified only three species capable of degrading glyphosate when exposed at a level of 1000 ppm: *Pseudomonas putida*, *P. aeruginosa* and *Acetobacter faecalis* [136]. Another study on marine species identified *Pseudomonas* as being among the rare microbial species that can utilize the phosphonate in glyphosate as a source of phosphorus [137]. It can be predicted that *Pseudomonas* and *Acinetobacter* species in the nasal or digestive tracts would have a substantial advantage over other microbes if they can degrade glyphosate. On the other hand, they would also be heavily exposed if they actively take it up, and it would not be unreasonable to assume that some of the glyphosate might end up in their synthesized proteins by mistake in place of glycine. Both *Pseudomonas aeruginosa* and *Acinetobacter* strains have recently become a serious problem in hospitals, and a public health issue, due to their multiple-antibiotic resistance [138]. Glyphosate has been

shown to induce generic antibiotic resistance in other microbial species, including *E. coli* and *Salmonella*, through the induction of a generic capability to export toxic chemicals through efflux pumps [139].

A PEP transferase enzyme synthesized by *Acinetobacter calcaceticus* has sequence homology with a bovine prion sequence, and antibodies against synthetic peptides containing the structurally related sequences were found to be significantly elevated in cattle with bovine spongiform encephalopathy (BSE) compared to negative controls [140]. Ebringer et al. (2005) [129] link MS to BSE, also known as “mad cow disease”, and to the related human disease, Creutzfeldt–Jakob disease (CJD). Cows suffering from BSE manifest hindquarters paralysis early after onset, similar to the mobility issues afflicting MS patients at onset. Ebringer et al. found elevated levels of antibodies to both *Acinetobacter* and *Pseudomonas*, along with autoantibodies to both white and grey matter components, in BSE-affected animals, as is also the case for MS [129].

Of particular note are the molecular similarities they identified between certain peptides found in these two microbes and peptides in MOG and MBP that are known to be allergenic. Strikingly, all three of the microbial sequences they identified and all three of their human protein analogues contain conserved glycines (Table 5).

Table 5. Amino acid sequences of three peptides from *Acinetobacter* and *Pseudomonas* and the corresponding human peptides from MBP that they mimic.<sup>a</sup>

Microbe	<i>Acinetobacter</i>	<i>Acinetobacter</i>	<i>Pseudomonas</i>
Protein	3-OACT-A	4-CMLD	Gamma-CMLD
Peptide	Leu-Tyr-Arg-Ala-Gly-Lys	Ser-Arg-Phe-Ala-Tyr-Gly	Thr-Arg-His-Ala-Tyr-Gly
MBP	Leu-Tyr-Arg-Asp-Gly-Lys	Ser-Arg-Phe-Ser-Tyr-Gly	Ser-Arg-Phe-Ser-Tyr-Gly

<sup>a</sup> Note that all six peptides have a glycine residue.

MOG is strongly implicated in the disease pathology of MS; autoantibodies recognizing MOG have been found in the CNS of MS patients [141]. One of the major encephalitogenic peptides in MOG is the sequence from residue 92 to residue 106, which contains a highly conserved glycine near its centre [142].

Both diabetes and MS are associated with abnormal T-cell immunity to proteins found in cow’s milk [143]. In a study conducted in dairy cows by Monsanto in 1973, <sup>14</sup>C-radiolabeled glyphosate was studied in the distribution of residues in milk, urine, faeces and other tissues of the lactating cow. Glyphosate contamination of milk ranged from 9 to 15 ppb with the highest accumulation in the kidney and rumen fluid (201 ppb and 109 ppb, respectively) [201]. An epitope of bovine serine albumin found in milk that is linked to MS but not to diabetes is BSA193. It shows

structural homology with exon 2 of MBP through the peptide sequence GLCHMYK. Note that the first peptide in this sequence is glycine. Exon 2 is a target peptide in both MS autoimmunity and in experimental autoimmune encephalitis (EAE), an animal model of MS [144–146]. Exon 2 of MBP is implicated in remyelination [144]. Its expression is largely restricted to the developing brain and to areas of myelin reconstruction, notably MS lesions [147].

The gly-ser-gly-lys tetrapeptide is highly conserved among MBPs from multiple species [148]. The serine in this sequence is the site of attachment of polyphosphoinositide. The highly conserved nature of this sequence suggests that the phospholipidation of MBP is important biologically. Substitution of glyphosate for either of the glycines would likely disrupt this modification.

## 9. MMR VACCINE AND AUTISM

In this section, we make a case for a direct link between the measles, mumps, and rubella (MMR) vaccine and autism, via autoantibody induction through molecular mimicry. In a paper provocatively titled, “Peptide cross-reactivity: the original sin of vaccines”, Kanduc makes the point that massive cross-reactivity between antigens in vaccines and similar sequences in human proteins makes it almost inevitable that vaccines lead to autoimmune disease through molecular mimicry [149]. Reported post-vaccination autoimmune diseases include systemic lupus erythematosus, rheumatoid arthritis, inflammatory myopathies, multiple sclerosis, Guillain-Barré syndrome and vasculitis [150].

It is becoming increasingly acknowledged that autism may be an autoimmune disease. Family members of autistic children have a significant increased risk to other known autoimmune diseases such as hypothyroidism, rheumatic fever and multiple sclerosis [151]. Several studies on both humans and monkeys have revealed a potential link between maternal antibodies directed against specific foetal brain proteins and a future autism diagnosis in the foetus [152–155]. Furthermore, it has already been demonstrated that vaccines are capable of inducing autoimmune antibodies against proteins in the brain. The narcolepsy epidemic in Europe following an aggressive immunization campaign against the H1N1 ‘flu virus was eventually conclusively resolved as being attributed to autoimmune reactions to the hypocretin receptor through molecular mimicry from a peptide in the surface-exposed region of the influenza nucleoprotein A that was present in the H1N1 vaccine [156] (hypocretin is an important regulator of sleep).

Much controversy surrounds the concept that the MMR vaccine may be contributing to the autism epidemic in the USA and elsewhere. In an immune-compromised child, the live measles virus from the vaccine is capable of infecting the brain and sustaining a chronic measles infection, resulting in loss of neurons, eosinophilic intranuclear inclusions and gliosis, a condition termed “subacute measles encephalitis”. This can result in a seizure disorder and developmental delay in language and motor skills (as was clearly observed in a case study involving an HIV-positive 2-year-old boy [157]).

Singh et al. have published a series of papers over the past two decades [14, 158–160] proposing that there is a subpopulation among the autism community who can be characterized as suffering from “autoimmune autistic disorder” [14]. The 1998 study by Singh et al. found that 90% of measles-IgG-positive autistic sera were also positive for anti-MBP antibodies, supporting the hypothesis that a virus-induced autoimmune response may be

causal in autism [158]. A follow-on serologic study of antibodies to viruses associated with autism published in 2003 revealed a statistically significantly elevated level of measles antibody in children with autism compared to their siblings ( $P = 0.0001$ ) or to unrelated children ( $P = 0.003$ ), but not with antibodies to mumps or rubella [159]. In a later study, 60% of 125 autistic children had significantly elevated levels of antibodies to measles haemagglutinin unique to the MMR strain of the virus, compared to the 92 control children [160]. Over 90% of the children who had elevated antibody levels also tested positive for MBP autoantibodies. It was suggested that this could be linked to virus-induced autoimmunity through mimicry.

In fact, there is a sequence homology of 78% between a peptide sequence from MBP (EISFKLGQEGRDSRSGTP) and one found in a measles virus protein, MP3 (EISDNLGQEGRASTSGTP) [161, Table 2, p. 7]. Three of the matches between these two sequences are glycines. Measles virus-neutralizing antibodies are mainly directed to haemagglutinin, implying that it is essential for acquired immunity from the vaccine [162]; yet over-production, particularly if the virus penetrates the blood–brain barrier, runs the risk of inducing an autoimmune response to the myelin sheath. In fact, high measles antibody titres have been previously linked to MS [163].

Gonzalez-Granow et al. found high titres of autoantibodies in both the IgG and IgA classes specific to MBP in the serum of patients with autism [15]. The IgA antibodies in particular were shown to act as serine proteinases to degrade MBP *in vitro*. They also induced a decrease in long-term potentiation in perfused rat hippocampi. Reduced long-term potentiation in the hippocampus is a feature of autism, as has been clearly demonstrated in studies using mouse models of autism [164].

Dr Andrew Wakefield was the first to reveal a possible connexion between MMR and autism. His controversial *Lancet* paper, published in 1998 and then later retracted, proposed that this vaccine caused an acute reaction in children with gut dysbiosis (abdominal pain, diarrhoea, food intolerances, bloating etc.) [9]. The paper reported on a group of 12 children who had experienced developmental delay following an MMR vaccine and who were diagnosed with autism. These children suffered from rash, fever, delirium and seizures following the vaccination with MMR. He and several colleagues later published additional papers elaborating the hypothesis that dysbiosis in the gut, combined with impaired protein hydrolysis, leads to autoimmune lesions in the duodenum that are associated with extensive colonic lymphoid hyperplasia. The release of undigested peptides

into the vasculature across a leaky gut barrier and, ultimately, from the vasculature across a leaky blood–brain barrier, could induce encephalopathy [165–167].

In an epidemiological study from 1998, encephalopathy was clearly demonstrated as an acute reaction to measles vaccine, where 48 cases were found following vaccination, with no cases identified after administration of either monovalent mumps or rubella [168]. Among these 48 children, eight died, and the remainder experienced mental regression, chronic seizures, movement disorders and sensory deficits in the subsequent months.

The FDA’s vaccine adverse event reporting system (VAERS) database is a valuable tool for uncovering trends in vaccine adverse reactions. Our earlier studies on VAERS comparing MMR with an age-matched, equal-sized distribution of all other vaccines showed a significant association of MMR with autism ( $P < 0.007$ ) [169]. This was puzzling, because MMR has never contained either aluminium or mercury, the two prime candidates for the kind of neurological damage that might lead to autism [170–174]. Strong associations also appeared with fever and rash. In that paper, we proposed that the adverse reaction might be caused by the acetaminophen administered to the child to try to curb the seizures.

Since glyphosate usage on crops has gone up dramatically since the GM Roundup Ready crops were

first introduced in 1996, we decided it would be worthwhile to compare the early data on MMR in VAERS with the later data. We defined a cutoff date on 1 January 2003, such that the events where MMR was included as an administered vaccine could be separated into “early” and “late”, based on whether they were before or after that date. Each dataset represented a 13-year interval. We found 10 639 events in the early set and 19 447 events in the late set; thus, the raw number of events nearly doubled in the later years.

We also tabulated the frequency of different adverse reactions in the two sets, and used a standard statistical analysis to compute the significance of any differences observed: we randomly down-sampled both sets as needed such that there was an identical total count and an identical distribution over age in the two datasets. Results were surprising: many symptoms associated with atopy or with an allergic reaction were significantly higher in the later set, and “hospitalization” was highly significantly overrepresented in the later set [Table 6]. Other overrepresented symptoms included seizures, dyspnea, hyperventilation, asthma, eczema, autism, hives, anaphylactic [shock], and irregular heart rate. Interestingly, the early set had more frequent occurrences of joint pain and arthritis, suggesting that the toxic elements in the vaccine impacted the joints rather than the brain.

Table 6. Frequency of various adverse reactions to MMR before and after January 2003 [US FDA, VAERS]. The  $P$ -values were computed according to a  $\chi^2$  goodness-of-fit test.

More common before 2003			
Reaction	Count < 2003	Count $\geq$ 2003	$P$ -value
Arthritis	52	18	0.045
Joint pain	175	75	0.012
More common after 2002			
Reaction	Count < 2003	Count $\geq$ 2003	$P$ -value
Hospital	132	423	0.00041
Seizures	314	534	0.0055
Dyspnea	139	279	0.0086
Hives	444	654	0.011
Anaphylactic	28	91	0.017
Eczema	10	47	0.028
Autism	105	184	0.031
Hyperventilation	18	57	0.035
General infection	77	136	0.044
Asthma	22	58	0.046
Immunoglobulin G	0	17	0.048
Ear infection	32	72	0.048
Heart rate irregular	11	39	0.049

To our knowledge, there have been no significant changes to the formulation of MMR since its introduction. The explanation for the significant changes in adverse reactions must, therefore, lie in external factors, one of which is likely to be glyphosate. We suggest that both chronic exposure to glyphosate from food, water and air and direct exposure to glyphosate residues in the vaccine are relevant factors. A child with a disrupted gut microbiome due to chronic glyphosate exposure will also suffer from a leaky blood–brain barrier, and this will lead to a much greater possibility of measles antigenic proteins entering the brain and causing anaphylaxis and seizures.

The measles virus is a member of the family of paramyxoviruses, which have two highly-conserved glycine residues at positions 3 and 7 in the hydrophobic fusion peptide (FP) region of the viral fusion-mediating glycoproteins [175]. This FP region is the most highly conserved region of the glycoproteins, and it plays a critical rôle in destabilizing the membrane of the host cell to gain entry. Substitutions of other amino acids for either the G3A or G7A glycines caused increases in both cell–cell fusion and the reactivity of the protein to antibodies, leading to both a higher infection rate and increased chances for an autoimmune reaction. Glyphosate substitution is likely to do the same, as well as leading to a form of the protein that would resist proteolysis.

The FPs of both the influenza virus and human immunodeficiency virus (HIV) gp41 contain numerous glycine residues at regular intervals, with glycine overall making up 29 and 26%, respectively, of the total peptide sequence [175]. Optic neuritis, an immune-mediated demyelinating injury of the optic nerve, has been recognized as a side effect of the influenza vaccine that can lead to blindness [176].

## 10. OTHER AUTOIMMUNE DISEASES

### 10.1 Neuromyelitis optica and aquaporin

Neuromyelitis optica is a rare severe inflammatory demyelinating disorder of the central nervous system, which is related to multiple sclerosis but distinctly different and manifested mainly by paralysis and optic nerve damage [177, 178]. It has been conclusively demonstrated that this condition is caused by an autoimmune reaction to aquaporin-4, which is highly expressed in the astrocyte membrane [177, 178].

Aquaporins are important membrane proteins, which can transport water molecules through pores into the cell while excluding protons [179]. They are highly expressed by astrocytes, one of whose rôles is to mediate water flow among the vasculature, the

cerebrospinal fluid and the lymph system [178]. Thus, aquaporins are implicated in brain oedema [180]. Plants produce aquaporins as well, and mimicry between plant and human aquaporins has been proposed as a mechanism for the development of an autoimmune sensitivity to this protein [181]. Plants considered to show aquaporin mimicry notably include corn and soy as well as tomato, tobacco and spinach [182].

Autoimmune sensitivity to aquaporin has also been found in association with MS [182]. Vojdani et al. found significant elevations in antibodies against both human and plant aquaporin 4, in addition to antibodies against MB, MOG and S100 calcium-binding protein B (S100B) in patients suffering from MS.

Among the aquaporins, aquaporin-6 is unique in that it operates as an anion channel instead of as a water channel. Analysis of the peptide sequence in comparison to other aquaporins reveals that aquaporin-6 has an asparagine substituted in place of a glycine at residue 60. This one small difference completely changes the way the molecule behaves in the membrane. A glycine at this position is conserved among all the other aquaporins. Furthermore, aquaporins are constructed of  $\alpha$ -helices, and there are three sites where the helices cross. Highly conserved glycine residues are found at all three sites [57, 183].

Aquaporin is also found in bacteria, although homology with human aquaporin is only about 20%. The bacterial aquaporin is a 27 kDa trypsin-resistant protein called aquaporin-Z, which was originally described in *E. coli* [184]. Sequence analysis conducted by Ren et al. [185] revealed four regions where homology was considerably stronger (90%, 60%, 50% and 45% respectively). They convincingly showed cross immunoreactivity between the human and bacterial versions of the protein. Antibodies to aquaporin Z bind to astrocytes, activate complement, and cause death.

Ren et al. [185] identified all the residues where the bacterial and human peptides were identical (Fig. 1 in [185]). A tally of counts reveals that glycine was by far the most common among these matched residues, representing 14 of the total 66 matches. The second most common amino acid was lysine with 8 matches. Alanine, isoleucine and valine had 7, 5 and 4 matches respectively, and all other amino acids had less than four.

Thus, it appears that glyphosate-substituted trypsin-resistant aquaporin from both gut microbes and from GM glyphosate-resistant corn and soy foods are plausible sources of antigens that could induce neuromyelitis optica and contribute to the disease process in MS through misincorporation.



## 10.2 Type 1 diabetes

Type 1 diabetes is considered a genetic disease, but its incidence has been increasing by 3–4% worldwide every year in the recent past [186, 168]. Although an environmental component is highly suspected, environmental factors have not yet been identified. An increased incidence of type 1 diabetes is associated with both MS [187] and autism [188]. The disease is characterized by an autoimmune reaction to various proteins expressed in the pancreatic islet cells. Specifically, antibodies against glutamic acid decarboxylase (GAD65) are often found [189]. Cross-reactivity with proteins from foods and microbes in the gut are both possibilities.

One microbe that may be inducing antibody production through mimicry is *Mycobacterium avium paratuberculosis* (MAP). Blast analysis revealed 75% homology between a previously identified antigenic region of GAD65 [190] and a MAP heat-shock protein (HSP65) [189]. The specific 16-residue matched sequence in HSP65 centrally contains a pair of glycines which could be substituted by glyphosate to cause resistance to proteolysis. This microbe has been linked to numerous other human diseases including ulcerative colitis, irritable bowel syndrome, sarcoidosis, Hashimoto's thyroiditis, MS and autism [188]. With respect to MS and autism, cross-reactivity between HSP65 and MBP through mimicry may provide the link.

Patients with type-1 diabetes commonly have an antibody reaction to bovine serum albumin, a component of cows' milk [191]. The hypothesized explanation is an autoimmune reaction to a beta-cell specific surface protein through mimicry.

Insulin-derived amyloidosis is a condition that can develop following long-term insulin therapy, whereby an "insulin ball" develops at the site of injection. This hard mass has been analysed and found to contain accumulations of insulin fibrils reminiscent of amyloid  $\beta$ -plaque in the Alzheimer's brain. Insulin amyloidosis is more common for animal (cows and pigs)-derived than human-derived insulin products. Nowadays, cows and pigs are chronically exposed to glyphosate in their feed. The rôle of glycine residues in proteins may indeed be to protect from aggregation into amyloid fibrils [192]. Substitution of glyphosate for any of these conserved glycines would therefore tend to promote amyloidosis.

Glutamic acid and glycine are by far the largest component amino acids of bovine proinsulin and make up 25% of the amino acid residues in the molecule [193]. The same is true for human insulin, which differs very little from the animal versions. The herbicide glufosinate is a natural noncoding amino acid analogue of glutamic

acid (Fig. 2). Substitution of either glufosinate for glutamic acid or glyphosate for glycine in insulin is likely to impair its function, and may also lead to amyloidosis.

The widespread appearance of glyphosate-resistant weeds among the glyphosate-resistant crops has forced some farmers to turn to glufosinate as the herbicide of choice [194]. Glufosinate-tolerant corn and soybean have been available on the US market since their approval by the USDA in 1995 and 1996, respectively. A tri-resistant form of soybean tolerant of glyphosate, glufosinate, and 2,4-D was approved by the FDA in September 2014. Dual resistance to glufosinate and glyphosate in corn was approved in November 2015.

## 10.3 Coeliac disease

Coeliac disease and, more generally, gluten intolerance, have reached epidemic proportions in the USA in the past decade [195]. Wheat grown there is being routinely sprayed with glyphosate for staging and desiccation just before harvest. This practice clears the field of weeds prior to harvest and planting of the next crop, but increases the amount of residual glyphosate in the grain. The practice has been increasing in popularity in step with the increase in gluten intolerance. Glyphosate is systemic in the plant and enters the seed as the plant dies, hence eventually ending up in wheat-based foods.

Proline residues make up 20% of the first 100 amino acids of both  $\alpha$ - and  $\gamma$ -gliadins [54]. Related proteins from rye and barley are also unusually proline-rich [56]. As we implied earlier, proline is inaccessible to most digestive proteases because the bond between the peptide nitrogen atom and the side group complicates hydrolytic attack. As a consequence, specialized prolyl aminopeptidases detach the amino-terminal proline from a peptide. These enzymes depend on manganese as a catalyst, and manganese is one of the metals most dramatically affected by glyphosate chelation [125]. Unhydrolysed gliadin peptides bind to HLA-DQ molecules (receptors on antigen-presenting cells) and trigger pathogenic T-cell responses [196]. Genetic variants of HLA-DQ are linked to both coeliac disease and type 1 diabetes [197, 198].

Analysis of the X-ray crystal structure of a human cytosolic prolyl aminopeptidase worked out in 2008 revealed that it is a dimer with a dependency on two manganese ions as the catalytic centres [199]. The full sequence of the catalytic domains of six prolyl peptidases from both human and microbial species is shown in Fig. 6 in ref. 199. Six of the twenty sites of fully conserved residues across all species were glycine residues, three were histidine, two were tyrosine and two were proline. The remaining seven were seven different amino acids.

## 11. CONCLUSION

In this paper, we have shown that widespread misincorporation of glyphosate for glycine during protein synthesis could explain the aetiology of multiple autoimmune diseases that are currently increasing in incidence in the USA. Misincorporation is plausible by analogy with multiple known toxins produced by organisms in defence against pathogens, including Aze, BMAA, L-canavanine and glufosinate, which work in a similar manner. We have shown that proteins from foods such as milk, wheat and sugar beet, as well as peptides derived from microbes resident in the gut or nasal tract or introduced iatrogenically through vaccination, are all potential causes of autoimmune disease induced through molecular mimicry. It is highly significant that two microbes linked to MS through molecular mimicry are among the very few microbes that can fully metabolize glyphosate. Using the VAERS database, we have shown that severe adverse reactions to the MMR vaccine have increased significantly over the past decade in step with the increased use of glyphosate. Glyphosate in MMR may originate from growth of the live virus on culture materials derived from glyphosate-exposed animals and/or from gelatin used as an excipient stabilizer. We have confirmed the presence of glyphosate contamination in MMR and in many other vaccines where the live virus is cultured in eggs, bovine protein or gelatin, or where animal products are used as an excipient component. Notably, some vaccines prepared without live culture on gelatin were free of glyphosate contamination. Substitution of glyphosate for glycine during protein synthesis could yield a peptide that resists proteolysis, making it more likely to induce an immune response. Furthermore, enzymes involved in proteolysis are likely to be disrupted due to their confirmed contamination with glyphosate. A non-exhaustive list of possible diseases that can be attributed to this mechanism include autism, multiple sclerosis, type 1 diabetes, coeliac disease, inflammatory bowel disease and neuromyelitis optica.

## ACKNOWLEDGMENT

This research is supported in part by Quanta Computers, Taiwan, under the auspices of the Qmulus program.

## REFERENCES

- Ashwood, P. & van de Water, J. Is autism an autoimmune disease? *Autoimmun. Rev.* **3** (2004) 557–562.
- Gulcher, J.R., Vartanian, T. & Stefansson, K. Is multiple sclerosis an autoimmune disease? *Clin. Neurosci.* **2** (1994) 246–252.
- Hertz-Picciotto, I., Croen, L.A., Hansen, R., Jones, C.R., van de Water, J. & Pessah, I.N. The CHARGE study: An epidemiologic investigation of genetic and environmental factors contributing to autism. *Environ. Health Perspectives* **114** (2006) 1119–1125.
- London, E.A. The environment as an etiologic factor in autism: a new direction for research. *Environ. Health Perspectives* **108** (Suppl. 3) (2000) 401–404.
- Milo, R. & Kahana, E. Multiple sclerosis: Geoepidemiology, genetics and the environment. *Autoimmunity Rev.* **9** (2010) A387–A394.
- Koch-Henriksen, N. & Sorensen P.S. The changing demographic pattern of multiple sclerosis epidemiology. *Lancet Neurol.* **9** (2010) 520–532.
- Edwards, L.J. & Constantinescu, C.S. A prospective study of conditions associated with multiple sclerosis in a cohort of 658 consecutive outpatients attending a multiple sclerosis clinic. *Multiple Sclerosis* **10** (2004) 575–581.
- Kotey, S., Ertel, K. & Whitcomb, B. Co-occurrence of autism and asthma in a nationally-representative sample of children in the United States. *J. Autism Devl Disorders* **44** (2014) 3083–3088.
- Wakefield, A.J., Murch, S.H., Anthony, A., Linnell, J., Casson, D.M., Malik, M., Berelowitz, M., Dhillon, A.P., Thomson, M.A., Harvey, P., Valentine, A., Davies, S.E. & Walker-Smith, J.A. Ileal-lymphoid-nodular hyperplasia, non-specific colitis, and pervasive developmental disorder in children. *Lancet* **351** (1998) 637–641 (retracted).
- Seneff, S., Davidson, R.M. & Liu, J. Is cholesterol sulfate deficiency a common factor in preeclampsia, autism, and pernicious anemia? *Entropy* **14** (2012) 2265–2290.
- Gillberg, C., Gillberg, C. & Kopp, S. Hypothyroidism and autism spectrum disorders. *J. Child Psychol. Psychiat.* **33** (1992) 531–542.
- Miyazawa, M. Molecular mimicry and mechanisms of autoantibody production. *Nihon Rinsho* **55** (1997) 1370–1376 [in Japanese].
- Shoenfeld, Y.F. & Aron-Maor, A. Vaccination and autoimmunity/vaccinosis: A dangerous liaison? *J. Autoimmunity* **14** (2000) 1–10.
- Singh, V.K. Phenotypic expression of autoimmune autistic disorder (AAD): a major subset of autism. *Ann. Clin. Psychiat.* **21** (2009) 148–161.
- Gonzalez-Gronow, M., Cuchacovich, M., Francos, R., Cuchacovich, S., Blanco, A., Sandoval, R., Gomez, C.F., Valenzuela, J.A., Ray, R. & Pizzo, S.V. Catalytic autoantibodies against myelin basic protein (MBP) isolated from serum of autistic children impair *in vitro* models of synaptic plasticity in rat hippocampus. *J. Neuroimmunol.* **287** (2015) 1–8.
- Weizman, A., Weizman, R., Szekely, G.A., Wijzenbeak, H. & Livni, E. Abnormal immune response to brain tissue antigen in the syndrome of autism. *Am. J. Psychiatr.* **139** (1982) 1462–1465.
- Herroelen, L., de Keyser, J. & Ebinger, G. Central nervous system demyelination after immunization with recombinant Hepatitis B vaccine. *Lancet* **338** (1991) 1174–1175.
- Genain, C.P., Cannella, B., Hauser, S.L. & Raine, C.S. Identification of autoantibodies associated with myelin damage in multiple sclerosis. *Nature Med.* **5** (1999) 170–175.
- Fredman, P., Vedeler, C.A., Nyland, H., Aarli, J.A. & Svennerholm, L. Antibodies in sera from patients with

- inflammatory demyelinating polyradiculoneuropathy reactive with ganglioside LM1 and sulfate of peripheral nerve myelin. *J. Neurol.* **238** (1991) 75–79.
20. Steinman, L. Multiple sclerosis: A two-stage disease. *Nature Immunol.* **2** (2001) 762–764.
  21. Drummond, D.A. & Wilke, C.O. The evolutionary consequences of erroneous protein synthesis. *Nature Rev. Genetics* **10** (2009) 715–724.
  22. Drummond, D.A. & Wilke, C.O. Mistranslation-induced protein misfolding as a dominant constraint on coding-sequence evolution. *Cell* **134** (2008) 341–352.
  23. Conrad, A., Schröter-Kermani, C., Hoppe, H.W., Rütger, M., Pieper, S. & Kolossa-Gehring, M. Glyphosate in German adults—time trend (2001 to 2015) of human exposure to a widely used herbicide. *Intl J. Hyg. Environ. Health* **220** (2017) 8–16.
  24. Swanson, N.L., Leu, A., Abrahamson, J. & Wallet, B. Genetically engineered crops, glyphosate and the deterioration of health in the United States of America. *J. Org. Syst.* **9** (2014) 6–37.
  25. Hoy, J., Swanson, N. & Seneff, S. The high cost of pesticides: Human and animal diseases. *Poultry Fish. Wildlife Sci.* **3** (2015) 132.
  26. Seneff, S., Swanson, N. & Li, C. Aluminum and glyphosate can synergistically induce pineal gland pathology: connection to gut dysbiosis and neurological disease. *Agric. Sci.* **6** (2015) 42–70.
  27. Malmborg, P. & Hildebrand, H. The emerging global epidemic of paediatric inflammatory bowel disease—causes and consequences. *J. Intern. Med.* **279** (2016) 241–258.
  28. Boorom, K.F. Is this recently characterized gastrointestinal pathogen responsible for rising rates of inflammatory bowel disease (IBD) and IBD associated autism in Europe and the United States in the 1990s? *Med. Hypotheses* **69** (2007) 652–659.
  29. Horvath, K., Papadimitriou, J.C., Rabsztyrn, A., Drachenberg, C. & Tildon, J.T. Gastrointestinal abnormalities in children with autistic disorder. *J. Pediatrics* **135** (1999) 559–563.
  30. Michielan, A. & D’Inca, R. Intestinal permeability in inflammatory bowel disease: pathogenesis, clinical evaluation, and therapy of leaky gut. *Mediators Inflammation* **2015** (2015) 628157.
  31. Stumpf, M., Krones, C.J., Klinge, U., Rosch, R., Junge, K. & Schumpelick, V. Collagen in colon disease. *Hernia* **10** (2006) 498–501.
  32. Samsel, A. & Seneff, S. Glyphosate, pathways to modern diseases V: Amino acid analogue of glycine in diverse proteins. *J. Biol. Phys. Chem.* **16** (2016) 9–46.
  33. Hundorfean, G., Neurath, M.F. & Sitaru, C. Autoimmunity against type VII collagen in inflammatory bowel disease. *J. Cell Molec. Med.* **14** (2010) 2393–2403.
  34. Koelink, P.J., Overbeek, S.A., Braber, S., Morgan, M.E., Henricks, P.A., Roda, A., Verspaget, H.W., Wolfkamp, S.C., te Velde, A.A., Jones, C.W., Jackson, P.L., Blalock, J.E., Sparidans, R.W., Kruijtzter, J.A., Garssen, J., Folkerts, G. & Kraneveld, A.D. Collagen degradation and neutrophilic infiltration: a vicious circle in inflammatory bowel disease. *Gut* **63** (2014) 578–587.
  35. Ridley, W.P. & Mirly, K. The metabolism of glyphosate in Sprague Dawley rats. Part I. Excretion and tissue distribution of glyphosate and its metabolites following intravenous and oral administration (unpublished study MSL-7215 conducted by Monsanto’s Environmental Health Laboratory and submitted to the EPA July 1988) (MRID#407671-01) (1988).
  36. Samsel, A. & Seneff, S. Glyphosate, pathways to modern diseases IV: cancer and related pathologies. *J. Biol. Phys. Chem.* **15** (2015) 121–159.
  37. Rubenstein, E. Misincorporation of the proline analog azetidine-2-carboxylic acid in the pathogenesis of multiple sclerosis: a hypothesis. *J. Neuropathol. Exp. Neurol.* **67** (2008) 1035–1040.
  38. Rubenstein, E., McLaughlin, T., Winant, R.C., Sanchez, A., Eckart, M., Krasinska, K.M. & Chien, A. Azetidine-2-carboxylic acid in the food chain. *Phytochemistry* **70** (2009) 100–104.
  39. Hoerlein, G. Glufosinate (phosphinothricin), a natural amino acid with unexpected herbicidal properties. *Rev. Environ. Contamination Toxicol.* **138** (1994) 73–145.
  40. Dunlop, R.A., Cox, P.A., Banack, S.A. & Rodgers, K.J. The non-protein amino acid BMAA is misincorporated into human proteins in place of L-serine causing protein misfolding and aggregation. *PLoS ONE* **8** (2013) e75376.
  41. Rosenthal, G.A. The biochemical basis for the deleterious effects of L-canavanine. *Phytochemistry* **30** (1990) 1055–1058.
  42. Krakauer, J., Long, Y., Kolbert, A., Thanedar, S. & Southard, J. Presence of L-canavanine in *Hedysarum alpinum* seeds and its potential rôle in the death of Chris McCandless. *Wilderness Environ. Med.* **26** (2015) 36–42.
  43. Krakauer, J. *Into the Wild*. New York: Anchor Books (1996).
  44. Rosenthal, G.A. Biochemical basis for the deleterious effects of L-canavanine. *Phytochemistry* **30** (1991) 1055–1058.
  45. Dahlman, D.L. & Rosenthal, G.A. Non-protein amino acid-insect interactions I. Growth effects and symptomology of L-canavanine consumption by tobacco hornworm, *Manduca sexta* (L.). *Comparative Biochem. Physiol. A* **51** (1975) 33–36.
  46. Melangeli, C., Rosenthal, G.A. & Dalmank, D.L. The biochemical basis for l-canavanine tolerance by the tobacco budworm *Heliothis virescens* (Noctuidae). *Proc. Natl Acad. Sci. USA* **94** (1997) 2255–2260.
  47. Padgett, S.R., Re, D.B., Gasser, C.S., Eichholtz, D.A., Frazier, R.B., Hironaka, C.M., Levine, E.B., Shah, D.M., Fraley, R.T. & Kishore, G.M. Site-directed mutagenesis of a conserved region of the 5-enolpyruvylshikimate-3-phosphate synthase active site. *J. Biol. Chem.* **266** (1991) 22364–22369.
  48. Eschenburg, S., Healy, M.L., Priestman, M.A., Lushington, G.H. & Schonbrunn, E. How the mutation glycine 96 to alanine confers glyphosate insensitivity to 5-enolpyruvyl shikimate-3-phosphate synthase from *Escherichia coli*. *Planta* **216** (2002) 129–135.
  49. Funke, T., Han, H., Healy-Fried, M.L., Fischer, M. & Schonbrunn, E. Molecular basis for the herbicide resistance of Roundup Ready crops. *Proc. Natl Acad. Sci. USA* **103** (2006) 13010–13015.

50. Beecham, J.E. & Seneff, S. The possible link between autism and glyphosate acting as glycine mimetic—a review of evidence from the literature with analysis. *J. Molec. Genet. Med.* **9** (2015) 187.
51. Cattani, D., de Liz Oliveira Cavalli, V.L., Heinz Rieg, C.E., Domingues, J.T., Dal-Cim, T., Tasca, C.I., Mena Barreto Silva, F.R. & Zamoner, A. Mechanisms underlying the neurotoxicity induced by glyphosate-based herbicide in immature rat hippocampus: involvement of glutamate excitotoxicity. *Toxicology* **320** (2014) 34–45.
52. Kitchen, L.M., Witt, W.W. & Rieck, C.E. Inhibition of  $\delta$ -aminolevulinic acid synthesis by glyphosate. *Weed Sci.* **29** (1981) 571–577.
53. Zuckermann, R.N., Martin, E.J., Spellmeyer, D.C., Stauber, G.B., Shoemaker, K.R., Kerr, J.M., Figliozzi, G.M., Goff, D.A., Siani, M.A., Simon, R.J. et al. Discovery of nanomolar ligands for 7-transmembrane G-protein-coupled receptors from a diverse N-(substituted)glycine peptoid library. *J. Med. Chem.* **37** (1994) 2678–2685.
54. Hausch, F., Shan, L., Santiago, N.A., Gray, G.M. & Khosla, C. Intestinal digestive resistance of immunodominant gliadin peptides. *Am. J. Physiol. Gastrointestinal Liver Physiol.* **283** (2002) G996–G1003.
55. Schuppan, D. Current concepts of celiac disease pathogenesis. *Gastroenterology* **119** (2000) 234–242.
56. Wieser, H. Relation between gliadin structure and coeliac toxicity. *Acta Paediatr. (Suppl.)* **412** (1996) 3–9.
57. Liu, J. & Sessa, W.C. Identification of covalently bound amino-terminal myristic acid in endothelial nitric oxide synthase. *J. Biol. Chem.* **269** (1994) 11691–11694.
58. Kang, M.-I., Kobayashi, A., Wakabayashi, N., Kim, S.-G. & Yamamoto, M. Scaffolding of Keap1 to the actin cytoskeleton controls the function of Nrf2 as key regulator of cytoprotective phase 2 genes. *Proc. Natl Acad. Sci. USA* **101** (2004) 2046–2051.
59. Aicart-Ramos, C., Valero, R.A. & Rodriguez-Crespo, I. Protein palmitoylation and subcellular trafficking. *Biochim. Biophys. Acta* **1808** (2011) 2981–2994.
60. Kleuss, C. & Krause, E. G $\alpha$ s is palmitoylated at the N-terminal glycine. *EMBO J.* **22** (2003) 826–832.
61. Hirsch, R.H., Augustin, D.J. Nitrosamine analyses of Roundup herbicide, Rodeo herbicide, MON 0139 and Polado Technical (unpublished study RD835). St Louis, Missouri: Monsanto Agricultural Company (4 November 1987).
62. Massey, R.C. Analysis of N-nitroso compounds in foods and human body fluids. In: *Nitrosamines Toxicology and Microbiology* (ed. M.H. Hill), p. 26, section 2.4.4. VCH (1988).
63. Tricker, A.R., Perkins, M.J., Massey, R.C. & McWeeny, D.J. Some nitrosoamino acids in bacon adipose tissue and their contribution to the total N-nitroso compound concentration. *Z. Lebensmittel Untersuchung Forschung* **180** (1985) 379–383.
64. Kubacki, S.J., Havery, D.C. & Fazio, T. Nonvolatile N-nitrosamine investigations: methods for the determination of N-nitrosoamino acids and preliminary results of the development of a method for the determination of nitrosopeptides N-terminal in proline. In: *N-Nitroso Compounds: Occurrence, Biological Effects and Relevance to Human Cancer* (eds I.K. O’Neill, R.C. von Borstel, C.T. Miller, J. Long & H. Bartsch), No. 57, pp. 145–158. Lyons: International Agency for Research on Cancer (1984).
65. Tricker, A.R., Perkins, M.J., Massey, R.C. & McWeeny, D.J. Characterization studies on insoluble total N-nitroso compounds in bacon adipose connective tissue. *Food Additives Contaminants* **3** (1986) 153–159.
66. Newman, M.M., Lorenz, N., Hoilett, N., Lee, N.R., Dick, R.P., Liles, M.R., Ramsier, C. & Kloepper, J.W. Changes in rhizosphere bacterial gene expression following glyphosate treatment. *Sci. Total Environ.* **553** (2016) 32–41.
67. Zulet, A., Gil-Monreal, M., Villamor, J.G., Zabalza, A., van der Hoorn, R.A. & Royuela, M. Proteolytic pathways induced by herbicides that inhibit amino acid biosynthesis. *PLoS ONE* **8** (2013) e73847.
68. Nakajima, Y., Ito, K., Sakata, M., Xu, Y., Nakashima, K., Matsubara, F., Hatakeyama, S. & Yoshimoto, T. Unusual extra space at the active site and high activity for acetylated hydroxyproline of prolyl aminopeptidase from *Serratia marcescens*. *J. Bacteriol.* **188** (2006) 1599–1606.
69. Beauvais, A., Monod, M., Wyniger, J., Debeaupuis, J.P., Grouzmann, E., Brakch, N., Svab, J., Hovanessian, A.G. & Latgé, J.P. Dipeptidyl-peptidase IV secreted by *Aspergillus fumigatus*, a fungus pathogenic to humans. *Infection Immunity* **65** (1997) 3042–3047.
70. Byun, T., Kofod, L., Blinkovsky, A. Synergistic action of an X-prolyl dipeptidyl aminopeptidase and a non-specific aminopeptidase in protein hydrolysis. *J. Agric. Food Chem.* **49** (2001) 2061–2063.
71. Barberis, C.L., Carranza, C.S., Chiacchiera, S.M., Magnoli, C.E. Influence of herbicide glyphosate on growth and aflatoxin B1 production by *Aspergillus* section Flavi strains isolated from soil on in vitro assay. *J. Environ. Sci. Health B* **48** (2013) 1070–1079.
72. Freij, B.J., Levy, H.L., Dudin, G., Mutasim, D., Deeb, M., Der Kaloustian, V.M. Clinical and biochemical characteristics of prolidase deficiency in sibs. *Am. J. Med. Genet.* **19** (1984) 561–571.
73. Kumar, A., Are, V.N., Ghosh, B., Agrawal, U., Jamdar, S.N., Makde, R.D., Sharma, S.M. Crystallization and preliminary X-ray diffraction analysis of Xaa-Pro dipeptidase from *Xanthomonas campestris*. *Acta Crystallogr. F. Struct. Biol. Commun.* **70** (2014) 1268–1271.
74. Davis, T.W., Berry, D.L., Boyer, G.L. & Gobler, C.J. The effects of temperature and nutrients on the growth and dynamics of toxic and non-toxic strains of *Microcystis* during cyanobacteria blooms. *Harmful Algae* **8** (2009) 715–725.
75. Al-Sammak, M.A., Rogers, D.G. & Hoagland, K.D. Acute  $\alpha$ -N-methylamino-L-alanine toxicity in a mouse model. *J. Toxicol.* **2015** (2015) 739–746.
76. Field, N.C., Metcalf, J.S., Caller, T.A., Banack, S.A., Cox, P.A. & Stommel, E.W. Linking  $\beta$ -N-methylamino-L-alanine exposure to sporadic amyotrophic lateral sclerosis in Annapolis, MD. *Toxicol.* **70** (2013) 179–183.
77. Masseret, E., Banack, S., Boumédiène, F., Abadie, E., Briant, L., Pernet, F., Juntas-Morales, R., Pageot, N., Metcalf, J., Cox, P. & Camu, W. French Network on ALS Clusters Detection and Investigation. Dietary BMAA exposure in an amyotrophic lateral sclerosis cluster from southern France. *PLoS ONE* **8** (2013) e83406.

78. Powell, H.A., Kerby, N.W. & Rowell, P. Natural tolerance of cyanobacteria to the herbicide glyphosate. *New Phytol.* **119** (1991) 421–426.
79. Forlani, G., Pavan, M., Gramek, M., Kafarski, P. & Lipok, J. Biochemical bases for a widespread tolerance of cyanobacteria to the phosphonate herbicide glyphosate. *Plant Cell Physiol.* **49** (2008) 443–456.
80. Michalak, A.M., Anderson, E.J., Beletsky, D., Boland, S., Bosch, N.S., Bridgeman, T.B., Chaffin, J.D., Cho, K., Confesor, R., Daloglu, I., DePinto, J.V. et al. Record-setting algal bloom in Lake Erie caused by agricultural and meteorological trends consistent with expected future conditions. *Proc. Natl Acad. Sci. USA* **110** (2013) 6448–6452.
81. Foran, E. & Trotti, D. Glutamate transporters and the excitotoxic path to motor neuron degeneration in amyotrophic lateral sclerosis. *Antioxidant Redox Signaling* **11** (2009) 1587–1602.
82. Slotboom, D.J., Sobczak, I., Konings, W.N. & Lolkema, J.S. A conserved serine-rich stretch in the glutamate transporter family forms a substrate-sensitive reentrant loop. *Proc. Natl Acad. Sci. USA* **96** (1999) 14282–14287.
83. Cox, P.A. & Sacks, O.W. Cycad neurotoxins, consumption of flying foxes, and ALS-PDC disease in Guam. *Neurology* **58** (2002) 956–959.
84. Ince, P.G. & Codd, G.A. Return of the cycad hypothesis does the amyotrophic lateral sclerosis/parkinsonism dementia complex (ALS/PDC) of Guam have new implications for global health? *Neuropathol. Appl. Neurobiol.* **31** (2005) 345–353.
85. Steele, J.C. & McGeer, P.L. The ALS/PDC syndrome of Guam and the cycad hypothesis. *Neurology* **70** (2008) 1984–1990.
86. Monson, C.S., Banack, S.A. & Cox, P.A. Conservation implications of Chamorro consumption of flying foxes as a possible cause of Amyotrophic Lateral Sclerosis Parkinsonism dementia complex in Guam. *Conservation Biol.* **17** (2003) 678–686.
87. Banack, S.A. & Cox, P.A. Biomagnification of cycad neurotoxins in flying foxes: implications for ALS-PDC in Guam. *Neurology* **61** (2003) 387–389.
88. Eastoe, J.E. The amino acid composition of mammalian collagen and gelatin. *Biochem. J.* **61** (1955) 589–600.
89. Ridley, W.P. & Chott, K.A. Uptake, depuration and bioconcentration of C-14 glyphosate to bluegill sunfish (*Lepomis macrochirus*) Part II: Characterization and quantitation of glyphosate and its metabolites. St Louis, Missouri: Monsanto Agricultural Company (unpublished study) (August 1989).
90. Yang, K.W., Brandt, J.J., Chatwood, L.L., Crowder, M.W. Phosphoramidate and phosphothioate dipeptides as potential inhibitors of VanX. *Bioorg. Med. Chem. Lett.* **10** (2000) 10850–10857.
91. Perlikowska, R., Fichna, J., do-Rego J.C., Gach, K., Janecka, A. Kinetic studies of novel inhibitors of endomorphin degrading enzymes. *Med. Chem. Res.* **21** (2012) 1445–1450.
92. Kramer, G.J., Mohd, A., Schwager, S.L.U., Masuyer, G., Acharya, K.R., Sturrock, E.D. & Bachmann, B.O. Interkingdom pharmacology of angiotensin-I converting enzyme inhibitor phosphonates produced by Actinomycetes. *ACS Med. Chem. Lett.* **5** (2014) 346–351.
93. Rubin, B. & Dennis, E. *Lipases, Part B: Enzyme Characterization and Utilization* (vol. 286). Academic Press (1997).
94. Chapus, C., Rovey, M., Sarda, L. & Verger, R. Minireview on pancreatic lipase and colipase. *Biochimie* **70** (1988) 1223–1234.
95. *Epidemiology and Prevention of Vaccine-Preventable Diseases*, 13th edn, Appendix B. Centers for Disease Control and Prevention (2015).
96. Graham, J. (ed.). *The Hive and the Honey Bee* (rev. edn). Watertown, Wisconsin: Dadant & Sons (1992).
97. Pool, V., Braun, M.M., Kelso, J.M., Mootrey, G., Chen, R.T., Yunginger, J.W., Jacobson, R.M., Gargiullo, P.M. & VAERS Team. US Vaccine Adverse Event Reporting System. Prevalence of anti-gelatin IgE antibodies in people with anaphylaxis after measles-mumps rubella vaccine in the United States. *Pediatrics* **110** (2002) e71.
98. Kelso, J.M., Jones, R.T. & Yunginger, J.W. Anaphylaxis to measles, mumps, and rubella vaccine mediated by IgE to gelatin. *J. Allergy Clin. Immunol.* **91** (1993) 867–872.
99. Sakaguchi, M., Nakayama, T. & Inouye, S. Food allergy to gelatin in children with systemic immediate-type reactions, including anaphylaxis, to vaccines. *J. Allergy Clin. Immunol.* **98** (1996) 1058–1061.
100. Sakaguchi, M., Yamanaka, T., Ikeda, K., Sano, Y., Fujita, H., Miura, T. & Inouye S. IgE-mediated systemic reactions to gelatin included in the varicella vaccine. *J. Allergy Clin. Immunol.* **9** (1997) 263–264.
101. Bogdanovic, J., Halsey, N.A., Wood, R.A. & Hamilton, R.G. Bovine and porcine gelatin sensitivity in milk and meat-sensitized children. *J. Allergy Clin. Immunol.* **124** (2009) 1108–1110.
102. Ece, I., Akbayram, S., Demiroren, K. & Uner, A. Is Kawasaki Disease a side effect of vaccination as well? *J. Vaccines Vaccination* **5** (2014) 234.
103. Hogenesch, H., Azcona-Olivera, J., Scott-Moncrieff, C., Snyder, P.W. & Glickman, L.T. Vaccine-induced autoimmunity in the dog. *Adv. Vet. Med.* **41** (1999) 733–747.
104. Beck, C.A., Metz, L.M., Svenson, L.W. & Patten, S.B. Regional variation of multiple sclerosis prevalence in Canada. *Multiple Sclerosis* **11** (2005) 516–519.
105. Gale, C.R. & Martyn, C.N. Migrant studies in multiple sclerosis. *Prog. Neurobiol.* **47** (1995) 425–448.
106. Houzen, H., Niino, M., Kikuchi, S., Fukazawa, T., Nogoshi, S., Matsumoto, H. & Tashiro, K. The prevalence and clinical expression of MS in northern Japan. *J. Neurol. Sci.* **211** (2003) 49–53.
107. Boggs, J.M. Myelin basic protein: a multifunctional protein. *Cell Molec. Life Sci.* **63** (2006) 1945–1961.
108. Smith, G.S., De Avila, M., Paez, P.M., Spreuer, V., Wills, M.K., Jones, N., Boggs, J.M. & Harauz, G. Proline substitutions and threonine pseudophosphorylation of the SH3 ligand of 18.5-kDa myelin basic protein decrease its affinity for the Fyn-SH3 domain and alter process development and protein localization in oligodendrocytes. *J. Neurosci. Res.* **90** (2012) 28–47.
109. Harauz, G. & Libich, D.S. The classic basic protein of myelin conserved structural motifs and the dynamic molecular barcode involved in membrane adhesion and protein-protein interactions. *Current Protein Peptide Sci.* **10** (2009) 196–215.

110. Homchaudhuri, L., Polverini, E., Gao, W., Harauz, G. & Boggs, J.M. Influence of membrane surface charge and post-translational modifications to myelin basic protein on its ability to tether the Fyn-SH3 domain to a membrane in vitro. *Biochemistry* **48** (2009) 2385–2393.
111. Machold, R., Hayashi, S., Rutlin, M., Muzumdar, M.D., Nery, S., Corbin, J.G., Gritli-Linde, A., Dellovade, T., Porter, J.A., Rubin, L.L., Dudek, H., McMahon, A.P. & Fishell, G. Sonic hedgehog is required for progenitor cell maintenance in telencephalic stem cell niches. *Neuron* **39** (2003) 937–950.
112. Manié, S.N., Astier, A., Haghayeghi, N., Canty, T., Druker, B.J., Hirai, H. & Freedman, A.S. Regulation of integrin-mediated p130(Cas) tyrosine phosphorylation in human B cells. A role for p59(Fyn) and SHP2. *J. Biol. Chem.* **272** (1997) 15636–15641.
113. Resh, M.D. Fyn, a Src family tyrosine kinase. *Intl J. Biochem. Cell Biol.* **30** (1998) 1159–1162.
114. Bessonov, K., Vassall, K.A. & Harauz, G. Parameterization of the proline analogue Aze (azetidine-2-carboxylic acid) for molecular dynamics simulations and evaluation of its effect on homo-pentapeptide conformations. *J. Molec. Graphics Modelling* **39** (2013) 118–125.
115. Bessonov, K., Bamm, V.V. & Harauz, G. Misincorporation of the proline homologue Aze (azetidine-2-carboxylic acid) into recombinant myelin basic protein. *Phytochemistry* **71** (2010) 502–507.
116. Kiewnick, S., Jacobsen, B.J., Braun-Kiewnick, A., Eckhoff, J.L.A. & Bergman, J.W. Integrated control of Rhizoctonia crown and root rot of sugar beet with fungicides and antagonistic bacteria. *Plant Diseases* **85** (2001) 718–722.
117. Bach, B., Gregson, R.P., Holland, G.S., Quinn, R.J. & Reichelt, J.L. L-Azetidine-2-carboxylic acid, the antidermatophyte constituent of two marine sponges. *Experientia* **34** (1978) 688.
118. Hayat, S., Hayat, Q., Alyemeni, M.N., Wani, A.S., Pichtel, J. & Ahmad, A. Role of proline under changing environments: a review. *Plant Signaling Behaviour* **7** (2012) 1456–1466.
119. Malosse, D., Perron, H., Sasco, A. & Seigneurin, J.M. Correlation between milk and dairy product consumption and multiple sclerosis prevalence: A worldwide study. *Neuroepidemiology* **11** (1992) 304–312.
120. Huhtanen, P. The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fiber digestion in the rumen of cattle given a silage diet. *Animal Feed Sci. Technol.* **20** (1988) 259–278.
121. Gordon, W.G., Semmett, W.F. & Alanine, M.B. Glycine and proline contents of casein and its components. *J. Am. Chem. Soc.* **72** (1950) 4282–4282.
122. Bodden, R.M., Patanella, J.E., Feng, P. *Metabolism Study of Synthetic <sup>13</sup>C/<sup>14</sup>C- Labeled Glyphosate and AMPA In Lactating Goats*, vols 1 & 2 (unpublished study). St. Louis, Missouri: Monsanto Company (February 1988).
123. Lowrie, C. Metabolism of [<sup>14</sup>C]-N-Acetylgllyphosate (IN-MCX20) in the Lactating Goat (Charles River Laboratories Project no. 210583, submitted by E.I. du Pont de Nemours and Company) (Report No DuPont-19796) (2007).
124. Morel, F., Gilbert, C., Geourjon, C., Frot-Coutaz, J., Portalier, R. & Atlan, D. The prolyl aminopeptidase from *Lactobacillus delbrueckii* subsp. *bulgaricus* belongs to the u/v hydrolase fold family. *Biochim. Biophys. Acta* **1429** (1999) 501–505.
125. Samsel, A. & Seneff, S. Glyphosate, pathways to modern diseases III: Manganese neurological diseases and associated pathologies. *Surg. Neurol. Intern.* **6** (2015) 45.
126. Coen, M., Menegatti, E., Salvi, F., Mascoli, F., Zamboni, P., Gabbiani, G. & Bochaton-Piallat, M.L. Altered collagen expression in jugular veins in multiple sclerosis. *Cardiovasc. Pathol.* **22** (2013) 33–38.
127. Tan, E.M., Ryhänen, L. & Uitto, J. Proline analogues inhibit human skin fibroblast growth and collagen production in culture. *J. Investigative Dermatol.* **80** (1983) 261–267.
128. Bhattacharjee, A. & Bansal, M. Collagen structure: the madras triple helix and the current scenario. *IUBMB Life* **57** (2005) 161–172.
129. Ebringer, A., Rashid, T. & Wilson, C. Bovine spongiform encephalopathy, multiple sclerosis, and Creutzfeldt-Jakob disease are probably autoimmune diseases evoked by *Acinetobacter* bacteria. *Ann. NY Acad. Sci.* **1050** (2005) 417–428.
130. Ebringer, A., Rashid, T. & Wilson, C. The role of *Acinetobacter* in the pathogenesis of multiple sclerosis examined by using Popper sequences. *Med. Hypotheses* **78** (2012) 763–769.
131. Hughes, L.E., Smith, P.A., Bonell, S., Natt, R.S., Wilson, C., Rashid, T., Amor, S., Thompson, E.J., Croker, J. & Ebringer, A. Cross-reactivity between related sequences found in *Acinetobacter* sp., *Pseudomonas aeruginosa*, myelin basic protein and myelin oligodendrocyte glycoprotein in multiple sclerosis. *J. Neuroimmunol.* **144** (2003) 105–115.
132. Hughes, L.E., Bonell, S., Natt, R.S., Wilson, C., Tiwana, H., Ebringer, A., Cunningham, P., Chamoun, V., Thompson, E.J., Croker, J. & Vowles, J. Antibody responses to *Acinetobacter* spp. and *Pseudomonas aeruginosa* in multiple sclerosis: prospects for diagnosis using the myelin *Acinetobacter* neurofilament antibody index. *Clin. Diagnostic Lab. Immunol.* **8** (2001) 1181–1188.
133. Lister, P.D., Wolter, D.J. & Hanson, N.D. Antibacterial-resistant *Pseudomonas aeruginosa*: Clinical impact and complex regulation of chromosomally encoded resistance mechanisms. *Clin. Microbiol. Rev.* **22** (2009) 582–610.
134. Karageorgopoulos, D.E. & Falagas, M.E. Current control and treatment of multidrug-resistant *Acinetobacter baumannii* infections. *Lancet Infectious Diseases* **8** (2008) 751–762.
135. Chung, N.-J., Han, H.-J., Lee, H.-H., Rhie, H.-G. & Lee, H.-S. Degradation of phosphonate herbicide glyphosate by *Acinetobacter lwoffii* HN401. *Molecules Cells* **6** (1996) 239–245.
136. Olawale, A.K. & Akintobi, O.A. Biodegradation of glyphosate pesticide by bacteria isolated from agricultural soil. *Report Opinion* **3** (2011) 124–128.
137. Moore, J.K., Braymer, H.D. & Larson, A.D. Isolation of a *Pseudomonas* sp. which utilizes the phosphonate herbicide glyphosate. *Appl. Environ. Microbiol.* **46** (1983) 316–320.
138. Landman, D., Quale, J.M., Mayorga, D., Adediji, A., Vangala, K., Ravishankar, J., Flores, C. & Brooks, S.

- Citywide clonal outbreak of multiresistant *Acinetobacter baumannii* and *Pseudomonas aeruginosa* in Brooklyn, NY: The preantibiotic era has returned. *Arch. Intern. Med.* **162** (2002) 1515–1520.
139. Kurenbach, B., Marjoshi, D., Amabile-Cuevas, C.F., Ferguson, G.C., Godsoe, W., Gibson, P. & Heinemann, J.A. Sublethal exposure to commercial formulations of the herbicides dicamba, 2,4-dichlorophenoxyacetic acid, and glyphosate cause changes in antibiotic susceptibility in *Escherichia coli* and *Salmonella enterica* serovar typhimurium. *nBio* **6** (2015) e00009.
  140. Wilson, C., Hughes, L., Rashid, T., Cunningham, P., Bansal, S., Ebringer, A. & Ettelaie, C. Antibodies to prion and *Acinetobacter* peptide sequences in bovine spongiform encephalopathy. *Vet. Immunol. Immunopathol.* **98** (2004) 1–7.
  141. O'Connor, K.C., Appel, H., Bregoli, L., Call, M.E., Catz, I., Chan, J.A., Moore, N.H., Warren, K.G., Wong, S.J., Hafler, D.A. & Wucherpfennig, K.W. Antibodies from inflamed central nervous system tissue recognize myelin oligodendrocyte glycoprotein. *J. Immunol.* **175** (2005) 1974–1982.
  142. Clements, C.S., Reid, H.H., Beddoe, T., Tynan, F.E., Perugini, M.A., Johns, T.G., Bernard, C.C. & Rossjohn, J. The crystal structure of myelin oligodendrocyte glycoprotein, a key autoantigen in multiple sclerosis. *Proc. Natl Acad. Sci. USA* **100** (2003) 11059–11064.
  143. Winer, S., Astsaturov, I., Cheung, R.K., Schrade, K., Gunaratnam, L., Wood, D.D., Moscarello, M.A., O'Connor, P., McKerlie, C., Becker, D.J. & Dosch, H.M. T cells of multiple sclerosis patients target a common environmental peptide that causes encephalitis in mice. *J. Immunol.* **166** (2001) 4751–4756.
  144. Segal, B.M., Raine, C.S., McFarlin, D.E., Voskul, R.R. & McFarland, H.F. Experimental allergic encephalomyelitis induced by the peptide encoded by exon 2 of the MBP gene, a peptide implicated in remyelination. *J. Neuroimmunol.* **51** (1994) 7–19.
  145. Fritz, R.B. & Zhao, M.L. Encephalitogenicity of myelin basic protein exon-2 peptide in mice. *J. Neuroimmunol.* **51** (1994) 1–6.
  146. Voskuhl, R.R., Robinson, E.D., Segal, B.M., Tranquill, L., Camphausen, K., Albert, P.S., Richert, J.R. & McFarland, H.F. HLA restriction and TCR usage of T lymphocytes specific for a novel candidate autoantigen, X2 MBP, in multiple sclerosis. *J. Immunol.* **153** (1994) 4834–4844.
  147. Capello, E.R., Voskuhl, R.R., McFarland, H.F. & Raine, C.S. 1997. Multiple sclerosis: re-expression of a developmental gene in chronic lesions correlates with remyelination. *Ann. Neurol.* **41** (1997) 797–805.
  148. Chang, P.C., Yang, J.C., Fujitaki, J.M., Chiu, K.C. & Smith, R.A. Covalent linkage of phospholipid to myelin basic protein: identification of serine-54 as the site of attachment. *Biochemistry* **25** (1986) 2682–2686.
  149. Kanduc, D. Peptide cross-reactivity: the original sin of vaccines. *Frontiers Biosci.* **4** (2012) 1393–1401.
  150. Orbach, H., Agmon-Levin, N. & Zandman-Goddard, G. Vaccines and autoimmune diseases of the adult. *Discovery Med.* **9** (2010) 90–97.
  151. Sweeten, T.L., Bowyer, S.L., Posey, D.J., Halberstadt, G.M. & McDougle, C.J. Increased prevalence of familial autoimmunity in probands with pervasive developmental disorders. *Pediatrics* **112** (2003) e420–e424.
  152. Brimberg, L., Sadiq, A., Gregersen, P.K. & Diamond, B. Brain-reactive IgG correlates with autoimmunity in mothers of a child with an autism spectrum disorder. *Molec. Psychol.* **18** (2013) 1171–1177.
  153. Bauman, M.D., Iosif, A. M., Ashwood, P., Braunschweig, D., Lee, A., Schumann, C.M. et al. Maternal antibodies from mothers of children with autism alter brain growth and social behavior development in the rhesus monkey. *Transl. Psychiat.* **3** (2013) e278.
  154. Atladóttir, H.O., Pedersen, M.G., Thorsen, P., Mortensen, P.B., Deleuran, B., Eaton, W.W. & Parner, E.T. Association of family history of autoimmune diseases and autism spectrum disorders. *Pediatrics* **124** (2009) 687–694.
  155. Braunschweig, D., Krakowiak, P., Duncanson, P., Boyce, R., Hansen, R.L., Ashwood, P. et al. Autism-specific maternal autoantibodies recognize critical proteins in developing brain. *Transl. Psychiat.* **3** (2013) e277.
  156. Ahmed, S.S., Volkmuth, W., Duca, J., Corti, L., Pallaoro, M., Pezzicoli, A., Karle, A., Rigat, F., Rappuoli, R., Narasimhan, V., Julkunen, I., Vuorela, A., Vaarala, O., Nohynek, H., Laghi Pasini, F., Montomoli, E., Trombetta, C., Adams, C.M., Rothbard, J. & Steinman, L. Antibodies from vaccine-associated narcolepsy sera cross-reacted with both influenza nucleoprotein and hypocretin receptor 2. *Sci. Translational Med.* **7** (2015) 294ra105.
  157. Poon, T.P., Tchertkoff, V. & Win, H. Subacute measles encephalitis with AIDS diagnosed by fine needle aspiration biopsy. A case report. *Acta Cytol.* **42** (1998) 729–733.
  158. Singh, V.K., Lin, S.X. & Yang, V.C. Serological association of measles virus and human herpes virus-6 with brain autoantibodies in autism. *Clin. Immunol. Immunopathol.* **89** (1998) 105–108.
  159. Singh, V.K. & Jensen, R.L. Elevated levels of measles antibodies in children with autism. *Pediat. Neurol.* **28** (2003) 292–294.
  160. Singh, V.K., Lin, S.X., Newell, E. & Nelson, C. Abnormal measles-mumps-rubella antibodies and CNS autoimmunity in children with autism. *J. Biomed. Sci.* **9** (2002) 359–364.
  161. Oldstone, M.B.A. (ed.). *Molecular Mimicry: Infection-Inducing Autoimmune Disease*. Springer (2006).
  162. de Swart, R.L., Yüksel, S. & Osterhaus, A.D.M.E. Relative contributions of measles virus hemagglutinin- and fusion protein-specific serum antibodies to virus neutralization. *J. Virol.* **79** (2005) 11547–11551.
  163. Alter, M. Is multiple sclerosis an age-dependent host response to measles? *Lancet* **28** (1976) 456–457.
  164. Kaphzan, H., Hernandez, P., Jung, J., Cowansage, K.K., Deinhardt, K., Chao, M.V., Abel, T. & Klann, E. Reversal of impaired hippocampal long-term potentiation and contextual fear memory deficits in Angelman syndrome model mice by ErbB inhibitors. *Biol. Psychiat.* **72** (2012) 182–190.
  165. Kawashima, H., Mori, T., Kashiwagi, Y., Takekuma, K., Hoshika, A. & Wakefield, A. Detection and sequencing of measles virus from peripheral mononuclear cells from patients with inflammatory bowel disease and autism. *Digestive Diseases Sci.* **45** (2000) 723–729.

166. Torrente, F., Ashwood, P., Day, R., Machado, N., Furlano, R.I., Anthony, A., Davies, S.E., Wakefield, A.J., Thomson, M.A., Walker-Smith, J.A. & Murch, S.H. Small intestinal enteropathy with epithelial IgG and complement deposition in children with regressive autism. *Molec. Psychol.* **7** (2002) 375–382.
167. Wakefield, A.J., Puleston, J.M., Montgomery, S.M., Anthony, A., O’Leary, J.J. & Murch, S.H. Review article: the concept of entero-colonic encephalopathy, autism and opioid receptor ligands. *Alimentary Pharmacol. Therapeut.* **16** (2002) 663–674.
168. Weibel, R.E., Caserta, V., Benor, D.E. & Evans, G. Acute encephalopathy followed by permanent brain injury or death associated with further attenuated measles vaccines: A review of claims submitted to the national vaccine injury compensation program. *Pediatrics* **101** (1998) 383–387.
169. Seneff, S., Davidson, R.M. & Liu, J. Empirical data confirm autism symptoms related to aluminum and acetaminophen exposure. *Entropy* **14** (2012) 2227–2253.
170. Dufault, R., Schnoll, R., Lukiw, W.J., LeBlanc, B., Cornett, C., Patrick, L., Wallinga, D., Gilbert, S.G. & Crider, R. Mercury exposure, nutritional deficiencies and metabolic disruptions may affect learning in children. *Behavioral Brain Functions* **5** (2009) 44.
171. Sharpe, M.A., Gist, T.L. & Baskin, D.S. B-lymphocytes from a population of children with autism spectrum disorder and their unaffected siblings exhibit hypersensitivity to thimerosal. *J. Toxicol.* **2013** (2013) 801517.
172. Shaw, C.A., Kette, S.D., Davidson, R.M. & Seneff, S. Aluminum’s role in CNS-immune system interactions leading to neurological disorders. *Immunome Res.* **9** (2013) 069.
173. Shaw, C.A., Seneff, S., Kette, S.D., Tomljenovic, L., Oller, J.W., Jr. & Davidson, R.M. Aluminum-induced entropy in biological systems: implications for neurological disease. *J. Toxicol.* **2014** (2014) 491316.
174. Tomljenovic, L. & Shaw, C.A. Mechanisms of aluminum adjuvant toxicity and autoimmunity in pediatric populations. *Lupus* **21** (2012) 223–230.
175. Russell, C.J., Jardetzky, T.S. & Lamb, R.A. Conserved glycine residues in the fusion peptide of the paramyxovirus fusion protein regulate activation of the native state. *J. Virol.* **78** (2004) 13727–13742.
176. Kawasaki, A., Purvin, V.A. & Tang, R. Bilateral anterior ischemic optic neuropathy following influenza vaccination. *J. Neuroophthalmol.* **18** (1998) 56–59.
177. Papadopoulos, M.C. & Verkman, A.S. Aquaporin 4 and neuromyelitis optica. *Lancet Neurol.* **11** (2012) 535–544.
178. Roemer, S.F., Parisi, J.E., Lennon, V.A., Benarroch, E.E., Lassmann, H., Bruck, W., Mandler, R.N., Weinshenker, B.G., Pittock, S.J., Wingerchuk, D.M. & Lucchinetti, C.F. Pattern-specific loss of aquaporin-4 immunoreactivity distinguishes neuromyelitis optica from multiple sclerosis. *Brain* **130** (2007) 1194–1205.
179. Liu, K., Kozono, D., Kato, Y., Agre, P., Hazama, A. & Yasui, M. Conversion of aquaporin 6 from an anion channel to a water-selective channel by a single amino acid substitution. *Proc. Natl Acad. Sci. USA* **102** (2005) 2192–2197.
180. Zador, Z., Bloch, O., Yao, X. & Manley, G.T. Aquaporins: role in cerebral edema and brain water balance. *Prog. Brain Res.* **161** (2007) 185–194.
181. Vaishnav, R.A., Liu, R., Chapman, J., Roberts, A.M., Ye, H., Rebolledo-Mendez, J.D., Tabira, T., Fitzpatrick, A.H., Achiron, A., Running, M.P. & Friedland, R.P. Aquaporin 4 molecular mimicry and implications for neuromyelitis optica. *J. Neuroimmunol.* **260** (2013) 92–98.
182. Vojdani, A., Mukherjee, P.S., Berookhim, J. & Kharrazian, D. Detection of antibodies against human and plant aquaporins in patients with multiple sclerosis. *Autoimmune Diseases* **2015** (2015) 905208.
183. Schneider, D., Liu, Y., Gerstein, M. & Engelman, D.M. Thermostability of membrane protein helix-helix interaction elucidated by statistical analysis. *FEBS Lett.* **523** (2002) 231–236.
184. Borgnia, M.J., Kozono, D., Calamita, G., Maloney, P.C. & Agre, P. Functional reconstitution and characterization of AqpZ, the *E. coli* water channel protein. *J. Molec. Biol.* **291** (1999) 1169–1179.
185. Ren, Z., Wang, Y., Duan, T., Patel, J., Liggett, T., Loda, E., Brahma, S., Goswami, R., Grouse, C., Byrne, R., Stefoski, D., Javed, A., Miller, S.D. & Balabanov, R. Cross-immunoreactivity between bacterial aquaporin-Z and human aquaporin-4: potential relevance to neuromyelitis optica. *J. Immunol.* **189** (2012) 4602–4611.
186. Tuomilehto, J. The emerging global epidemic of type 1 diabetes. *Current Diabetes Rep.* **13** (2013) 795–804.
187. Tetley, P., Simpson, S. Jr., Taylor, B.V. & van der Mei, I.A.F. The co-occurrence of multiple sclerosis and type 1 diabetes: Shared aetiologic features and clinical implication for MS aetiology. *J. Neurol. Sci.* **348** (2015) 126–131.
188. Dow, C.T. *Mycobacterium paratuberculosis* and autism: is this a trigger? *Med. Hypotheses* **77** (2011) 977–981.
189. Naser, S.A., Thanigachalam, S., Dow, S.T. & Collins, M.T. Exploring the role of *Mycobacterium avium* subspecies *paratuberculosis* in the pathogenesis of type 1 diabetes mellitus: a pilot study. *Gut Pathogens* **5** (2013) 14.
190. Capitani, G., De Biase, D., Gut, H., Ahmed, S. & Grutter, M.G. Structural model of human GAD65: prediction and interpretation of biochemical and immunogenic features. *Proteins* **59** (2005) 7–14.
191. Karjalainen, J., Martin, J.M., Knip, M., Ilonen, J., Robinson, B.H., Savilahti, E., Akerblom, H.K. & Dosch, H.M. A bovine albumin peptide as a possible trigger of insulin-dependent diabetes mellitus. *N. Engl. J. Med.* **327** (1992) 302–307.
192. Parrini, C., Taddei, N., Ramazzotti, M., Degl’Innocenti, D., Ramponi, G., Dobson, C.M. & Chiti, F. Glycine residues appear to be evolutionarily conserved for their ability to inhibit aggregation. *Structure* **13** (2005) 1143–1151.
193. Nolan, C., Margoliash, E., Peterson, J.D. & Steiner, D.F. The structure of bovine proinsulin. *J. Biol. Chem.* **246** (1971) 2780–2795.
194. Green, J.M. & Castle, L.A. Transitioning from single to multiple herbicide-resistant crops. In: *Glyphosate Resistance in Crops and Weeds: History, Development, and Management* (ed. V.K. Nandula), ch. 4, p. 112. Wiley (2010).
195. Samsel, A. & Seneff, S. Glyphosate, pathways to modern diseases II: Celiac sprue and gluten intolerance. *Interdisciplinary Toxicol.* **6** (2013) 159–184.



196. Janssen, G., Christis, C., Kooy-Winkelaar, Y., Edens, L., Smith, D., van Veelen, P. & Koning, F. Ineffective degradation of immunogenic gluten epitopes by currently available digestive enzyme supplements. *PLoS ONE* **10** (2015) e0128065.
197. Todd, J.A., Bell, J. & McDevitt, H.O. HLA-DQ beta gene contributes to susceptibility and resistance to insulin-dependent diabetes mellitus. *Nature* **329** (1987) 599–604.
198. Hogberg, L., Falth-Magnusson, K. & Grodzinsky, E. Familial prevalence of coeliac disease: a twenty-year follow-up study. *Scand. J. Gastroenterol.* **38** (2003) 61–65.
199. Li, X., Lou, Z., Li, X., Zhou, W., Ma, M., Cao, Y., Geng, Y., Bartlam, M., Zhang, X.C. & Rao, Z. Structure of human cytosolic X-prolyl aminopeptidase: a double Mn(II)-dependent dimeric enzyme with a novel three-domain subunit. *J. Biol. Chem.* **283** (2008) 22858–22866.
200. Rubio, F., Veldhuis, L.J., Clegg, S., Fleeker, J.R., & Hall, J.C. Comparison of a direct ELISA and an HPLC method for glyphosate determinations in water. *J. Agric. Food Chem.* **51** (2003) 691–696.
201. Jenkins, D.H., Grapenthien, N. & Keplinger, M.L. Milk and tissue residue study with N-phosphonomethylglycine (CP 67573) (unpublished study) prepared by Industrial Biotest Laboratories, Inc., submitted by Monsanto to US EPA, Washington, DC EPA MRID #0178 06004) (16 October 1973).